



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 07:34 pm GMT

PDB ID : 3ZS9  
Title : *S. cerevisiae* Get3-ADP-AIF4- complex with a cytosolic Get2 fragment  
Authors : Mariappan, M.; Mateja, A.; Dobosz, M.; Bove, E.; Hegde, R.S.; Keenan, R.J.  
Deposited on : 2011-06-24  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4608 (2.10-2.10)
Clashscore	122126	5109 (2.10-2.10)
Ramachandran outliers	120053	5059 (2.10-2.10)
Sidechain outliers	120020	5060 (2.10-2.10)
RSRZ outliers	108989	4497 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	<p>6% 70% 12% • 17%</p>
1	B	354	<p>4% 69% 14% • 15%</p>
2	C	38	<p>18% 71% 11% 18%</p>
2	D	38	<p>16% 84% 8% 8%</p>

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5530 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ATPASE GET3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	293	Total	C	N	O	S	0	0	0
			2310	1462	382	449	17			
1	B	302	Total	C	N	O	S	0	0	0
			2382	1512	396	457	17			

- Molecule 2 is a protein called GOLGI TO ER TRAFFIC PROTEIN 2.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	31	Total	C	N	O	0	0	0
			255	154	56	45			
2	D	35	Total	C	N	O	0	0	0
			279	167	61	51			

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	27	10	5	10	2	0	0
3	B	1	27	10	5	10	2	0	0

- Molecule 4 is TETRAFLUOROALUMINATE ION (three-letter code: ALF) (formula:  $\text{AlF}_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Al	F		
4	A	1	5	1	4	0	0
4	B	1	5	1	4	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	B	1	1	1	0	0
5	A	1	1	1	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	B	1	1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	130	Total 130	O 130	0	0
7	B	98	Total 98	O 98	0	0
7	C	3	Total 3	O 3	0	0
7	D	6	Total 6	O 6	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATPASE GET3




- Molecule 1: ATPASE GET3




- Molecule 2: GOLGI TO ER TRAFFIC PROTEIN 2



- Molecule 2: GOLGI TO ER TRAFFIC PROTEIN 2





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.56Å 77.32Å 165.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.54 – 2.10 36.54 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.9 (36.54-2.10) 96.9 (36.54-2.10)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 2.10Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.182 , 0.238 0.179 , 0.234	Depositor DCC
$R_{free}$ test set	2023 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.2	Xtrriage
Anisotropy	0.266	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 66.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5530	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ALF, ZN, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/2347	0.50	0/3164
1	B	0.37	0/2420	0.51	0/3261
2	C	0.31	0/255	0.43	0/334
2	D	0.31	0/279	0.40	0/366
All	All	0.36	0/5301	0.50	0/7125

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2310	0	2298	30	0
1	B	2382	0	2385	42	0
2	C	255	0	276	3	0
2	D	279	0	297	2	0
3	A	27	0	12	0	0
3	B	27	0	12	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	1	0	0	0	0
6	B	1	0	0	0	0
7	A	130	0	0	2	0
7	B	98	0	0	0	0
7	C	3	0	0	0	0
7	D	6	0	0	0	0
All	All	5530	0	5280	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:292:TRP:NE1	1:B:296:LYS:HD2	1.88	0.88
1:B:68:GLU:OE1	1:B:75:ARG:NH1	2.09	0.81
1:A:285:CYS:N	7:A:2095:HOH:O	2.17	0.77
1:A:150:LYS:HG3	1:A:154:GLN:NE2	2.06	0.71
1:B:250:TYR:CD2	1:B:254:ARG:NH1	2.59	0.69
1:B:229:GLN:O	1:B:236:THR:HG23	1.96	0.65
1:A:169:PRO:HG2	1:B:169:PRO:HG2	1.78	0.65
1:A:129:LEU:CD1	1:B:185:LYS:HB3	2.28	0.64
1:A:308:ASP:OD2	2:D:520:LYS:HE3	1.97	0.63
1:B:88:ILE:HD13	1:B:141:SER:HA	1.81	0.62
1:B:230:PHE:HA	1:B:236:THR:HG21	1.84	0.60
1:B:159:THR:HG22	1:B:160:PHE:H	1.66	0.60
1:A:183:LEU:O	1:A:187:LEU:HD22	2.04	0.58
1:A:213:SER:O	1:A:216:LEU:HB3	2.04	0.58
1:A:179:LEU:O	1:A:183:LEU:HB2	2.03	0.58
1:B:69:LYS:HD3	1:B:70:PHE:O	2.04	0.58
1:A:146:MET:HG2	1:A:226:ILE:HD11	1.87	0.56
1:A:278:GLU:OE2	1:A:296:LYS:NZ	2.35	0.56
1:A:181:ASN:OD1	1:A:227:ARG:NH2	2.39	0.56
1:B:278:GLU:CD	1:B:296:LYS:HZ1	2.10	0.55
1:B:292:TRP:CE2	1:B:296:LYS:HD2	2.42	0.54
1:B:250:TYR:CE2	1:B:254:ARG:NH1	2.76	0.54
1:B:126:LEU:HA	1:B:129:LEU:HD22	1.90	0.53
1:B:317:CYS:HB3	1:B:350:LEU:HD21	1.91	0.52
2:D:515:GLU:HG3	2:D:516:ARG:N	2.23	0.52
1:B:185:LYS:O	1:B:189:LYS:HD3	2.09	0.51
2:C:504:LEU:HD12	2:C:504:LEU:N	2.25	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:149:ILE:O	1:B:152:GLN:HB2	2.10	0.51
1:B:129:LEU:HD23	1:B:130:THR:N	2.26	0.51
1:B:76:LYS:HE3	1:B:81:ASN:O	2.11	0.50
1:A:130:THR:HA	1:A:133:ILE:CD1	2.42	0.49
1:B:292:TRP:CD1	1:B:296:LYS:HD2	2.46	0.49
1:B:69:LYS:HD3	1:B:70:PHE:N	2.29	0.48
1:A:329:LYS:O	1:A:332:GLN:HG2	2.13	0.48
1:B:69:LYS:HD3	1:B:69:LYS:C	2.35	0.47
1:A:129:LEU:O	1:A:133:ILE:HG13	2.14	0.47
1:B:266:VAL:HG22	1:B:309:PHE:HE2	1.79	0.47
1:A:224:GLU:O	1:A:228:GLN:HG3	2.15	0.47
1:A:129:LEU:H	1:A:129:LEU:CD2	2.28	0.47
1:A:24:GLY:HA3	1:A:238:PHE:CZ	2.50	0.46
1:A:20:TRP:HZ2	1:A:149:ILE:HD11	1.80	0.46
1:A:285:CYS:HB2	1:B:285:CYS:HB3	2.00	0.44
1:B:90:PRO:HA	1:B:140:LEU:HD13	2.00	0.44
2:C:529:ARG:O	2:C:532:LYS:HG2	2.17	0.44
1:A:130:THR:HA	1:A:133:ILE:HD12	1.99	0.44
1:A:183:LEU:HA	1:A:183:LEU:HD12	1.73	0.44
1:A:68:GLU:OE1	1:A:75:ARG:NH2	2.51	0.44
1:B:317:CYS:HA	1:B:347:ILE:HB	1.99	0.44
1:A:88:ILE:O	1:A:90:PRO:HD3	2.18	0.44
1:B:197:LEU:O	1:B:198:GLY:C	2.55	0.44
1:B:126:LEU:HD12	1:B:129:LEU:CD2	2.48	0.43
1:B:26:LYS:O	1:B:29:VAL:HG22	2.17	0.43
1:B:258:GLU:HG3	1:B:262:TYR:CZ	2.54	0.43
1:B:179:LEU:O	1:B:183:LEU:HB2	2.18	0.43
1:B:216:LEU:HA	1:B:216:LEU:HD23	1.82	0.43
1:B:176:PHE:O	1:B:179:LEU:HB2	2.19	0.43
1:A:252:THR:O	1:A:256:ILE:HG13	2.17	0.43
1:A:217:ASN:O	1:A:220:LYS:HB3	2.19	0.42
1:B:49:LYS:O	1:B:82:ASN:HB2	2.19	0.42
1:A:129:LEU:HD22	1:A:129:LEU:H	1.84	0.42
1:B:19:LYS:HG3	1:B:235:LEU:HD22	2.01	0.42
1:B:250:TYR:HE1	2:C:530:LEU:HD22	1.84	0.42
1:B:20:TRP:HB2	1:B:236:THR:HB	2.02	0.41
1:A:63:SER:HB2	7:A:2040:HOH:O	2.19	0.41
1:B:350:LEU:H	1:B:350:LEU:HD22	1.85	0.41
1:A:13:ILE:HA	1:A:42:MET:HG2	2.02	0.41
1:A:129:LEU:HD11	1:B:185:LYS:HB3	2.01	0.41
1:B:20:TRP:HD1	1:B:236:THR:HG22	1.86	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:LYS:HE2	1:A:297:LYS:HB3	1.89	0.40
1:B:56:THR:OG1	1:B:167:THR:HA	2.20	0.40
1:B:329:LYS:O	1:B:332:GLN:HG2	2.22	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/354 (80%)	277 (97%)	8 (3%)	0	100	100
1	B	292/354 (82%)	289 (99%)	3 (1%)	0	100	100
2	C	29/38 (76%)	28 (97%)	1 (3%)	0	100	100
2	D	33/38 (87%)	33 (100%)	0	0	100	100
All	All	639/784 (82%)	627 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/309 (85%)	254 (97%)	8 (3%)	43	45
1	B	270/309 (87%)	259 (96%)	11 (4%)	33	33

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	27/32 (84%)	27 (100%)	0	100	100
2	D	29/32 (91%)	29 (100%)	0	100	100
All	All	588/682 (86%)	569 (97%)	19 (3%)	42	44

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	MET
1	A	128	ASP
1	A	130	THR
1	A	159	THR
1	A	183	LEU
1	A	187	LEU
1	A	247	LEU
1	A	305	LEU
1	B	129	LEU
1	B	159	THR
1	B	179	LEU
1	B	183	LEU
1	B	212	ILE
1	B	224	GLU
1	B	236	THR
1	B	264	MET
1	B	266	VAL
1	B	299	LEU
1	B	350	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	154	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ADP	A	401	5,4	25,29,29	1.41	4 (16%)	25,45,45	1.42	2 (8%)
4	ALF	A	402	3,5	0,4,4	0.00	-	0,6,6	0.00	-
3	ADP	B	401	5,4	25,29,29	1.36	3 (12%)	25,45,45	1.40	4 (16%)
4	ALF	B	402	3,5	0,4,4	0.00	-	0,6,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	401	5,4	-	0/12/32/32	0/3/3/3
4	ALF	A	402	3,5	-	0/0/0/0	0/0/0/0
3	ADP	B	401	5,4	-	0/12/32/32	0/3/3/3
4	ALF	B	402	3,5	-	0/0/0/0	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	ADP	PB-O3A	-2.21	1.56	1.60
3	A	401	ADP	C5-C4	2.07	1.45	1.40
3	B	401	ADP	C5-C4	2.28	1.45	1.40
3	B	401	ADP	O4'-C1'	2.69	1.45	1.41
3	A	401	ADP	O4'-C1'	2.84	1.45	1.41
3	B	401	ADP	C8-N9	3.85	1.41	1.36
3	A	401	ADP	C8-N9	4.27	1.42	1.36

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	ADP	C1'-N9-C4	-3.47	120.64	126.64
3	B	401	ADP	C1'-N9-C4	-3.31	120.92	126.64
3	B	401	ADP	O3'-C3'-C4'	-2.00	105.27	111.06
3	B	401	ADP	O2B-PB-O1B	2.22	119.26	110.60
3	B	401	ADP	PA-O3A-PB	3.88	145.66	132.63
3	A	401	ADP	PA-O3A-PB	4.28	147.01	132.63

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	293/354 (82%)	0.29	23 (7%) 13 17	14, 28, 78, 102	0
1	B	302/354 (85%)	0.36	15 (4%) 29 35	15, 29, 65, 88	0
2	C	31/38 (81%)	1.22	7 (22%) 0 0	32, 45, 64, 76	0
2	D	35/38 (92%)	0.85	6 (17%) 1 1	28, 43, 58, 63	0
All	All	661/784 (84%)	0.39	51 (7%) 13 17	14, 30, 72, 102	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	129	LEU	8.1
1	A	214	GLY	7.7
1	B	127	ALA	7.3
1	A	127	ALA	6.3
1	B	283	HIS	6.0
1	B	100	MET	5.6
1	A	100	MET	5.3
1	A	186	LEU	5.3
1	A	129	LEU	5.1
1	B	101	ALA	4.9
1	A	219	LEU	4.7
1	A	215	LYS	4.4
1	B	284	ASN	4.4
2	C	524	GLY	4.4
1	B	126	LEU	4.2
1	B	97	MET	4.1
1	B	98	ASN	4.0
1	A	97	MET	3.9
1	B	99	ASP	3.8
1	A	128	ASP	3.6
2	D	523	ASN	3.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	C	523	ASN	3.4
1	A	212	ILE	3.3
1	A	187	LEU	3.2
1	B	212	ILE	3.1
1	B	128	ASP	3.1
2	C	504	LEU	3.1
2	C	525	GLY	2.9
1	A	216	LEU	2.8
2	C	507	ALA	2.8
1	A	130	THR	2.7
1	A	158	GLU	2.7
1	B	95	LYS	2.7
2	C	522	SER	2.6
1	A	98	ASN	2.6
1	A	96	ASP	2.5
1	B	29	VAL	2.5
1	A	217	ASN	2.5
1	A	241	VAL	2.5
2	D	522	SER	2.3
1	A	185	LYS	2.3
1	B	270	ILE	2.3
2	D	506	GLU	2.3
2	D	512	LEU	2.2
1	A	27	GLY	2.2
2	D	510	ARG	2.2
2	D	504	LEU	2.1
1	A	29	VAL	2.1
1	A	341	ILE	2.1
2	C	506	GLU	2.0
1	A	181	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ALF	B	402	5/5	0.95	0.22	24,26,28,31	0
5	MG	A	403	1/1	0.96	0.12	18,18,18,18	0
5	MG	B	403	1/1	0.97	0.13	17,17,17,17	0
4	ALF	A	402	5/5	0.97	0.24	19,20,32,41	0
3	ADP	B	401	27/27	0.98	0.12	12,17,28,30	0
6	ZN	B	1351	1/1	0.98	0.04	34,34,34,34	0
3	ADP	A	401	27/27	0.99	0.12	11,16,22,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.