



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2020 – 03:20 PM BST

PDB ID : 6ZUN  
Title : Crystal Structure of Thrombin in complex with compound20a  
Authors : Schafer, M.  
Deposited on : 2020-07-23  
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	36	
2	H	259	
3	I	11	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2688 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prothrombin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	27	222	140	36	45	1	0	0	0


- Molecule 2 is a protein called Prothrombin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	250	2025	1292	358	361	14	0	0	0

- Molecule 3 is a protein called Hirudin-2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	I	11	99	61	11	26	1	0	0	0

- Molecule 4 is [2-[(3-chlorophenyl)methylamino]-7-methoxy-1,3-benzoxazol-5-yl]-[(3 {R}, 4 {R})-3-methyl-4-oxidanyl-piperidin-1-yl]methanone (three-letter code: QQ5) (formula: C<sub>22</sub>H<sub>24</sub>ClN<sub>3</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
4	H	1	30	22	1	3	4	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	H	1	14	8	1	5	0	0

- Molecule 6 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	L	32	Total 32	O 32	0	0
6	H	258	Total 258	O 258	0	0
6	I	8	Total 8	O 8	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Prothrombin



- Molecule 2: Prothrombin



- Molecule 3: Hirudin-2



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.73Å 71.04Å 71.79Å 90.00° 99.52° 90.00°	Depositor
Resolution (Å)	70.80 – 1.79 33.16 – 1.79	Depositor EDS
% Data completeness (in resolution range)	87.6 (70.80-1.79) 87.7 (33.16-1.79)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.53 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.227 , 0.262 0.216 , 0.260	Depositor DCC
$R_{free}$ test set	1467 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.2	Xtrriage
Anisotropy	0.173	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 54.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2688	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: QQ5, NAG, TYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	1.57	3/224 (1.3%)	1.30	3/298 (1.0%)
2	H	1.55	14/2077 (0.7%)	1.49	30/2806 (1.1%)
3	I	0.56	0/83	0.57	0/108
All	All	1.53	17/2384 (0.7%)	1.45	33/3212 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	3

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	207	TRP	CE3-CZ3	8.14	1.52	1.38
2	H	141	TRP	CE3-CZ3	7.44	1.51	1.38
1	L	14(E)	GLU	CD-OE1	6.72	1.33	1.25
2	H	76	TYR	CE1-CZ	-6.69	1.29	1.38
1	L	1(A)	ASP	CB-CG	-6.63	1.37	1.51
2	H	217	GLU	CD-OE1	6.53	1.32	1.25
2	H	195	SER	CA-CB	6.09	1.62	1.52
2	H	45	SER	CB-OG	-6.06	1.34	1.42
2	H	51	TRP	CB-CG	-5.97	1.39	1.50
2	H	86	GLU	CD-OE2	5.94	1.32	1.25
2	H	89	TYR	CG-CD2	-5.77	1.31	1.39
2	H	77(A)	ARG	CG-CD	5.74	1.66	1.51
2	H	97	ARG	CZ-NH2	-5.53	1.25	1.33
1	L	13	GLU	CD-OE2	-5.27	1.19	1.25
2	H	101	ARG	CZ-NH2	-5.24	1.26	1.33

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	239	GLN	CD-NE2	5.22	1.46	1.32
2	H	67	ARG	CZ-NH1	5.21	1.39	1.33

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	37	SER	N-CA-C	-21.01	54.28	111.00
2	H	35	ARG	NE-CZ-NH2	13.87	127.24	120.30
2	H	35	ARG	NE-CZ-NH1	-12.06	114.27	120.30
2	H	206	ARG	NE-CZ-NH2	-9.67	115.47	120.30
2	H	187	ARG	NE-CZ-NH2	-8.98	115.81	120.30
2	H	73	ARG	NE-CZ-NH1	8.23	124.41	120.30
2	H	187	ARG	NE-CZ-NH1	8.19	124.39	120.30
2	H	97(A)	GLU	OE1-CD-OE2	-8.14	113.53	123.30
2	H	50	ARG	NE-CZ-NH1	-8.10	116.25	120.30
2	H	101	ARG	NE-CZ-NH1	8.09	124.34	120.30
2	H	101	ARG	NE-CZ-NH2	-8.05	116.28	120.30
2	H	243	ASP	CB-CG-OD1	7.82	125.34	118.30
2	H	39	GLU	OE1-CD-OE2	-7.78	113.97	123.30
2	H	49	ASP	CB-CG-OD1	7.72	125.25	118.30
1	L	1(A)	ASP	CB-CG-OD2	-7.58	111.48	118.30
2	H	126	ARG	NE-CZ-NH1	7.15	123.87	120.30
2	H	81	LYS	CD-CE-NZ	6.46	126.56	111.70
2	H	63	ASP	CB-CG-OD2	-6.28	112.65	118.30
2	H	243	ASP	CB-CG-OD2	-6.26	112.66	118.30
2	H	37	SER	CA-C-O	6.03	132.77	120.10
2	H	206	ARG	NE-CZ-NH1	5.96	123.28	120.30
2	H	49	ASP	CB-CG-OD2	-5.81	113.07	118.30
1	L	1(A)	ASP	N-CA-CB	-5.81	100.15	110.60
2	H	33	LEU	CB-CG-CD2	5.65	120.61	111.00
2	H	75	ARG	NE-CZ-NH2	-5.63	117.49	120.30
2	H	73	ARG	NE-CZ-NH2	-5.62	117.49	120.30
2	H	186(A)	ASP	CB-CG-OD1	5.60	123.34	118.30
2	H	37	SER	O-C-N	-5.60	110.46	121.10
2	H	64	LEU	CB-CG-CD2	5.49	120.34	111.00
2	H	50	ARG	CD-NE-CZ	-5.25	116.26	123.60
2	H	130	LEU	CA-CB-CG	5.19	127.24	115.30
2	H	186(A)	ASP	CB-CG-OD2	-5.07	113.74	118.30
1	L	14	ASP	CB-CG-OD1	-5.07	113.74	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	36	LYS	Peptide
2	H	37	SER	Mainchain
2	H	97(A)	GLU	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	222	0	225	0	0
2	H	2025	0	2000	18	0
3	I	99	0	77	0	0
4	H	30	0	0	0	0
5	H	14	0	13	1	0
6	H	258	0	0	2	0
6	I	8	0	0	0	0
6	L	32	0	0	0	0
All	All	2688	0	2315	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:60(H):ASN:HD22	5:H:1002:NAG:H83	1.50	0.75
2:H:139:THR:HG22	2:H:157:VAL:HG22	1.71	0.73
2:H:77(A):ARG:O	6:H:1101:HOH:O	2.15	0.65
2:H:144:LEU:HD21	2:H:152:PRO:HB3	1.85	0.59
2:H:60(I):PHE:HB3	2:H:64:LEU:HD21	1.90	0.53
2:H:60(K):THR:HG22	2:H:63:ASP:OD2	2.10	0.51
2:H:35:ARG:O	2:H:38:GLN:HA	2.12	0.48
2:H:212:ILE:HB	2:H:229:THR:HB	1.96	0.48
2:H:77(A):ARG:O	2:H:78:ASN:HB2	2.14	0.47
2:H:29:TRP:CG	2:H:121:VAL:HB	2.52	0.45
2:H:211:GLY:HA2	2:H:229:THR:O	2.17	0.45
2:H:99:LEU:HD12	2:H:215:TRP:CD1	2.52	0.45
2:H:64:LEU:CD1	2:H:85:LEU:HD12	2.48	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:68:ILE:HD12	2:H:112:VAL:HG21	1.99	0.44
2:H:201:MET:SD	2:H:210:MET:HG3	2.58	0.44
2:H:203:SER:HB3	2:H:204(B):ASN:ND2	2.34	0.42
2:H:50:ARG:NH1	6:H:1120:HOH:O	2.54	0.41
2:H:235:LYS:HE2	2:H:239:GLN:HE22	1.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	25/36 (69%)	24 (96%)	1 (4%)	0	100	100
2	H	246/259 (95%)	235 (96%)	11 (4%)	0	100	100
3	I	8/11 (73%)	8 (100%)	0	0	100	100
All	All	279/306 (91%)	267 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	25/31 (81%)	25 (100%)	0	100	100
2	H	219/225 (97%)	212 (97%)	7 (3%)	39	25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	I	9/9 (100%)	9 (100%)	0	100	100
All	All	253/265 (96%)	246 (97%)	7 (3%)	43	30

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	29	TRP
2	H	37	SER
2	H	64	LEU
2	H	75	ARG
2	H	147	THR
2	H	204(B)	ASN
2	H	239	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	78	ASN
2	H	156	GLN
2	H	204(B)	ASN
2	H	239	GLN
2	H	244	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TYS	I	18	3	15,16,17	1.94	2 (13%)	18,22,24	1.26	3 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TYS	I	18	3	-	0/10/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	18	TYS	OH-S	-5.64	1.49	1.58
3	I	18	TYS	OH-CZ	-4.25	1.35	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	18	TYS	O3-S-OH	3.83	115.04	105.83
3	I	18	TYS	CB-CA-C	-2.16	107.42	111.47
3	I	18	TYS	CG-CB-CA	-2.02	110.01	114.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	H	1002	2	14,14,15	0.92	1 (7%)	17,19,21	1.32	3 (17%)
4	QQ5	H	1001	-	29,33,33	1.48	5 (17%)	35,47,47	2.31	10 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	H	1002	2	-	4/6/23/26	0/1/1/1
4	QQ5	H	1001	-	-	0/13/28/28	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	1001	QQ5	C2-C1	-3.54	1.36	1.41
4	H	1001	QQ5	O28-C14	3.20	1.42	1.36
4	H	1001	QQ5	C8-C7	2.51	1.59	1.52
4	H	1001	QQ5	C23-CL1	2.40	1.79	1.74
5	H	1002	NAG	O5-C1	2.18	1.47	1.43
4	H	1001	QQ5	C22-C21	2.03	1.42	1.39

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1001	QQ5	C7-C8-C9	7.63	119.99	111.69
4	H	1001	QQ5	C10-C11-N6	6.64	119.53	110.17
4	H	1001	QQ5	C23-C22-C21	-3.38	115.79	119.71
4	H	1001	QQ5	C8-C7-N6	-3.03	106.21	110.82
4	H	1001	QQ5	C2-C1-C15	-2.96	118.13	121.12
5	H	1002	NAG	O4-C4-C5	2.87	116.43	109.30
4	H	1001	QQ5	C20-N19-C17	-2.86	118.29	122.23
4	H	1001	QQ5	O28-C14-C13	2.56	127.90	123.34
4	H	1001	QQ5	C29-O28-C14	-2.44	114.33	117.75
5	H	1002	NAG	C8-C7-N2	2.19	119.81	116.10
4	H	1001	QQ5	C26-C25-C24	-2.17	117.17	120.25
4	H	1001	QQ5	O5-C4-C3	-2.13	116.09	120.23

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	1002	NAG	C1-O5-C5	2.06	114.98	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	H	1002	NAG	O5-C5-C6-O6
5	H	1002	NAG	C8-C7-N2-C2
5	H	1002	NAG	O7-C7-N2-C2
5	H	1002	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	1002	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	27/36 (75%)	2.15	11 (40%) 0 0	48, 51, 53, 54	1 (3%)
2	H	250/259 (96%)	2.78	160 (64%) 0 0	47, 50, 57, 64	1 (0%)
3	I	10/11 (90%)	9.49	10 (100%) 0 0	212, 232, 277, 286	0
All	All	287/306 (93%)	2.95	181 (63%) 0 0	47, 51, 60, 286	2 (0%)

All (181) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	I	11	PHE	18.2
3	I	9	GLY	16.7
3	I	14	ILE	11.8
2	H	147	THR	11.8
3	I	19	LEU	10.7
2	H	37	SER	8.9
2	H	186(A)	ASP	8.8
3	I	10	ASP	8.2
2	H	212	ILE	8.1
2	H	138	VAL	7.2
2	H	213	VAL	7.2
2	H	53	LEU	6.8
3	I	16	GLU	6.6
2	H	245	PHE	6.5
2	H	199	PHE	6.2
2	H	29	TRP	6.1
3	I	15	PRO	6.0
2	H	31	VAL	6.0
2	H	103	ILE	5.9
3	I	12	GLU	5.9
3	I	17	GLU	5.7
2	H	242	ILE	5.7
2	H	105	LEU	5.6

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	H	44	ALA	5.5
2	H	200	VAL	5.5
2	H	244	GLN	5.4
2	H	47	ILE	5.2
1	L	1(B)	ALA	5.1
2	H	60(K)	THR	5.1
3	I	13	GLU	5.1
1	L	14(D)	ARG	5.0
2	H	198	PRO	4.9
2	H	197	GLY	4.9
2	H	41	LEU	4.8
2	H	121	VAL	4.8
2	H	141	TRP	4.6
2	H	231	VAL	4.6
2	H	181	PHE	4.5
1	L	14(J)	TYR	4.5
2	H	52	VAL	4.4
2	H	228	TYR	4.4
2	H	54	THR	4.4
2	H	128	THR	4.4
2	H	208	TYR	4.3
2	H	186(C)	GLY	4.3
2	H	158	VAL	4.2
2	H	50	ARG	4.2
1	L	14(K)	ILE	4.1
2	H	16	ILE	4.1
2	H	190	ALA	4.1
2	H	184	TYR	4.0
2	H	33	LEU	4.0
2	H	207	TRP	4.0
2	H	238	ILE	4.0
2	H	62	ASN	4.0
2	H	168	CYS	4.0
2	H	60(E)	ASP	4.0
2	H	40	LEU	3.9
2	H	211	GLY	3.9
2	H	186	PRO	3.9
2	H	176	ILE	3.8
2	H	157	VAL	3.8
2	H	191	CYS	3.8
2	H	195	SER	3.8
2	H	243	ASP	3.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	H	196	GLY	3.8
2	H	118	ILE	3.8
2	H	46	LEU	3.8
2	H	114	PHE	3.7
1	L	14(F)	LEU	3.7
2	H	241	VAL	3.7
2	H	162	ILE	3.7
2	H	155	LEU	3.7
2	H	60(D)	TRP	3.7
2	H	209	GLN	3.7
2	H	154	VAL	3.7
2	H	236	LYS	3.6
2	H	87	LYS	3.6
2	H	123	LEU	3.6
2	H	163	VAL	3.6
2	H	55	ALA	3.5
2	H	42	CYS	3.5
2	H	32	MET	3.5
2	H	240	LYS	3.5
2	H	60(A)	TYR	3.4
2	H	43	GLY	3.4
2	H	129(C)	LEU	3.4
1	L	1(A)	ASP	3.4
2	H	229	THR	3.4
2	H	38	GLN	3.4
2	H	51	TRP	3.4
2	H	113	ALA	3.3
2	H	215	TRP	3.3
2	H	99	LEU	3.3
2	H	160	LEU	3.3
2	H	37(A)	PRO	3.3
2	H	139	THR	3.3
2	H	214	SER	3.3
2	H	56	ALA	3.3
2	H	151	GLN	3.2
2	H	68	ILE	3.2
2	H	174	ILE	3.2
2	H	152	PRO	3.2
2	H	82	ILE	3.2
2	H	234	LEU	3.2
2	H	45	SER	3.2
2	H	84	MET	3.1

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	H	201	MET	3.1
2	H	161	PRO	3.1
2	H	193	GLY	3.1
2	H	140	GLY	3.1
2	H	239	GLN	3.0
2	H	129(A)	ALA	3.0
2	H	167	VAL	3.0
2	H	225	TYR	2.9
2	H	227	PHE	2.9
2	H	130	LEU	2.9
2	H	210	MET	2.9
2	H	66	VAL	2.9
2	H	126	ARG	2.9
2	H	30	GLN	2.9
2	H	34	PHE	2.8
2	H	237	TRP	2.8
2	H	88	ILE	2.8
2	H	19	GLY	2.8
2	H	64	LEU	2.8
2	H	17	VAL	2.8
2	H	183	ALA	2.8
2	H	74	THR	2.7
2	H	28	PRO	2.7
2	H	93	ARG	2.7
2	H	232	PHE	2.7
2	H	65	LEU	2.6
2	H	104	ALA	2.6
2	H	115	SER	2.6
2	H	27	SER	2.6
2	H	226	GLY	2.6
2	H	24	ILE	2.6
2	H	110	LYS	2.5
2	H	63	ASP	2.5
2	H	94	TYR	2.5
2	H	180	MET	2.5
2	H	177	THR	2.5
1	L	11	SER	2.4
2	H	182	CYS	2.4
2	H	35	ARG	2.4
1	L	12	LEU	2.4
2	H	194	ASP	2.4
2	H	122	CYS	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	H	22	ALA	2.4
2	H	108	LEU	2.4
2	H	78	ASN	2.4
2	H	111	PRO	2.4
1	L	1	CYS	2.4
2	H	70	LYS	2.4
2	H	204(A)	PHE	2.4
2	H	60(C)	PRO	2.4
2	H	165	ARG	2.4
2	H	59	LEU	2.3
2	H	166	PRO	2.3
2	H	89	TYR	2.3
2	H	125	ASP	2.3
2	H	112	VAL	2.3
2	H	90	ILE	2.3
2	H	134	TYR	2.3
2	H	188	GLY	2.3
2	H	129	ALA	2.2
2	H	172	THR	2.2
1	L	5	PRO	2.2
2	H	221	ARG	2.2
1	L	14(G)	LEU	2.2
2	H	222	ASP	2.2
2	H	25	GLY	2.1
2	H	49	ASP	2.1
2	H	76	TYR	2.1
2	H	96	TRP	2.1
2	H	79	ILE	2.1
2	H	20	SER	2.0
2	H	170	ASP	2.0
2	H	60(B)	PRO	2.0
2	H	204	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	TYS	I	18	16/17	0.36	0.37	245,260,284,294	0

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	H	1002	14/15	0.67	0.35	49,65,75,77	0
4	QQ5	H	1001	30/30	0.84	0.18	47,48,49,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers

There are no such residues in this entry.