



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 7, 2022 – 01:25 pm BST

PDB ID : 7ZVN
Title : Crystal structure of human Annexin A2 in complex with full phosphorothioate 5-10 2'-methoxyethyl DNA gapmer antisense oligonucleotide solved at 1.87 Å resolution
Authors : Hyjek-Skladanowska, M.; Anderson, B.; Mykhaylyk, V.; Orr, C.; Wagner, A.; Skowronek, K.; Seth, P.; Nowotny, M.
Deposited on : 2022-05-16
Resolution : 1.87 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.30
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.30


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	 91% 9%
2	C	15	 27% 33% 7% 33%
2	D	15	 47% 47% 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Annexin A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	307	2472	1548	425	488	11	0	5	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	SER	-	expression tag	UNP P07355
A	293	ALA	VAL	variant	UNP P07355

- Molecule 2 is a DNA chain called 2'-methoxyethyl DNA gapmer antisense oligonucleotide.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
2	C	10	365	178	66	91	15	15	0	6	0
2	D	15	523	257	88	134	22	22	0	8	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	7	Total Ca 7 7	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	320	Total O 320 320	0	0
6	C	12	Total O 12 12	0	0
6	D	20	Total O 20 20	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Annexin A2



- Molecule 2: 2'-methoxyethyl DNA gapmer antisense oligonucleotide



- Molecule 2: 2'-methoxyethyl DNA gapmer antisense oligonucleotide



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.62Å 57.50Å 70.48Å 90.00° 90.24° 90.00°	Depositor
Resolution (Å)	43.75 – 1.87 43.75 – 1.87	Depositor EDS
% Data completeness (in resolution range)	99.8 (43.75-1.87) 99.8 (43.75-1.87)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.95 (at 1.87Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.167 , 0.198 0.166 , 0.198	Depositor DCC
R_{free} test set	2011 reflections (5.48%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtrriage
Anisotropy	0.179	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.006 for -k,-h,-l 0.011 for k,h,-l 0.032 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3768	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K39, N7X, AS, GS, MG, PST, GOL, OKN, K2F, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/2503	0.66	0/3363

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2472	0	2458	18	0
2	C	365	0	96	1	0
2	D	523	0	142	1	0
3	A	48	0	64	2	0
4	A	7	0	0	0	0
5	A	1	0	0	0	0
6	A	320	0	0	5	0
6	C	12	0	0	0	0
6	D	20	0	0	1	0
All	All	3768	0	2760	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:318:TYR:OH	6:A:501:HOH:O	2.10	0.69
1:A:230[B]:ASP:OD2	6:A:502:HOH:O	2.10	0.68
1:A:168:ARG:HD2	6:A:697:HOH:O	2.01	0.60
1:A:115:LYS:NZ	6:A:505:HOH:O	2.26	0.59
1:A:284:ARG:NH1	6:A:510:HOH:O	2.36	0.58
2:C:6[A]:GS:S2P	2:C:6[A]:GS:N2	2.68	0.54
1:A:145:ARG:NH1	1:A:149:GLU:OE1	2.41	0.53
1:A:281:LYS:O	3:A:405:GOL:H2	2.10	0.52
1:A:237:PRO:HG2	1:A:238:TYR:CE2	2.46	0.51
1:A:284:ARG:HG3	3:A:405:GOL:H31	1.92	0.50
1:A:129:ILE:O	1:A:133:CYS:HB2	2.14	0.48
1:A:148:LYS:HD3	1:A:154:ASP:OD1	2.15	0.47
2:D:4[B]:K2F:SP1	6:D:102:HOH:O	2.59	0.46
1:A:73:PHE:HE2	1:A:77:ARG:HH21	1.64	0.44
1:A:87:LEU:HD12	1:A:103:LEU:HD11	2.01	0.43
1:A:43:GLU:OE1	1:A:78:ARG:HD3	2.19	0.42
1:A:238:TYR:CE1	1:A:246:LYS:HE3	2.55	0.42
1:A:36:GLU:OE2	1:A:77:ARG:NH2	2.51	0.41
1:A:292:MET:O	1:A:296[B]:SER:HB2	2.21	0.41
1:A:47:LYS:HA	1:A:47:LYS:HD3	1.81	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	310/307 (101%)	304 (98%)	5 (2%)	1 (0%)	41 30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	51	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	264/269 (98%)	262 (99%)	2 (1%)	81 80

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	162	ASP
1	A	322	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

39 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	OKN	D	7[A]	2	17,21,22	0.21	0	24,30,33	0.45	0
2	GS	C	10[A]	2	18,24,25	1.02	2 (11%)	19,35,38	0.55	0
2	K39	C	1	2	21,26,30	0.97	2 (9%)	23,37,44	0.69	1 (4%)
2	PST	D	14	2	18,21,22	0.22	0	26,30,33	0.55	0
2	GS	C	6[A]	2	18,24,25	1.02	2 (11%)	19,35,38	0.59	0
2	K2F	D	4[A]	2	21,28,29	0.67	0	22,39,42	0.73	1 (4%)
2	GS	D	9	2	18,24,25	1.03	2 (11%)	19,35,38	0.56	0
2	GS	D	10[B]	2	18,24,25	1.01	2 (11%)	19,35,38	0.67	0
2	GS	D	15	2	18,24,25	1.07	2 (11%)	19,35,38	0.64	0
2	GS	C	10[B]	2	18,24,25	1.02	2 (11%)	19,35,38	0.56	0
2	OKN	D	7[B]	2	17,21,22	0.21	0	24,30,33	0.47	0
2	K39	C	5	2	21,29,30	0.97	2 (9%)	23,41,44	0.61	1 (4%)
2	GS	C	6[B]	2	18,24,25	1.01	2 (11%)	19,35,38	0.56	0
2	N7X	C	3	2	22,26,27	0.28	0	30,36,39	0.66	0
2	K2F	D	4[B]	2	21,28,29	0.66	0	22,39,42	0.72	1 (4%)
2	K39	D	5[A]	2	21,29,30	0.95	2 (9%)	23,41,44	0.64	1 (4%)
2	GS	C	9[A]	2	18,24,25	1.03	2 (11%)	19,35,38	0.55	0
2	N7X	D	3[A]	2	22,26,27	0.25	0	30,36,39	0.48	0
2	PST	D	11	2	18,21,22	0.19	0	26,30,33	0.34	0
2	PST	D	12[A]	2	18,21,22	0.19	0	26,30,33	0.27	0
2	K39	D	5[B]	2	21,29,30	0.96	2 (9%)	23,41,44	0.62	1 (4%)
2	GS	C	9[B]	2	18,24,25	1.03	2 (11%)	19,35,38	0.56	0
2	K39	D	1	2	21,26,30	0.95	2 (9%)	23,37,44	0.61	1 (4%)
2	GS	D	6[A]	2	18,24,25	1.02	2 (11%)	19,35,38	0.53	0
2	N7X	C	2	2	22,26,27	0.27	0	30,36,39	0.43	0
2	N7X	D	3[B]	2	22,26,27	0.26	0	30,36,39	0.57	0
2	PST	C	8[A]	2	18,21,22	0.18	0	26,30,33	0.28	0
2	PST	D	12[B]	2	18,21,22	0.18	0	26,30,33	0.22	0
2	K2F	C	4[A]	2	21,28,29	0.59	0	22,39,42	0.66	1 (4%)
2	GS	D	6[B]	2	18,24,25	1.02	2 (11%)	19,35,38	0.57	0
2	PST	C	8[B]	2	18,21,22	0.20	0	26,30,33	0.33	0
2	OKN	C	7[A]	2	17,21,22	0.22	0	24,30,33	0.50	0
2	N7X	D	2[A]	2	22,26,27	0.24	0	30,36,39	0.38	0
2	K2F	C	4[B]	2	21,28,29	0.60	0	22,39,42	0.68	1 (4%)
2	PST	D	8	2	18,21,22	0.27	0	26,30,33	0.31	0
2	OKN	C	7[B]	2	17,21,22	0.24	0	24,30,33	0.49	0
2	N7X	D	2[B]	2	22,26,27	0.22	0	30,36,39	0.35	0
2	AS	D	13	2	18,23,24	0.62	0	17,33,36	0.77	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GS	D	10[A]	2	18,24,25	1.01	2 (11%)	19,35,38	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OKN	D	7[A]	2	-	0/6/21/22	0/2/2/2
2	GS	C	10[A]	2	-	0/2/21/22	0/3/3/3
2	K39	C	1	2	-	0/7/27/31	0/3/3/3
2	PST	D	14	2	-	2/7/21/22	0/2/2/2
2	GS	C	6[A]	2	-	0/2/21/22	0/3/3/3
2	K2F	D	4[A]	2	-	3/8/30/31	0/3/3/3
2	GS	D	9	2	-	0/2/21/22	0/3/3/3
2	GS	D	10[B]	2	-	0/2/21/22	0/3/3/3
2	GS	D	15	2	-	0/2/21/22	0/3/3/3
2	GS	C	10[B]	2	-	0/2/21/22	0/3/3/3
2	OKN	D	7[B]	2	-	0/6/21/22	0/2/2/2
2	K39	C	5	2	-	2/8/30/31	0/3/3/3
2	GS	C	6[B]	2	-	2/2/21/22	0/3/3/3
2	N7X	C	3	2	-	1/12/30/31	0/2/2/2
2	K2F	D	4[B]	2	-	4/8/30/31	0/3/3/3
2	K39	D	5[A]	2	-	5/8/30/31	0/3/3/3
2	GS	C	9[A]	2	-	0/2/21/22	0/3/3/3
2	N7X	D	3[A]	2	-	4/12/30/31	0/2/2/2
2	PST	D	11	2	-	0/7/21/22	0/2/2/2
2	PST	D	12[A]	2	-	0/7/21/22	0/2/2/2
2	K39	D	5[B]	2	-	4/8/30/31	0/3/3/3
2	GS	C	9[B]	2	-	2/2/21/22	0/3/3/3
2	K39	D	1	2	-	4/7/27/31	0/3/3/3
2	GS	D	6[A]	2	-	1/2/21/22	0/3/3/3
2	N7X	C	2	2	-	1/12/30/31	0/2/2/2
2	N7X	D	3[B]	2	-	3/12/30/31	0/2/2/2
2	PST	C	8[A]	2	-	0/7/21/22	0/2/2/2
2	PST	D	12[B]	2	-	0/7/21/22	0/2/2/2
2	K2F	C	4[A]	2	-	3/8/30/31	0/3/3/3
2	GS	D	6[B]	2	-	2/2/21/22	0/3/3/3
2	PST	C	8[B]	2	-	0/7/21/22	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OKN	C	7[A]	2	-	0/6/21/22	0/2/2/2
2	N7X	D	2[A]	2	-	3/12/30/31	0/2/2/2
2	K2F	C	4[B]	2	-	3/8/30/31	0/3/3/3
2	PST	D	8	2	-	0/7/21/22	0/2/2/2
2	OKN	C	7[B]	2	-	0/6/21/22	0/2/2/2
2	N7X	D	2[B]	2	-	3/12/30/31	0/2/2/2
2	AS	D	13	2	-	0/3/21/22	0/3/3/3
2	GS	D	10[A]	2	-	0/2/21/22	0/3/3/3

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	K39	C5-C6	-2.60	1.42	1.47
2	C	9[B]	GS	C5-C6	-2.58	1.42	1.47
2	D	6[B]	GS	C5-C6	-2.58	1.42	1.47
2	C	9[A]	GS	C5-C6	-2.58	1.42	1.47
2	D	9	GS	C5-C6	-2.56	1.42	1.47
2	C	10[A]	GS	C5-C6	-2.56	1.42	1.47
2	C	10[B]	GS	C5-C6	-2.55	1.42	1.47
2	D	15	GS	C5-C6	-2.55	1.42	1.47
2	D	6[A]	GS	C5-C6	-2.54	1.42	1.47
2	C	6[A]	GS	C5-C6	-2.53	1.42	1.47
2	D	10[A]	GS	C5-C6	-2.51	1.42	1.47
2	C	6[B]	GS	C5-C6	-2.50	1.42	1.47
2	D	5[B]	K39	C5-C6	-2.50	1.42	1.47
2	D	5[A]	K39	C5-C6	-2.49	1.42	1.47
2	D	10[B]	GS	C5-C6	-2.49	1.42	1.47
2	D	1	K39	C5-C6	-2.46	1.42	1.47
2	C	5	K39	C5-C6	-2.44	1.42	1.47
2	C	1	K39	C8-N7	-2.41	1.30	1.35
2	D	5[A]	K39	C8-N7	-2.38	1.31	1.35
2	D	6[B]	GS	C8-N7	-2.38	1.31	1.35
2	D	5[B]	K39	C8-N7	-2.38	1.31	1.35
2	C	9[B]	GS	C8-N7	-2.37	1.31	1.35
2	C	6[A]	GS	C8-N7	-2.37	1.31	1.35
2	C	9[A]	GS	C8-N7	-2.37	1.31	1.35
2	D	15	GS	C8-N7	-2.37	1.31	1.35
2	D	1	K39	C8-N7	-2.36	1.31	1.35
2	C	10[A]	GS	C8-N7	-2.36	1.31	1.35
2	C	10[B]	GS	C8-N7	-2.36	1.31	1.35
2	D	10[A]	GS	C8-N7	-2.36	1.31	1.35
2	D	6[A]	GS	C8-N7	-2.35	1.31	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	10[B]	GS	C8-N7	-2.34	1.31	1.35
2	C	5	K39	C8-N7	-2.34	1.31	1.35
2	C	6[B]	GS	C8-N7	-2.33	1.31	1.35
2	D	9	GS	C8-N7	-2.32	1.31	1.35

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4[A]	K2F	C5-C6-N6	2.24	123.75	120.35
2	C	4[B]	K2F	C5-C6-N6	2.23	123.74	120.35
2	D	4[A]	K2F	C5-C6-N6	2.20	123.70	120.35
2	D	4[B]	K2F	C5-C6-N6	2.20	123.69	120.35
2	D	13	AS	C5-C6-N6	2.17	123.65	120.35
2	C	5	K39	O6-C6-C5	2.17	128.61	124.37
2	D	5[A]	K39	O6-C6-C5	2.15	128.57	124.37
2	D	1	K39	O6-C6-C5	2.15	128.57	124.37
2	D	5[B]	K39	O6-C6-C5	2.15	128.57	124.37
2	C	1	K39	O6-C6-C5	2.11	128.50	124.37

There are no chirality outliers.

All (52) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	2[A]	N7X	C1'-C2'-O2'-C
2	D	2[B]	N7X	C1'-C2'-O2'-C
2	D	3[A]	N7X	C1'-C2'-O2'-C
2	D	3[B]	N7X	C1'-C2'-O2'-C
2	D	6[B]	GS	C3'-C4'-C5'-O5'
2	C	9[B]	GS	O4'-C4'-C5'-O5'
2	C	9[B]	GS	C3'-C4'-C5'-O5'
2	D	1	K39	C3'-C4'-C5'-O5'
2	D	1	K39	O4'-C4'-C5'-O5'
2	D	5[A]	K39	C3'-C4'-C5'-O5'
2	D	5[A]	K39	O2'-CA'-CB'-OC'
2	D	5[B]	K39	O2'-CA'-CB'-OC'
2	D	3[A]	N7X	O2'-C-C1-O12
2	D	2[A]	N7X	O2'-C-C1-O12
2	D	3[B]	N7X	O2'-C-C1-O12
2	C	6[B]	GS	O4'-C4'-C5'-O5'
2	D	2[A]	N7X	C-C1-O12-C12
2	D	4[A]	K2F	O2'-CA'-CB'-OC'
2	D	4[B]	K2F	O2'-CA'-CB'-OC'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	D	5[B]	K39	CA'-CB'-OC'-CD'
2	D	14	PST	O4'-C4'-C5'-O5'
2	D	14	PST	C3'-C4'-C5'-O5'
2	C	5	K39	O2'-CA'-CB'-OC'
2	D	1	K39	O2'-CA'-CB'-OC'
2	D	5[A]	K39	CA'-CB'-OC'-CD'
2	C	4[A]	K2F	C3'-C4'-C5'-O5'
2	C	6[B]	GS	C3'-C4'-C5'-O5'
2	D	6[B]	GS	O4'-C4'-C5'-O5'
2	C	4[A]	K2F	C1'-C2'-O2'-CA'
2	C	4[B]	K2F	C1'-C2'-O2'-CA'
2	C	4[B]	K2F	C3'-C4'-C5'-O5'
2	D	5[A]	K39	O4'-C4'-C5'-O5'
2	C	2	N7X	C-C1-O12-C12
2	D	2[B]	N7X	O2'-C-C1-O12
2	D	6[A]	GS	C3'-C4'-C5'-O5'
2	D	3[B]	N7X	C1-C-O2'-C2'
2	D	5[A]	K39	CB'-CA'-O2'-C2'
2	D	5[B]	K39	CB'-CA'-O2'-C2'
2	D	5[B]	K39	C3'-C4'-C5'-O5'
2	D	2[B]	N7X	C-C1-O12-C12
2	C	4[B]	K2F	CA'-CB'-OC'-CD'
2	D	3[A]	N7X	C1-C-O2'-C2'
2	D	4[A]	K2F	CB'-CA'-O2'-C2'
2	D	1	K39	CA'-CB'-OC'-CD'
2	C	4[A]	K2F	CA'-CB'-OC'-CD'
2	D	4[B]	K2F	CB'-CA'-O2'-C2'
2	C	3	N7X	C-C1-O12-C12
2	D	4[A]	K2F	C1'-C2'-O2'-CA'
2	D	4[B]	K2F	C1'-C2'-O2'-CA'
2	D	3[A]	N7X	C-C1-O12-C12
2	D	4[B]	K2F	C3'-C2'-O2'-CA'
2	C	5	K39	CA'-CB'-OC'-CD'

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	6[A]	GS	1	0
2	D	4[B]	K2F	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	408	-	5,5,5	0.84	0	5,5,5	0.87	0
3	GOL	A	407	-	5,5,5	0.60	0	5,5,5	1.21	1 (20%)
3	GOL	A	404	-	5,5,5	0.62	0	5,5,5	1.39	0
3	GOL	A	405	-	5,5,5	0.99	0	5,5,5	0.88	0
3	GOL	A	403	-	5,5,5	0.93	0	5,5,5	1.00	0
3	GOL	A	401	-	5,5,5	0.94	0	5,5,5	1.14	1 (20%)
3	GOL	A	402	-	5,5,5	0.95	0	5,5,5	0.83	0
3	GOL	A	406	-	5,5,5	0.77	0	5,5,5	1.23	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	408	-	-	0/4/4/4	-
3	GOL	A	407	-	-	4/4/4/4	-
3	GOL	A	404	-	-	4/4/4/4	-
3	GOL	A	405	-	-	4/4/4/4	-
3	GOL	A	403	-	-	2/4/4/4	-
3	GOL	A	401	-	-	2/4/4/4	-
3	GOL	A	402	-	-	2/4/4/4	-
3	GOL	A	406	-	-	2/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	406	GOL	C3-C2-C1	-2.37	102.47	111.70
3	A	401	GOL	C3-C2-C1	-2.02	103.86	111.70
3	A	407	GOL	C3-C2-C1	-2.01	103.89	111.70

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	GOL	C1-C2-C3-O3
3	A	403	GOL	C1-C2-C3-O3
3	A	403	GOL	O2-C2-C3-O3
3	A	404	GOL	O1-C1-C2-C3
3	A	404	GOL	C1-C2-C3-O3
3	A	405	GOL	C1-C2-C3-O3
3	A	406	GOL	O1-C1-C2-C3
3	A	407	GOL	O1-C1-C2-C3
3	A	402	GOL	O2-C2-C3-O3
3	A	406	GOL	O1-C1-C2-O2
3	A	401	GOL	C1-C2-C3-O3
3	A	405	GOL	O1-C1-C2-C3
3	A	407	GOL	C1-C2-C3-O3
3	A	404	GOL	O1-C1-C2-O2
3	A	405	GOL	O2-C2-C3-O3
3	A	407	GOL	O1-C1-C2-O2
3	A	404	GOL	O2-C2-C3-O3
3	A	405	GOL	O1-C1-C2-O2
3	A	407	GOL	O2-C2-C3-O3
3	A	401	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	405	GOL	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	307/307 (100%)	-0.16	1 (0%) 94 94	18, 26, 46, 76	0
2	C	0/15	-	-	-	-
2	D	0/15	-	-	-	-
All	All	307/337 (91%)	-0.16	1 (0%) 94 94	18, 26, 46, 76	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	339	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GS	C	10[A]	22/23	0.73	0.23	70,79,97,103	22
2	GS	C	10[B]	22/23	0.73	0.23	70,81,97,106	22
2	GS	C	9[A]	22/23	0.81	0.21	66,72,90,96	22
2	GS	C	9[B]	22/23	0.81	0.21	66,72,89,100	22
2	GS	D	6[A]	22/23	0.82	0.20	50,61,77,87	22
2	GS	D	6[B]	22/23	0.82	0.20	53,62,78,88	22
2	K39	D	5[A]	27/28	0.85	0.16	56,71,80,83	27
2	K39	D	5[B]	27/28	0.85	0.16	56,71,79,83	27
2	OKN	D	7[B]	20/21	0.86	0.18	45,56,80,82	20
2	N7X	D	2[A]	25/26	0.86	0.18	53,66,79,89	25
2	N7X	D	2[B]	25/26	0.86	0.18	46,66,79,89	25
2	K39	D	1	24/28	0.86	0.28	72,79,85,87	0
2	OKN	D	7[A]	20/21	0.86	0.18	45,56,82,83	20

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	N7X	D	3[A]	25/26	0.88	0.16	55,64,75,80	25
2	N7X	D	3[B]	25/26	0.88	0.16	55,64,73,80	25
2	PST	C	8[A]	20/21	0.88	0.16	57,67,79,89	20
2	PST	C	8[B]	20/21	0.88	0.16	57,67,78,88	20
2	GS	C	6[A]	22/23	0.88	0.19	45,52,62,64	22
2	GS	C	6[B]	22/23	0.88	0.19	47,53,63,65	22
2	K2F	D	4[A]	26/27	0.88	0.18	55,66,76,83	26
2	K2F	D	4[B]	26/27	0.88	0.18	58,66,76,83	26
2	OKN	C	7[A]	20/21	0.90	0.18	51,57,71,77	20
2	OKN	C	7[B]	20/21	0.90	0.18	51,57,69,78	20
2	GS	D	9	22/23	0.93	0.12	32,40,53,54	0
2	K2F	C	4[B]	26/27	0.94	0.13	32,43,50,57	26
2	K39	C	5	27/28	0.94	0.12	41,49,56,69	0
2	PST	D	8	20/21	0.94	0.13	37,49,79,83	0
2	K2F	C	4[A]	26/27	0.94	0.13	32,43,50,57	26
2	PST	D	11	20/21	0.94	0.13	29,38,44,55	0
2	K39	C	1	24/28	0.96	0.09	22,28,36,46	0
2	GS	D	10[A]	22/23	0.96	0.08	30,34,42,47	22
2	GS	D	10[B]	22/23	0.96	0.08	30,34,42,50	22
2	N7X	C	3	25/26	0.96	0.10	27,37,47,60	0
2	PST	D	12[A]	20/21	0.96	0.09	29,32,40,40	20
2	PST	D	12[B]	20/21	0.96	0.09	28,32,38,38	20
2	GS	D	15	22/23	0.96	0.09	24,28,35,47	0
2	AS	D	13	21/22	0.97	0.08	22,28,33,35	0
2	PST	D	14	20/21	0.97	0.09	27,33,41,44	0
2	N7X	C	2	25/26	0.97	0.08	22,30,40,63	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	403	6/6	0.79	0.24	28,48,54,55	0
3	GOL	A	404	6/6	0.86	0.29	31,40,57,72	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	405	6/6	0.88	0.17	28,42,54,58	0
3	GOL	A	408	6/6	0.88	0.27	39,55,60,60	0
5	MG	A	416	1/1	0.91	0.06	40,40,40,40	0
3	GOL	A	407	6/6	0.93	0.35	37,50,58,59	0
3	GOL	A	406	6/6	0.94	0.14	32,42,48,51	0
4	CA	A	409	1/1	0.95	0.07	42,42,42,42	0
3	GOL	A	401	6/6	0.95	0.10	26,27,32,35	0
4	CA	A	415	1/1	0.96	0.07	32,32,32,32	1
3	GOL	A	402	6/6	0.97	0.08	26,40,46,58	0
4	CA	A	414	1/1	0.97	0.06	30,30,30,30	0
4	CA	A	412	1/1	0.98	0.05	42,42,42,42	0
4	CA	A	410	1/1	0.99	0.07	25,25,25,25	0
4	CA	A	411	1/1	0.99	0.11	24,24,24,24	0
4	CA	A	413	1/1	1.00	0.13	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.