



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 6, 2024 – 06:55 AM EDT

PDB ID : 1ZWY
Title : Crystal structure of protein VC0702 from *Vibrio cholerae*
Authors : Chang, C.; Wu, R.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2005-06-06
Resolution : 1.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	185	 5% 77% 11% • 11%
1	B	185	 % 83% 12% • •
1	C	185	 3% 79% 11% • 7%
1	D	185	 77% 14% • 6%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 6062 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical UPF0244 protein VC0702.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	165	1313	829	231	247	1	5	0	4	0
1	B	180	1431	907	254	263	1	6	0	3	0
1	C	172	1377	871	244	256	1	5	0	4	0
1	D	173	1374	868	244	256	1	5	0	2	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ASN	-	cloning artifact	UNP Q9KU27
A	0	ALA	-	cloning artifact	UNP Q9KU27
A	1	MSE	MET	modified residue	UNP Q9KU27
A	10	MSE	MET	modified residue	UNP Q9KU27
A	55	MSE	MET	modified residue	UNP Q9KU27
A	97	MSE	MET	modified residue	UNP Q9KU27
A	114	MSE	MET	modified residue	UNP Q9KU27
A	133	MSE	MET	modified residue	UNP Q9KU27
B	-1	ASN	-	cloning artifact	UNP Q9KU27
B	0	ALA	-	cloning artifact	UNP Q9KU27
B	1	MSE	MET	modified residue	UNP Q9KU27
B	10	MSE	MET	modified residue	UNP Q9KU27
B	55	MSE	MET	modified residue	UNP Q9KU27
B	97	MSE	MET	modified residue	UNP Q9KU27
B	114	MSE	MET	modified residue	UNP Q9KU27
B	133	MSE	MET	modified residue	UNP Q9KU27
C	-1	ASN	-	cloning artifact	UNP Q9KU27
C	0	ALA	-	cloning artifact	UNP Q9KU27
C	1	MSE	MET	modified residue	UNP Q9KU27
C	10	MSE	MET	modified residue	UNP Q9KU27
C	55	MSE	MET	modified residue	UNP Q9KU27

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	97	MSE	MET	modified residue	UNP Q9KU27
C	114	MSE	MET	modified residue	UNP Q9KU27
C	133	MSE	MET	modified residue	UNP Q9KU27
D	-1	ASN	-	cloning artifact	UNP Q9KU27
D	0	ALA	-	cloning artifact	UNP Q9KU27
D	1	MSE	MET	modified residue	UNP Q9KU27
D	10	MSE	MET	modified residue	UNP Q9KU27
D	55	MSE	MET	modified residue	UNP Q9KU27
D	97	MSE	MET	modified residue	UNP Q9KU27
D	114	MSE	MET	modified residue	UNP Q9KU27
D	133	MSE	MET	modified residue	UNP Q9KU27


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	100	Total O 100 100	0	0
2	B	168	Total O 168 168	0	0
2	C	122	Total O 122 122	0	0
2	D	177	Total O 177 177	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hypothetical UPF0244 protein VC0702




- Molecule 1: Hypothetical UPF0244 protein VC0702



- Molecule 1: Hypothetical UPF0244 protein VC0702



- Molecule 1: Hypothetical UPF0244 protein VC0702



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.40Å 105.35Å 75.27Å 90.00° 104.57° 90.00°	Depositor
Resolution (Å)	50.00 – 1.90 50.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.4 (50.00-1.90) 98.4 (50.00-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 1.88Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.180 , 0.226 0.195 , 0.235	Depositor DCC
R_{free} test set	3176 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtrriage
Anisotropy	0.100	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6062	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1345	0.69	1/1811 (0.1%)
1	B	0.81	0/1464	0.80	2/1972 (0.1%)
1	C	0.68	0/1412	0.69	1/1901 (0.1%)
1	D	0.85	0/1404	0.76	0/1894
All	All	0.75	0/5625	0.74	4/7578 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	11	ARG	NE-CZ-NH2	-7.99	116.30	120.30
1	A	181	PRO	N-CA-CB	7.44	112.23	103.30
1	B	97	MSE	CG-SE-CE	-6.55	84.50	98.90
1	C	105	ARG	NE-CZ-NH2	-6.42	117.09	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1313	0	1300	14	0
1	B	1431	0	1443	22	0
1	C	1377	0	1379	20	0
1	D	1374	0	1373	30	0
2	A	100	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	168	0	0	2	0
2	C	122	0	0	3	0
2	D	177	0	0	3	0
All	All	6062	0	5495	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:84[B]:LEU:HG	1:B:97:MSE:HE1	1.18	1.14
1:B:84[B]:LEU:HG	1:B:97:MSE:CE	1.83	1.09
1:A:133:MSE:SE	1:A:137:PHE:HE2	2.00	0.94
1:B:84[B]:LEU:CG	1:B:97:MSE:HE1	1.97	0.94
1:B:155[B]:HIS:HB3	1:C:155[B]:HIS:HB3	1.47	0.93
2:C:208:HOH:O	1:D:92:LYS:HE2	1.68	0.91
1:D:11:ARG:HG3	1:D:39:TRP:CZ3	2.12	0.85
1:C:104[A]:GLN:HE21	1:C:105:ARG:H	1.28	0.81
1:D:104[B]:GLN:HE21	1:D:105:ARG:H	1.26	0.80
1:B:144:GLN:HB3	2:B:331:HOH:O	1.84	0.77
1:C:159:ARG:NH1	2:C:192:HOH:O	2.18	0.76
1:D:53:GLN:HE22	1:D:129:LEU:H	1.35	0.74
1:B:24:ASN:ND2	1:B:27:ARG:HH21	1.86	0.72
1:C:33[A]:VAL:HG21	1:C:171:LEU:HD13	1.69	0.72
2:C:208:HOH:O	1:D:92:LYS:CE	2.32	0.69
1:A:115:LEU:HD22	1:A:133:MSE:HE1	1.75	0.69
1:B:67:ARG:NH1	1:B:67:ARG:HG2	2.07	0.68
1:D:139:THR:HG22	1:D:143:LYS:HE2	1.77	0.67
1:B:91:ASN:HD21	1:B:114:MSE:HE2	1.58	0.66
1:C:9:VAL:O	1:C:10:MSE:HB2	1.96	0.66
1:A:133:MSE:SE	1:A:137:PHE:CE2	2.91	0.65
1:D:122:ARG:HD2	1:D:125:GLN:NE2	2.11	0.64
1:A:115:LEU:HD22	1:A:133:MSE:CE	2.28	0.64
1:C:91:ASN:HB2	1:C:120:LEU:HD11	1.79	0.63
1:D:53:GLN:NE2	1:D:129:LEU:H	1.98	0.60
1:D:11:ARG:CG	1:D:39:TRP:CZ3	2.84	0.60
1:C:145:LYS:HD3	1:C:151:LEU:HD11	1.83	0.60
1:C:92[B]:LYS:HE2	1:C:114:MSE:HG3	1.85	0.58
1:B:67:ARG:HG2	1:B:67:ARG:HH11	1.69	0.58
1:B:46:VAL:O	1:B:67:ARG:NH1	2.34	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:12:LYS:HD3	1:C:42:ILE:CD1	2.33	0.58
1:D:139:THR:HG22	1:D:143:LYS:CE	2.33	0.58
1:D:158:THR:HG22	1:D:161:THR:H	1.69	0.58
1:D:102:ASP:OD1	1:D:103:GLN:HG2	2.02	0.58
1:D:158:THR:CG2	1:D:160:SER:OG	2.52	0.57
1:A:178:GLU:CD	1:A:178:GLU:H	2.08	0.57
1:C:67:ARG:HE	1:C:85:GLU:CD	2.08	0.57
1:B:84[B]:LEU:CD2	1:B:97:MSE:HE1	2.35	0.56
1:D:122:ARG:HD2	1:D:125:GLN:HE21	1.70	0.54
1:D:11:ARG:HG2	1:D:39:TRP:CE3	2.44	0.54
1:A:104:GLN:HB3	1:A:174:PHE:CE1	2.43	0.53
1:B:24:ASN:HD21	1:B:27:ARG:HH21	1.56	0.53
1:D:60:THR:OG1	1:D:87:GLY:HA3	2.09	0.53
1:D:75:HIS:HE1	2:D:211:HOH:O	1.92	0.53
1:B:155[B]:HIS:CB	1:C:155[B]:HIS:HB3	2.31	0.52
1:B:154:ARG:O	1:B:155[B]:HIS:HB2	2.10	0.51
1:D:92:LYS:HE3	1:D:114:MSE:HG3	1.93	0.51
1:A:154:ARG:HG3	1:B:175:ILE:CG2	2.41	0.49
1:C:24:ASN:HD22	1:C:27:ARG:NH1	2.09	0.49
1:D:122:ARG:HH11	1:D:125:GLN:HE22	1.60	0.49
1:B:10:MSE:SE	2:B:269:HOH:O	2.80	0.49
1:A:119:VAL:HG11	1:A:133:MSE:HE2	1.95	0.49
1:D:158:THR:HG21	1:D:160:SER:OG	2.13	0.48
1:C:104[A]:GLN:NE2	1:C:105:ARG:H	2.04	0.48
1:C:154:ARG:NH1	1:D:175:ILE:O	2.47	0.47
1:D:85:GLU:OE1	2:D:254:HOH:O	2.20	0.47
1:A:11:ARG:HG3	1:A:79:GLU:HB2	1.96	0.47
1:D:122:ARG:HH11	1:D:125:GLN:NE2	2.11	0.47
1:D:11:ARG:CG	1:D:39:TRP:CE3	2.97	0.47
1:D:24:ASN:ND2	1:D:27:ARG:HH11	2.12	0.46
1:C:104[A]:GLN:HE21	1:C:105:ARG:N	2.06	0.46
1:B:60:THR:OG1	1:B:87:GLY:HA3	2.16	0.45
1:B:92:LYS:HE3	1:B:114:MSE:HG3	1.99	0.45
1:B:155[B]:HIS:HB3	1:C:155[B]:HIS:CB	2.32	0.45
1:A:74:ARG:HE	1:A:74:ARG:HB2	1.65	0.44
1:B:0:ALA:O	1:D:75:HIS:HD2	2.01	0.44
1:B:90:GLU:HG3	1:B:91:ASN:H	1.83	0.43
1:A:137:PHE:HZ	1:A:152:LEU:HD21	1.84	0.42
1:D:155:HIS:ND1	2:D:317:HOH:O	2.22	0.42
1:A:48:SER:HB3	1:A:67:ARG:CZ	2.50	0.42
1:D:97:MSE:N	1:D:97:MSE:SE	3.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:TYR:O	1:A:181:PRO:CB	2.67	0.41
1:C:16:ALA:HB1	1:C:67:ARG:HB3	2.01	0.41
1:C:91:ASN:CB	1:C:120:LEU:HD11	2.49	0.41
1:A:128:GLU:O	1:A:132:VAL:HG23	2.20	0.41
1:C:33[A]:VAL:HG21	1:C:171:LEU:CD1	2.44	0.41
1:D:82:VAL:HG12	1:D:84[A]:LEU:HD22	2.03	0.41
1:D:104[B]:GLN:HG3	1:D:105:ARG:N	2.36	0.41
1:B:116:PRO:HA	1:B:117:PRO:HD3	1.97	0.40
1:C:33[A]:VAL:CG2	1:C:171:LEU:HD13	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/185 (88%)	161 (99%)	2 (1%)	0	100	100
1	B	179/185 (97%)	176 (98%)	3 (2%)	0	100	100
1	C	172/185 (93%)	167 (97%)	2 (1%)	3 (2%)	7	2
1	D	173/185 (94%)	170 (98%)	3 (2%)	0	100	100
All	All	687/740 (93%)	674 (98%)	10 (2%)	3 (0%)	37	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	10	MSE
1	C	155[A]	HIS
1	C	155[B]	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/151 (94%)	135 (95%)	7 (5%)	21	13
1	B	156/151 (103%)	146 (94%)	10 (6%)	14	7
1	C	150/151 (99%)	145 (97%)	5 (3%)	33	26
1	D	148/151 (98%)	141 (95%)	7 (5%)	22	14
All	All	596/604 (99%)	567 (95%)	29 (5%)	23	13

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	ARG
1	A	103	GLN
1	A	113	LEU
1	A	123	LEU
1	A	125	GLN
1	A	171	LEU
1	A	178	GLU
1	B	4	ILE
1	B	84[A]	LEU
1	B	84[B]	LEU
1	B	107	GLU
1	B	128	GLU
1	B	141	ASN
1	B	144	GLN
1	B	155[A]	HIS
1	B	155[B]	HIS
1	B	178	GLU
1	C	124	ARG
1	C	143	LYS
1	C	151	LEU
1	C	159	ARG
1	C	178	GLU
1	D	11	ARG
1	D	84[A]	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	84[B]	LEU
1	D	97	MSE
1	D	144	GLN
1	D	158	THR
1	D	178	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	24	ASN
1	A	70	ASN
1	B	24	ASN
1	B	70	ASN
1	B	104	GLN
1	B	141	ASN
1	C	24	ASN
1	C	70	ASN
1	C	179	HIS
1	D	18	GLN
1	D	19	ASN
1	D	24	ASN
1	D	53	GLN
1	D	70	ASN
1	D	75	HIS
1	D	125	GLN
1	D	144	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	160/185 (86%)	0.43	10 (6%) 27 28	12, 24, 42, 54	4 (2%)
1	B	174/185 (94%)	-0.15	2 (1%) 77 79	9, 18, 36, 42	3 (1%)
1	C	167/185 (90%)	0.26	5 (2%) 52 55	9, 23, 33, 42	4 (2%)
1	D	168/185 (90%)	-0.31	0 100 100	8, 16, 30, 38	2 (1%)
All	All	669/740 (90%)	0.05	17 (2%) 58 60	8, 20, 35, 54	13 (1%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	9	VAL	4.3
1	A	129	LEU	3.9
1	A	126	ALA	3.6
1	C	9	VAL	3.5
1	A	181	PRO	3.4
1	B	120	LEU	3.3
1	A	123	LEU	2.8
1	A	137	PHE	2.8
1	C	124	ARG	2.7
1	A	136	VAL	2.6
1	A	132	VAL	2.6
1	A	130	GLY	2.4
1	B	123	LEU	2.3
1	C	38	GLU	2.2
1	C	138	GLY	2.2
1	C	128	GLU	2.1
1	A	138	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.