



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 31, 2025 – 10:08 PM JST

PDB ID : 8ZWY / pdb_00008zwy
Title : Structure-Based Mechanism and Specificity of Human Galactosyltransferase B3GalT5
Authors : Lo, J.M.; Ma, C.
Deposited on : 2024-06-13
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 2.0rc1
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	278	
1	B	278	
2	E	2	
2	G	2	
3	F	2	
4	H	6	
5	I	2	
5	J	2	

2 Entry composition i

There are 9 unique types of molecules in this entry. The entry contains 5179 atoms, of which 261 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-1,3-galactosyltransferase 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	267	Total	C	N	O	S	0	3	0
			2206	1432	371	390	13			
1	B	266	Total	C	N	O	S	0	0	0
			2189	1418	370	388	13			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	E	2	Total	C	H	N	O	0	0	0
			55	16	27	2	10			
2	G	2	Total	C	H	N	O	0	0	0
			55	16	27	2	10			

- Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	F	2	Total	C	H	N	O	0	0	0
			36	14	12	1	9			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	H	N				O
4	H	6	138	40	67	2	29	0	0	0

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-methyl beta-D-galactopyranoside.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	H	N				O
5	I	2	54	15	27	1	11	0	0	0
5	J	2	54	15	27	1	11	0	0	0

- Molecule 6 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mn	0	0
			1	1		
6	B	1	Total	Mn	0	0
			1	1		

- Molecule 7 is URIDINE-5'-DIPHOSPHATE (CCD ID: UDP) (formula: C₉H₁₄N₂O₁₂P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
7	A	1	36	9	11	2	12	2	0	0
7	B	1	36	9	11	2	12	2	0	0

- Molecule 8 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYLAMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: B3P) (formula: $C_{11}H_{26}N_2O_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
8	A	1	45	11	26	2	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
8	B	1	45	11	26	2	6	0	0

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	171	Total	O	0	0
			171	171		
9	B	57	Total	O	0	0
			57	57		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

- Molecule 1: Beta-1,3-galactosyltransferase 5




- Molecule 1: Beta-1,3-galactosyltransferase 5



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 3: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%

MAG1
FUC2

- Molecule 4: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-[α -L-fucopyranose-(1-6)]2-acetamido-2-deoxy- β -D-glucopyranose

Chain H:  83% 17%

MAG1
MAG2
BMA3
MAN4
MAN5
FUC6

- Molecule 5: 2-acetamido-2-deoxy- β -D-glucopyranose-(1-3)-methyl β -D-galactopyranoside

Chain I:  50% 50%

MAG1
MAG2

- Molecule 5: 2-acetamido-2-deoxy- β -D-glucopyranose-(1-3)-methyl β -D-galactopyranoside

Chain J:  50% 50%

MAG1
MAG2

4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	47.65Å 86.70Å 87.00Å 90.00° 95.25° 90.00°	Depositor
Resolution (Å)	47.44 – 1.95	Depositor
% Data completeness (in resolution range)	92.8 (47.44-1.95)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.48 (at 1.95Å)	Xtrriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, R_{free}	0.208 , 0.227	Depositor
Wilson B-factor (Å ²)	17.3	Xtrriage
Anisotropy	0.354	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5179	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, MN, MBG, B3P, FUC, NAG, MAN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/2274	0.85	0/3075
1	B	0.55	0/2248	0.71	0/3040
All	All	0.59	0/4522	0.78	0/6115

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	6
All	All	0	9

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	178	ARG	Sidechain
1	A	272	ARG	Sidechain
1	A	53	ARG	Sidechain
1	B	115	ARG	Sidechain
1	B	144	ARG	Sidechain
1	B	256	ARG	Sidechain
1	B	272	ARG	Sidechain
1	B	303	ARG	Sidechain
1	B	53	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2206	0	2205	23	0
1	B	2189	0	2175	43	0
2	E	28	27	25	0	0
2	G	28	27	25	4	0
3	F	24	12	22	0	0
4	H	71	67	61	0	0
5	I	27	27	26	2	0
5	J	27	27	26	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	25	11	11	0	0
7	B	25	11	11	0	0
8	A	19	26	26	0	0
8	B	19	26	26	0	0
9	A	171	0	0	3	0
9	B	57	0	0	0	0
All	All	4918	261	4639	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:288:LYS:HE3	1:B:290:ARG:HB3	1.57	0.85
1:B:46:LYS:NZ	1:B:110:ASP:HA	1.93	0.83
1:A:107:LYS:HA	1:A:107:LYS:HE2	1.70	0.74
1:B:47:LEU:HD12	1:B:48:PRO:HD2	1.69	0.73
1:B:193:GLN:HE21	1:B:195:PHE:HZ	1.37	0.72
1:B:46:LYS:HZ3	1:B:110:ASP:HA	1.58	0.68
1:B:56:PRO:HB3	1:B:148:GLN:HB3	1.74	0.68
1:A:232:VAL:HG21	1:A:252:ARG:HG2	1.75	0.66
1:B:46:LYS:HZ1	1:B:110:ASP:HA	1.63	0.63
1:A:232:VAL:CG2	1:A:252:ARG:HG2	2.29	0.62
1:B:79:THR:HG21	1:B:292:LEU:HG	1.82	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:193:GLN:NE2	1:B:195:PHE:CZ	2.63	0.60
1:A:189:PHE:CE1	1:A:209:ARG:HG2	2.36	0.60
1:B:46:LYS:HZ2	1:B:109:VAL:HG12	1.69	0.58
1:B:154:LYS:NZ	1:B:216:GLY:O	2.35	0.56
1:B:288:LYS:CE	1:B:290:ARG:HB3	2.33	0.56
2:G:1:NAG:H4	2:G:2:NAG:H83	1.87	0.56
1:B:187:ASN:OD1	1:B:209:ARG:NH2	2.37	0.56
1:A:107:LYS:O	1:A:111:GLN:HG3	2.05	0.56
1:B:193:GLN:NE2	1:B:195:PHE:HZ	2.03	0.56
1:A:56:PRO:HB3	1:A:148:GLN:HB3	1.89	0.54
1:B:232:VAL:HG21	1:B:252:ARG:HG2	1.89	0.54
1:A:303:ARG:HD3	1:A:304:GLY:N	2.23	0.53
1:B:189:PHE:CE1	1:B:209:ARG:HG2	2.44	0.52
1:A:231:ASN:HB2	9:A:620:HOH:O	2.10	0.51
1:B:152:VAL:HB	1:B:221:PHE:CE1	2.45	0.51
1:B:180:PHE:CD2	1:B:221:PHE:HB3	2.46	0.50
1:A:80:TRP:CZ3	1:A:161[B]:ILE:HD12	2.47	0.50
1:A:307:CYS:O	1:A:308:PRO:C	2.49	0.50
1:A:67:HIS:CD2	1:A:109:VAL:HG21	2.47	0.50
1:A:232:VAL:HG21	1:A:252:ARG:CG	2.42	0.50
1:B:72:GLU:HA	1:B:289:PRO:HG3	1.94	0.49
1:B:217:THR:HG23	5:J:2:NAG:H81	1.94	0.49
1:B:232:VAL:CG2	1:B:252:ARG:HG2	2.43	0.48
1:A:172:LYS:HG3	9:A:534:HOH:O	2.13	0.48
1:B:99:THR:HG21	1:B:123:ASP:HA	1.95	0.48
1:B:99:THR:CG2	1:B:123:ASP:HA	2.43	0.48
1:B:141:TRP:CD1	1:B:145:PHE:HD2	2.32	0.47
1:A:107:LYS:O	1:A:107:LYS:HD3	2.14	0.47
1:B:45:LEU:HB2	1:B:121:GLN:HG2	1.97	0.47
1:B:125:LEU:HD22	2:G:2:NAG:H81	1.97	0.46
1:B:97:LEU:O	1:B:121:GLN:HA	2.15	0.46
1:B:46:LYS:HB2	1:B:121:GLN:HB3	1.98	0.46
1:A:80:TRP:CH2	1:A:161[B]:ILE:HD12	2.50	0.46
1:A:189:PHE:CD1	1:A:209:ARG:HG2	2.51	0.45
1:B:63:VAL:HG22	1:B:155:THR:CG2	2.46	0.45
1:B:56:PRO:HB3	1:B:148:GLN:CB	2.46	0.44
1:B:44:PHE:HB3	1:B:47:LEU:HD13	1.99	0.44
1:B:189:PHE:HE1	1:B:209:ARG:HG2	1.81	0.44
1:A:146:CYS:N	1:A:147:PRO:CD	2.81	0.44
1:A:190:PRO:HD2	1:A:208:ASP:O	2.17	0.44
2:G:1:NAG:H61	2:G:2:NAG:H4	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:284:CYS:O	1:A:287[B]:ILE:HD12	2.18	0.43
1:B:99:THR:HG21	1:B:123:ASP:OD1	2.17	0.43
1:B:46:LYS:HZ1	1:B:110:ASP:CA	2.28	0.43
1:A:241:LEU:HD13	5:I:2:NAG:H62	2.00	0.43
1:A:125:LEU:HG	1:A:127:VAL:HG13	1.99	0.43
1:B:43:ASN:O	1:B:122:LYS:HG2	2.18	0.43
1:B:172:LYS:HE3	1:B:172:LYS:HB2	1.73	0.42
1:B:273:PHE:HB2	1:B:295:TYR:CZ	2.55	0.42
1:B:103:ALA:O	1:B:107:LYS:HG3	2.19	0.42
9:A:615:HOH:O	5:I:2:NAG:H83	2.19	0.42
1:A:44:PHE:HA	1:A:122:LYS:HG2	2.02	0.42
1:A:176:THR:HG23	1:A:177:THR:HG23	2.02	0.42
1:B:146:CYS:N	1:B:147:PRO:CD	2.83	0.42
1:B:285:HIS:CD2	1:B:286:PHE:H	2.37	0.42
1:B:91:LEU:HD13	1:B:163:VAL:HG11	2.01	0.41
1:B:125:LEU:HB3	2:G:1:NAG:H2	2.03	0.41
1:B:162:ASN:HB3	1:B:281:ILE:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/278 (96%)	262 (98%)	6 (2%)	0	100	100
1	B	264/278 (95%)	254 (96%)	10 (4%)	0	100	100
All	All	532/556 (96%)	516 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/255 (97%)	247 (100%)	0	100	100
1	B	244/255 (96%)	244 (100%)	0	100	100
All	All	491/510 (96%)	491 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	E	1	2,1	14,14,15	0.46	0	17,19,21	0.93	1 (5%)
2	NAG	E	2	2	14,14,15	0.72	0	17,19,21	0.55	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	F	1	3,1	14,14,15	0.42	0	17,19,21	0.47	0
3	FUC	F	2	3	10,10,11	0.85	0	14,14,16	0.95	1 (7%)
2	NAG	G	1	2,1	14,14,15	0.43	0	17,19,21	1.44	1 (5%)
2	NAG	G	2	2	14,14,15	0.40	0	17,19,21	0.71	1 (5%)
4	NAG	H	1	4,1	14,14,15	0.44	0	17,19,21	1.00	1 (5%)
4	NAG	H	2	4	14,14,15	0.45	0	17,19,21	0.89	0
4	BMA	H	3	4	11,11,12	0.60	0	15,15,17	0.52	0
4	MAN	H	4	4	11,11,12	0.33	0	15,15,17	0.57	0
4	MAN	H	5	4	11,11,12	0.35	0	15,15,17	0.75	0
4	FUC	H	6	4	10,10,11	0.31	0	14,14,16	0.51	0
5	MBG	I	1	5	13,13,13	0.26	0	18,18,18	0.39	0
5	NAG	I	2	5	14,14,15	0.42	0	17,19,21	0.67	0
5	MBG	J	1	5	13,13,13	0.45	0	18,18,18	0.96	1 (5%)
5	NAG	J	2	5	14,14,15	0.46	0	17,19,21	1.74	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
3	NAG	F	1	3,1	-	0/6/23/26	0/1/1/1
3	FUC	F	2	3	-	-	0/1/1/1
2	NAG	G	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	G	2	2	-	5/6/23/26	0/1/1/1
4	NAG	H	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	H	2	4	-	0/6/23/26	0/1/1/1
4	BMA	H	3	4	-	0/2/19/22	0/1/1/1
4	MAN	H	4	4	-	0/2/19/22	0/1/1/1
4	MAN	H	5	4	-	2/2/19/22	0/1/1/1
4	FUC	H	6	4	-	-	0/1/1/1
5	MBG	I	1	5	-	0/4/24/24	0/1/1/1
5	NAG	I	2	5	-	0/6/23/26	0/1/1/1
5	MBG	J	1	5	-	4/4/24/24	0/1/1/1
5	NAG	J	2	5	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	J	2	NAG	O5-C1-C2	-5.91	101.96	111.29
2	G	1	NAG	O5-C1-C2	-5.29	102.93	111.29
4	H	1	NAG	C2-N2-C7	2.84	126.95	122.90
2	G	2	NAG	C1-O5-C5	2.64	115.78	112.19
2	E	1	NAG	C2-N2-C7	2.13	125.94	122.90
3	F	2	FUC	C1-O5-C5	2.10	117.53	112.78
5	J	1	MBG	C1-O5-C5	-2.07	109.62	113.69

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	G	2	NAG	C8-C7-N2-C2
2	G	2	NAG	O7-C7-N2-C2
4	H	1	NAG	C8-C7-N2-C2
4	H	1	NAG	O7-C7-N2-C2
5	J	2	NAG	O7-C7-N2-C2
5	J	2	NAG	C8-C7-N2-C2
5	J	1	MBG	O5-C1-O1-C7
5	J	1	MBG	C2-C1-O1-C7
2	G	2	NAG	O5-C5-C6-O6
2	G	2	NAG	C4-C5-C6-O6
5	J	1	MBG	C4-C5-C6-O6
4	H	5	MAN	C4-C5-C6-O6
4	H	5	MAN	O5-C5-C6-O6
2	G	1	NAG	C8-C7-N2-C2
2	G	2	NAG	C3-C2-N2-C7
5	J	1	MBG	O5-C5-C6-O6
2	G	1	NAG	O7-C7-N2-C2


There are no ring outliers.


4 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	2	NAG	3	0
2	G	1	NAG	3	0
5	J	2	NAG	1	0
5	I	2	NAG	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	B3P	A	403	-	18,18,18	0.98	0	21,23,23	1.47	3 (14%)
7	UDP	B	402	6	24,26,26	0.55	0	37,40,40	0.52	0
7	UDP	A	402	6	24,26,26	1.54	1 (4%)	37,40,40	0.62	1 (2%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	B3P	B	403	-	18,18,18	0.74	0	21,23,23	1.53	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	B3P	A	403	-	-	0/28/28/28	-
7	UDP	B	402	6	-	7/16/32/32	0/2/2/2
7	UDP	A	402	6	-	2/16/32/32	0/2/2/2
8	B3P	B	403	-	-	3/28/28/28	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	402	UDP	PB-O1B	7.13	1.73	1.50

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	403	B3P	C2-N2-C8	-4.02	110.37	116.08
8	A	403	B3P	C7-C4-C5	-3.46	102.72	110.04
8	B	403	B3P	O1-C9-C8	-2.80	105.96	111.63
8	A	403	B3P	O1-C9-C8	-2.34	106.89	111.63
7	A	402	UDP	O2B-PB-O1B	2.22	119.39	110.68
8	A	403	B3P	C3-N1-C4	-2.22	112.94	116.08
8	B	403	B3P	O2-C10-C8	-2.19	107.19	111.63
8	B	403	B3P	C11-C8-C10	-2.08	105.65	110.04
8	B	403	B3P	C3-N1-C4	-2.04	113.18	116.08

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	402	UDP	O4'-C4'-C5'-O5'
7	B	402	UDP	C5'-O5'-PA-O1A
7	B	402	UDP	PB-O3A-PA-O5'
8	B	403	B3P	N1-C4-C6-O5
8	B	403	B3P	C5-C4-C6-O5

Continued on next page...


Continued from previous page...



Mol	Chain	Res	Type	Atoms
8	B	403	B3P	C7-C4-C6-O5
7	B	402	UDP	C5'-O5'-PA-O2A
7	A	402	UDP	PB-O3A-PA-O1A
7	B	402	UDP	C4'-C5'-O5'-PA
7	B	402	UDP	C5'-O5'-PA-O3A
7	A	402	UDP	PB-O3A-PA-O2A
7	B	402	UDP	PB-O3A-PA-O1A


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.


6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

6.3 Carbohydrates [i](#)




EDS failed to run properly - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








6.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MN A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UDP A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UDP B 402:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around B3P A 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.