



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 05:59 pm GMT

PDB ID : 3ZY2  
Title : Crystal structure of POFUT1 in complex with GDP (High resolution dataset)  
Authors : Lira-Navarrete, E.; Valero-Gonzalez, J.; Villanueva, R.; Martinez-Julvez, M.;  
Tejero, T.; Merino, P.; Panjekar, S.; Hurtado-Guerrero, R.  
Deposited on : 2011-08-17  
Resolution : 1.54 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2055 (1.56-1.52)
Clashscore	122126	2124 (1.56-1.52)
Ramachandran outliers	120053	2083 (1.56-1.52)
Sidechain outliers	120020	2081 (1.56-1.52)
RSRZ outliers	108989	2027 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GDP	A	1381[A]	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PUTATIVE GDP-FUCOSE PROTEIN O-FUCOSYLTRANSFERASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	342	2734	1763	461	495	15	0	0	0


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLU	-	expression tag	UNP Q18014
A	23	ALA	-	expression tag	UNP Q18014
A	24	GLU	-	expression tag	UNP Q18014

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
2	A	1	1	1	0	0

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	56	20	10	22	4	0	1


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	351	351	351	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PUTATIVE GDP-FUCOSE PROTEIN O-FUCOSYLTRANSFERASE 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	145.02Å 38.25Å 68.20Å 90.00° 103.07° 90.00°	Depositor
Resolution (Å)	70.63 – 1.54 19.50 – 1.54	Depositor EDS
% Data completeness (in resolution range)	98.4 (70.63-1.54) 98.5 (19.50-1.54)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.24 (at 1.54Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.205 , 0.237 0.202 , 0.237	Depositor DCC
$R_{free}$ test set	1010 reflections (1.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.7	Xtrriage
Anisotropy	0.480	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 57.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3142	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/2814	0.67	2/3812 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	77	MET	CB-CG-SD	-5.79	95.03	112.40
1	A	40	ARG	NE-CZ-NH2	-5.74	117.43	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2734	0	2663	28	0
2	A	1	0	0	0	0
3	A	56	0	17	16	0
4	A	351	0	0	7	0
All	All	3142	0	2680	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1381[B]:GDP:O2A	3:A:1381[B]:GDP:PA	1.15	1.53
3:A:1381[A]:GDP:C3'	3:A:1381[A]:GDP:O3'	1.74	1.34
3:A:1381[A]:GDP:C8	3:A:1381[A]:GDP:N9	1.83	1.32
3:A:1381[A]:GDP:C1'	3:A:1381[A]:GDP:O4'	1.77	1.30
3:A:1381[A]:GDP:C2'	3:A:1381[A]:GDP:C3'	2.19	1.19
3:A:1381[B]:GDP:O2A	3:A:1381[B]:GDP:O3A	1.86	0.91
3:A:1381[A]:GDP:C1'	3:A:1381[A]:GDP:C3'	2.49	0.89
3:A:1381[B]:GDP:O5'	3:A:1381[B]:GDP:O2A	1.91	0.86
1:A:291:GLN:HE22	1:A:379:GLY:H	1.25	0.84
3:A:1381[A]:GDP:C1'	3:A:1381[A]:GDP:C4'	2.61	0.78
1:A:210:LYS:HD2	4:A:2191:HOH:O	1.91	0.70
1:A:256:THR:HG23	4:A:2236:HOH:O	1.94	0.67
1:A:98:MET:O	1:A:102:THR:HB	2.00	0.61
3:A:1381[A]:GDP:H2'	3:A:1381[A]:GDP:N3	2.15	0.60
1:A:102:THR:HG21	4:A:2102:HOH:O	2.02	0.60
1:A:359:HIS:HD2	4:A:2025:HOH:O	1.84	0.60
1:A:329:HIS:HD2	4:A:2115:HOH:O	1.85	0.58
1:A:77:MET:HG3	1:A:149:GLN:HG2	1.85	0.57
1:A:312:HIS:HE1	1:A:334:ASP:OD2	1.87	0.57
3:A:1381[A]:GDP:C2'	3:A:1381[A]:GDP:N3	2.64	0.57
1:A:359:HIS:HE1	4:A:2325:HOH:O	1.85	0.56
3:A:1381[A]:GDP:C1'	3:A:1381[A]:GDP:C8	2.80	0.56
1:A:257:ASN:HD21	1:A:276:LEU:H	1.52	0.56
1:A:99:GLN:HG3	1:A:103:LYS:HD3	1.88	0.55
1:A:312:HIS:HD2	4:A:2277:HOH:O	1.91	0.54
3:A:1381[A]:GDP:O3B	3:A:1381[A]:GDP:O5'	2.27	0.53
1:A:291:GLN:NE2	1:A:379:GLY:H	2.02	0.51
1:A:164:PRO:HD2	1:A:177:LYS:HE2	1.93	0.49
1:A:257:ASN:ND2	1:A:271:HIS:HD2	2.12	0.48
1:A:137:SER:HB3	1:A:193:SER:HB2	1.96	0.47
1:A:138:LYS:CE	1:A:152:VAL:O	2.63	0.47
1:A:71:LYS:HZ3	1:A:71:LYS:H	1.63	0.46
1:A:253:ASP:OD1	1:A:255:THR:OG1	2.32	0.46
1:A:247:ARG:O	1:A:250:GLU:HG2	2.15	0.46
3:A:1381[A]:GDP:C3'	3:A:1381[A]:GDP:HO3'	2.17	0.46
1:A:263:SER:OG	1:A:271:HIS:HE1	1.98	0.46
1:A:257:ASN:ND2	1:A:276:LEU:H	2.14	0.45
1:A:309:ASP:HB3	3:A:1381[B]:GDP:C8	2.53	0.43
1:A:102:THR:HG22	1:A:103:LYS:HG3	2.00	0.43
3:A:1381[A]:GDP:C2'	3:A:1381[A]:GDP:O3'	2.66	0.43

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:HIS:HB2	1:A:74:GLU:O	2.19	0.42
1:A:106:MET:HG3	1:A:110:TRP:HE3	1.84	0.42
1:A:223:LYS:HE2	1:A:345:ARG:NH2	2.35	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	336/362 (93%)	333 (99%)	3 (1%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	297/312 (95%)	289 (97%)	8 (3%)	48 15

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	LYS
1	A	102	THR
1	A	167	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	184	SER
1	A	190	LEU
1	A	193	SER
1	A	286	GLN
1	A	359	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	ASN
1	A	257	ASN
1	A	271	HIS
1	A	291	GLN
1	A	312	HIS
1	A	329	HIS
1	A	331	GLN
1	A	359	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GDP	A	1381[A]	-	25,30,30	12.95	23 (92%)	27,47,47	7.64	20 (74%)
3	GDP	A	1381[B]	-	25,30,30	7.27	19 (76%)	27,47,47	5.39	18 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	1381[A]	-	-	0/12/32/32	0/3/3/3
3	GDP	A	1381[B]	-	-	0/12/32/32	0/3/3/3

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1381[A]	GDP	C2-N2	-16.58	1.00	1.33
3	A	1381[B]	GDP	PB-O3A	-16.13	1.35	1.60
3	A	1381[B]	GDP	C8-N9	-14.11	1.19	1.36
3	A	1381[A]	GDP	C6-N1	-13.91	1.08	1.33
3	A	1381[B]	GDP	O4'-C1'	-11.89	1.24	1.41
3	A	1381[B]	GDP	PA-O2A	-8.11	1.15	1.55
3	A	1381[A]	GDP	C2-N1	-5.96	1.24	1.35
3	A	1381[B]	GDP	PA-O1A	-5.89	1.29	1.50
3	A	1381[B]	GDP	C6-C5	-5.85	1.31	1.41
3	A	1381[A]	GDP	C3'-C4'	-5.32	1.39	1.53
3	A	1381[A]	GDP	C5'-C4'	-5.18	1.35	1.51
3	A	1381[B]	GDP	C2'-C3'	-4.97	1.40	1.53
3	A	1381[A]	GDP	PA-O2A	-4.48	1.33	1.55
3	A	1381[B]	GDP	C4-N3	-4.08	1.29	1.35
3	A	1381[A]	GDP	O6-C6	-3.93	1.14	1.24
3	A	1381[A]	GDP	O5'-C5'	-3.89	1.29	1.44
3	A	1381[B]	GDP	C2-N3	-3.65	1.16	1.35
3	A	1381[A]	GDP	O4'-C4'	-3.35	1.37	1.45
3	A	1381[B]	GDP	PB-O3B	-2.85	1.43	1.54
3	A	1381[A]	GDP	O2'-C2'	-2.74	1.36	1.43
3	A	1381[A]	GDP	C8-N7	-2.58	1.29	1.34
3	A	1381[B]	GDP	C5-C4	2.44	1.46	1.40
3	A	1381[A]	GDP	PB-O3B	2.57	1.65	1.54
3	A	1381[B]	GDP	O3'-C3'	2.60	1.49	1.43

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1381[B]	GDP	C8-N7	3.41	1.41	1.34
3	A	1381[B]	GDP	O6-C6	3.41	1.33	1.24
3	A	1381[B]	GDP	O2'-C2'	3.58	1.51	1.43
3	A	1381[B]	GDP	PB-O1B	4.40	1.65	1.50
3	A	1381[A]	GDP	C2-N3	4.45	1.58	1.35
3	A	1381[A]	GDP	PA-O5'	5.07	1.79	1.59
3	A	1381[A]	GDP	C4-N3	5.64	1.44	1.35
3	A	1381[A]	GDP	PA-O1A	6.57	1.74	1.50
3	A	1381[A]	GDP	PB-O3A	8.12	1.72	1.60
3	A	1381[A]	GDP	C5-C4	8.50	1.59	1.40
3	A	1381[B]	GDP	C2-N2	9.22	1.52	1.33
3	A	1381[A]	GDP	O3'-C3'	12.66	1.74	1.43
3	A	1381[B]	GDP	C6-N1	12.67	1.55	1.33
3	A	1381[B]	GDP	C2-N1	14.22	1.60	1.35
3	A	1381[A]	GDP	C6-C5	18.59	1.74	1.41
3	A	1381[A]	GDP	C2'-C3'	25.33	2.19	1.53
3	A	1381[A]	GDP	O4'-C1'	26.13	1.77	1.41
3	A	1381[A]	GDP	C8-N9	38.25	1.83	1.36

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1381[B]	GDP	N2-C2-N1	-13.01	96.81	117.25
3	A	1381[A]	GDP	C5-C6-N1	-12.79	105.28	123.47
3	A	1381[B]	GDP	C6-C5-C4	-12.38	108.70	120.85
3	A	1381[A]	GDP	C2-N3-C4	-11.07	102.24	115.16
3	A	1381[A]	GDP	C6-C5-C4	-10.37	110.68	120.85
3	A	1381[A]	GDP	N2-C2-N3	-9.81	99.83	117.75
3	A	1381[A]	GDP	C4-C5-N7	-9.29	100.44	109.41
3	A	1381[A]	GDP	O3'-C3'-C2'	-8.56	84.43	111.83
3	A	1381[B]	GDP	C4-C5-N7	-7.94	101.74	109.41
3	A	1381[B]	GDP	C6-N1-C2	-6.05	107.35	116.06
3	A	1381[B]	GDP	O2'-C2'-C3'	-4.79	96.48	111.83
3	A	1381[A]	GDP	C1'-N9-C4	-4.75	118.43	126.64
3	A	1381[A]	GDP	PA-O5'-C5'	-4.71	94.05	121.68
3	A	1381[A]	GDP	O3A-PB-O1B	-4.71	83.06	111.48
3	A	1381[B]	GDP	O2A-PA-O5'	-4.10	88.71	107.75
3	A	1381[A]	GDP	C5'-C4'-C3'	-3.73	101.26	115.29
3	A	1381[A]	GDP	O2'-C2'-C3'	-3.23	101.49	111.83
3	A	1381[B]	GDP	O4'-C4'-C5'	-3.18	98.81	109.39
3	A	1381[B]	GDP	O4'-C4'-C3'	-3.00	99.21	105.15
3	A	1381[B]	GDP	O3'-C3'-C4'	-2.32	104.35	111.06

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1381[B]	GDP	N3-C2-N1	-2.26	124.09	127.41
3	A	1381[A]	GDP	O3B-PB-O2B	2.01	115.56	107.59
3	A	1381[A]	GDP	O2A-PA-O1A	2.25	123.56	112.14
3	A	1381[B]	GDP	C5-C6-N1	2.40	126.89	123.47
3	A	1381[B]	GDP	C5'-C4'-C3'	3.37	127.97	115.29
3	A	1381[A]	GDP	C2'-C3'-C4'	3.65	109.62	102.62
3	A	1381[B]	GDP	C4'-O4'-C1'	3.82	113.81	109.83
3	A	1381[B]	GDP	C2-N3-C4	4.20	120.07	115.16
3	A	1381[B]	GDP	PA-O3A-PB	4.29	147.04	132.63
3	A	1381[A]	GDP	O2B-PB-O1B	4.74	129.11	110.60
3	A	1381[A]	GDP	O5'-C5'-C4'	5.02	126.46	109.00
3	A	1381[B]	GDP	O3'-C3'-C2'	5.74	130.21	111.83
3	A	1381[A]	GDP	O3'-C3'-C4'	6.10	128.73	111.06
3	A	1381[B]	GDP	O2A-PA-O1A	6.38	144.58	112.14
3	A	1381[A]	GDP	O4'-C4'-C5'	8.81	138.74	109.39
3	A	1381[A]	GDP	N2-C2-N1	11.47	135.28	117.25
3	A	1381[B]	GDP	N2-C2-N3	11.68	139.09	117.75
3	A	1381[A]	GDP	C6-N1-C2	22.41	148.30	116.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1381[A]	GDP	12	0
3	A	1381[B]	GDP	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	342/362 (94%)	0.67	35 (10%) <b>7</b> <b>7</b>	10, 21, 39, 46	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	25	ALA	7.4
1	A	164	PRO	5.3
1	A	251	HIS	5.2
1	A	255	THR	4.4
1	A	120	TRP	4.4
1	A	180	GLU	4.2
1	A	247	ARG	3.8
1	A	179	LEU	3.5
1	A	165	GLY	3.4
1	A	177	LYS	3.3
1	A	243	ALA	3.3
1	A	168	ASP	3.2
1	A	163	ILE	3.1
1	A	123	ARG	3.0
1	A	250	GLU	3.0
1	A	169	LEU	3.0
1	A	254	THR	2.8
1	A	380	ILE	2.8
1	A	354	VAL	2.8
1	A	132	GLU	2.8
1	A	167	PHE	2.7
1	A	256	THR	2.7
1	A	33	VAL	2.4
1	A	292	ILE	2.4
1	A	71	LYS	2.4
1	A	72	HIS	2.3
1	A	162	ASP	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	244	ASP	2.2
1	A	246	VAL	2.2
1	A	184	SER	2.1
1	A	63	VAL	2.1
1	A	183	PRO	2.1
1	A	286	GLN	2.1
1	A	252	ILE	2.0
1	A	272	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MN	A	1380	1/1	0.95	0.14	14,14,14,14	1
3	GDP	A	1381[A]	28/28	0.98	0.07	10,23,29,50	28
3	GDP	A	1381[B]	28/28	0.98	0.07	5,9,13,14	28

## 6.5 Other polymers [i](#)

There are no such residues in this entry.