



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 07:00 AM UTC

PDB ID : 1NGE / pdb_00001nge
Title : STRUCTURAL BASIS OF THE 70-KILODALTON HEAT SHOCK COGNATE PROTEIN ATP HYDROLYTIC ACTIVITY, II. STRUCTURE OF THE ACTIVE SITE WITH ADP OR ATP BOUND TO WILD TYPE AND MUTANT ATPASE FRAGMENT
Authors : Flaherty, K.M.; Wilbanks, S.M.; Deluca-Flaherty, C.; Mckay, D.B.
Deposited on : 1994-05-17
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtrriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3150 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HEAT-SHOCK COGNATE 70 kD PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	378	2926	1838	514	566	8	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	199	SER	ASP	conflict	UNP P19120

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	193	Total 193	O 193	0	0

SEQUENCE-PLOTS INFOmissingINFO

3 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	145.30 Å 65.00 Å 46.90 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.05	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.05)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.180 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3150	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

4 Model quality [i](#)

4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/2972	0.87	6/4014 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	48	GLU	N-CA-C	7.53	118.54	108.38
1	A	343	ILE	CA-C-N	5.70	125.17	119.24
1	A	343	ILE	C-N-CA	5.70	125.17	119.24
1	A	130	ILE	N-CA-C	-5.47	105.38	110.53
1	A	29	ILE	N-CA-C	5.36	115.96	108.89
1	A	248	LYS	N-CA-C	5.06	116.80	111.28

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2926	0	2934	39	0
2	A	31	0	12	0	0
3	A	193	0	0	2	0
All	All	3150	0	2946	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:ASP:O	1:A:108:LYS:HG2	1.90	0.70
1:A:310:PHE:O	1:A:313:THR:HB	1.91	0.70
1:A:319:LYS:HD2	1:A:322:ARG:NH1	2.10	0.66
1:A:151:ASN:H	1:A:154:GLN:HE21	1.43	0.66
1:A:251:LYS:NZ	1:A:289:GLU:HG3	2.16	0.61
1:A:21:PHE:CZ	1:A:24:GLY:HA2	2.39	0.58
1:A:87:MET:HA	1:A:90:TRP:CE3	2.38	0.57
1:A:38:THR:OG1	1:A:127:MET:HE1	2.05	0.56
1:A:319:LYS:HD2	1:A:322:ARG:HH11	1.71	0.55
1:A:257:LYS:O	1:A:261:ARG:HG3	2.08	0.53
1:A:160:ASP:O	1:A:164:ILE:HG23	2.08	0.53
1:A:269:ARG:HG2	1:A:269:ARG:HH11	1.74	0.52
1:A:379:ILE:HG13	1:A:380:LEU:N	2.28	0.49
1:A:235:ASN:OD1	1:A:264:ARG:NH1	2.45	0.49
1:A:60:ALA:HA	1:A:261:ARG:HD3	1.95	0.49
1:A:121:SER:CB	1:A:164:ILE:HG12	2.44	0.47
1:A:175:GLU:N	1:A:176:PRO:HD2	2.31	0.46
1:A:49:ARG:NH1	3:A:672:HOH:O	2.48	0.46
1:A:151:ASN:H	1:A:154:GLN:NE2	2.13	0.45
1:A:62:ASN:OD1	1:A:65:ASN:HB2	2.17	0.45
1:A:44:PHE:CD2	1:A:105:VAL:HG21	2.51	0.44
1:A:119:VAL:O	1:A:122:MET:HB2	2.18	0.44
1:A:146:VAL:HG22	1:A:150:PHE:CD2	2.53	0.44
1:A:315:ASP:HB2	1:A:316:PRO:HD3	2.01	0.43
1:A:216:ILE:N	1:A:216:ILE:HD12	2.33	0.43
1:A:210:LEU:CD2	1:A:219:VAL:HG22	2.49	0.42
1:A:233:PHE:HA	1:A:306:ASN:HD21	1.85	0.42
1:A:269:ARG:HG2	1:A:269:ARG:NH1	2.34	0.42
1:A:331:ILE:HD11	1:A:334:ILE:CG1	2.50	0.41
1:A:128:LYS:NZ	1:A:132:GLU:OE2	2.53	0.41
1:A:199:SER:HA	1:A:337:VAL:O	2.21	0.41
1:A:200:LEU:HD11	1:A:228:LEU:HD13	2.02	0.41
1:A:246:LYS:O	1:A:250:LYS:HA	2.21	0.41
1:A:328:LYS:NZ	3:A:535:HOH:O	2.53	0.41
1:A:23:HIS:O	1:A:25:LYS:HG3	2.20	0.41
1:A:315:ASP:CB	1:A:316:PRO:HD3	2.51	0.41
1:A:126:LYS:HA	1:A:126:LYS:HD2	1.95	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:LEU:HD23	1:A:162:GLY:HA2	2.03	0.40
1:A:150:PHE:CD2	1:A:154:GLN:HB3	2.56	0.40

There are no symmetry-related clashes.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/118 (319%)	364 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

4.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/98 (319%)	279 (89%)	34 (11%)	6	2

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	GLN
1	A	32	ASP
1	A	38	THR
1	A	50	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	58	GLN
1	A	73	LEU
1	A	79	ASP
1	A	110	GLU
1	A	124	LEU
1	A	138	THR
1	A	146	VAL
1	A	156	GLN
1	A	159	LYS
1	A	164	ILE
1	A	167	LEU
1	A	170	LEU
1	A	171	ARG
1	A	228	LEU
1	A	250	LYS
1	A	251	LYS
1	A	255	GLU
1	A	274	LEU
1	A	283	GLU
1	A	297	ILE
1	A	301	ARG
1	A	309	LEU
1	A	313	THR
1	A	321	LEU
1	A	331	ILE
1	A	335	VAL
1	A	350	LEU
1	A	358	GLU
1	A	359	LEU
1	A	380	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	57	ASN
1	A	154	GLN
1	A	156	GLN
1	A	194	ASN
1	A	227	HIS
1	A	249	HIS
1	A	306	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	351	GLN
1	A	376	GLN

4.3.3 RNA ⓘ

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

4.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ATP	A	486	-	32,33,33	1.42	3 (9%)	48,52,52	1.18	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	486	-	-	1/22/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	486	ATP	PB-O3A	-4.41	1.54	1.59
2	A	486	ATP	PA-O3A	-3.52	1.55	1.59
2	A	486	ATP	C2-N1	-2.04	1.30	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	486	ATP	O3'-C3'-C4'	-2.76	103.14	111.08
2	A	486	ATP	O2A-PA-O3A	2.63	114.39	107.27
2	A	486	ATP	O3G-PG-O2G	2.60	117.54	107.80
2	A	486	ATP	C1'-N9-C8	2.40	132.41	127.09
2	A	486	ATP	C4-N9-C1'	-2.13	121.64	126.63
2	A	486	ATP	O3B-PB-O1B	-2.09	104.43	110.70

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	486	ATP	PA-O3A-PB-O1B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

5.3 Carbohydrates

EDS was not executed - this section is therefore empty.

5.4 Ligands

EDS was not executed - this section is therefore empty.

5.5 Other polymers

EDS was not executed - this section is therefore empty.