



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:43 PM UTC

PDB ID : 2O5E / pdb\_00002o5e  
Title : Structure of E. coli topoisomerase III in complex with an 8-base single stranded oligonucleotide. Frozen in glucose pH 7.0  
Authors : Changela, A.; DiGate, R.J.; Mondragon, A.  
Deposited on : 2006-12-05  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	8	 88% 12%
1	D	8	 25% 62% 12%
2	A	659	 3% 86% 9% . .
2	B	659	 3% 85% 10% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TDR	D	9	-	X	-	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10496 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*CP\*GP\*CP\*AP\*AP\*CP\*TP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	7	Total	C	N	O	P	0	0	0
			138	67	26	39	6			
1	D	7	Total	C	N	O	P	0	0	0
			138	67	26	39	6			


- Molecule 2 is a protein called DNA topoisomerase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	634	Total	C	N	O	S	0	0	0
			5042	3184	918	920	20			
2	B	631	Total	C	N	O	S	0	0	0
			5020	3170	913	917	20			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	654	HIS	-	expression tag	UNP P14294
A	655	HIS	-	expression tag	UNP P14294
A	656	HIS	-	expression tag	UNP P14294
A	657	HIS	-	expression tag	UNP P14294
A	658	HIS	-	expression tag	UNP P14294
A	659	HIS	-	expression tag	UNP P14294
B	654	HIS	-	expression tag	UNP P14294
B	655	HIS	-	expression tag	UNP P14294
B	656	HIS	-	expression tag	UNP P14294
B	657	HIS	-	expression tag	UNP P14294
B	658	HIS	-	expression tag	UNP P14294
B	659	HIS	-	expression tag	UNP P14294

- Molecule 3 is THYMINE (CCD ID: TDR) (formula: C<sub>5</sub>H<sub>6</sub>N<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	D	1	Total	C	N	O	0	0
			9	5	2	2		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Cl	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	3	Total	O	0	0
			3	3		
5	D	4	Total	O	0	0
			4	4		
5	A	85	Total	O	0	0
			85	85		
5	B	55	Total	O	0	0
			55	55		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(\*CP\*GP\*CP\*AP\*AP\*CP\*TP\*T)-3'

Chain C: 





- Molecule 1: 5'-D(\*CP\*GP\*CP\*AP\*AP\*CP\*TP\*T)-3'

Chain D: 





- Molecule 2: DNA topoisomerase 3


Chain A: 



- Molecule 2: DNA topoisomerase 3

Chain B: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.26Å 102.26Å 445.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.13 – 2.50 29.13 – 2.50	Depositor EDS
% Data completeness (in resolution range)	95.7 (29.13-2.50) 95.7 (29.13-2.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.44 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.217 , 0.260 0.209 , 0.249	Depositor DCC
$R_{free}$ test set	3992 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	58.1	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 41.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10496	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, TDR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	C	0.51	0/154	1.10	0/235
1	D	0.50	0/154	1.13	1/235 (0.4%)
2	A	0.64	0/5154	0.85	0/6989
2	B	0.61	0/5130	0.87	3/6955 (0.0%)
All	All	0.62	0/10592	0.87	4/14414 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	1	DC	N1-C1'-C2'	-5.83	104.75	113.50
2	B	614	PHE	N-CA-C	5.51	120.12	113.17
2	B	613	GLN	N-CA-C	5.32	122.12	110.80
2	B	293	SER	N-CA-C	-5.03	103.29	110.59

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	138	0	80	0	0
1	D	138	0	80	5	0
2	A	5042	0	5042	42	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5020	0	5023	34	0
3	D	9	0	6	0	0
4	A	2	0	0	0	0
5	A	85	0	0	3	0
5	B	55	0	0	0	0
5	C	3	0	0	0	0
5	D	4	0	0	0	0
All	All	10496	0	10231	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:419:VAL:HB	2:B:443:GLU:HG3	1.55	0.89
2:A:229:VAL:HG23	2:A:235:ARG:HG2	1.66	0.76
2:A:21:LYS:O	5:A:806:HOH:O	2.04	0.75
2:B:75:LEU:HG	2:B:647:VAL:HG21	1.74	0.70
2:B:11:LEU:HD12	2:B:137:LEU:HD21	1.73	0.69
2:A:247:GLU:H	2:A:248:PRO:CD	2.05	0.68
2:A:564:MET:HG3	2:A:570:MET:SD	2.35	0.67
2:A:187:ALA:HB2	2:A:611:VAL:HG13	1.78	0.64
2:A:259:ARG:C	2:A:261:LEU:H	2.07	0.62
2:A:304:ARG:HG3	2:A:304:ARG:HH11	1.64	0.62
2:A:177:THR:HG23	2:A:194:SER:HA	1.81	0.61
2:B:332:ASP:HB3	2:B:525:GLU:HG3	1.84	0.60
2:A:656:HIS:CD2	5:A:850:HOH:O	2.55	0.58
2:B:230:THR:HG22	2:B:232:ALA:H	1.69	0.57
2:B:6:ALA:HB1	2:B:11:LEU:HD23	1.87	0.56
2:A:240:TRP:HB2	2:A:434:PHE:CE2	2.42	0.55
2:A:176:MET:CE	2:A:603:ILE:HD13	2.38	0.54
2:A:11:LEU:HD12	2:A:137:LEU:HD21	1.89	0.54
2:B:653:ALA:HA	2:B:656:HIS:ND1	2.23	0.53
2:A:264:HIS:O	2:A:268:ARG:HB2	2.08	0.53
2:A:304:ARG:HG3	2:A:304:ARG:NH1	2.24	0.53
1:D:6:DC:H4'	1:D:7:DT:C7	2.40	0.52
2:A:247:GLU:H	2:A:248:PRO:HD3	1.73	0.51
1:D:3:DC:O4'	2:B:173:GLY:HA3	2.11	0.51
2:A:124:GLU:O	2:A:128:GLN:HG2	2.11	0.51
2:A:247:GLU:N	2:A:248:PRO:CD	2.74	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:197:ARG:HG2	2:B:571:THR:HG23	1.94	0.50
2:B:522:LEU:CD1	2:B:571:THR:HG21	2.42	0.50
2:B:510:LYS:O	2:B:513:LYS:HB3	2.11	0.50
2:A:117:ASP:HB3	2:A:647:VAL:HG13	1.93	0.49
2:B:308:SER:OG	2:B:311:ASN:HB2	2.12	0.49
2:B:522:LEU:HD11	2:B:571:THR:HG21	1.94	0.49
2:B:328:TYR:CZ	2:B:330:ARG:HB2	2.48	0.49
2:A:176:MET:HE2	2:A:603:ILE:HD13	1.95	0.48
2:A:587:ARG:HB3	2:A:656:HIS:CD2	2.49	0.48
2:B:87:VAL:O	2:B:90:ARG:HB3	2.14	0.48
2:A:225:LYS:CG	2:A:237:THR:HG23	2.44	0.48
2:A:431:LYS:HA	2:A:431:LYS:HE2	1.96	0.48
2:B:177:THR:HG23	2:B:194:SER:HA	1.95	0.47
2:B:240:TRP:HB2	2:B:434:PHE:CE2	2.49	0.47
2:A:92:LEU:HD22	2:A:121:LEU:HD21	1.96	0.47
2:A:110:LEU:O	2:A:114:GLU:HB2	2.14	0.47
2:B:240:TRP:O	2:B:242:PRO:HD3	2.15	0.47
2:A:247:GLU:N	2:A:248:PRO:HD3	2.30	0.46
2:A:137:LEU:HD12	2:A:321:GLU:HA	1.95	0.46
2:A:328:TYR:CZ	2:A:330:ARG:HB2	2.51	0.46
2:A:225:LYS:HG2	2:A:237:THR:HG23	1.97	0.46
2:A:180:TYR:O	2:A:193:LEU:HD12	2.15	0.45
2:A:283:ARG:HG3	2:A:421:ARG:HD3	1.98	0.45
2:A:453:LYS:HG3	2:A:454:GLU:H	1.80	0.45
2:B:612:ARG:O	2:B:614:PHE:N	2.50	0.45
2:A:583:GLU:O	2:A:585:GLN:HG2	2.17	0.45
2:B:230:THR:HA	2:B:473:LEU:HD22	1.98	0.45
2:A:259:ARG:C	2:A:261:LEU:N	2.73	0.44
2:B:99:VAL:HG22	2:B:130:GLN:HB2	2.00	0.44
2:B:224:VAL:HB	2:B:240:TRP:HB3	2.00	0.44
2:A:440:PHE:CZ	2:A:463:PRO:HD3	2.53	0.43
2:B:614:PHE:O	2:B:617:ILE:HG12	2.17	0.43
2:B:78:ARG:HA	2:B:79:PRO:HD3	1.89	0.43
2:B:428:ASP:C	2:B:429:ILE:HG13	2.42	0.43
2:B:426:GLU:HG2	2:B:435:VAL:HG22	2.01	0.43
2:A:176:MET:CE	2:A:603:ILE:CD1	2.96	0.42
2:A:54:TYR:CE1	2:A:70:PRO:HB3	2.54	0.42
2:B:178:ARG:O	2:B:182:ILE:HG13	2.18	0.42
2:B:405:LEU:HD23	2:B:405:LEU:HA	1.94	0.42
2:A:508:GLN:HB2	5:A:877:HOH:O	2.19	0.42
2:A:240:TRP:CZ2	2:A:242:PRO:HA	2.54	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:589:GLN:H	2:B:589:GLN:CD	2.27	0.42
2:B:200:THR:HB	2:B:201:PRO:HD3	2.01	0.41
2:A:305:PHE:CD1	2:A:401:LYS:HD3	2.55	0.41
2:B:100:HIS:CE1	2:B:112:VAL:HB	2.56	0.41
2:B:146:ILE:O	2:B:149:LEU:HG	2.21	0.41
1:D:6:DC:H4'	1:D:7:DT:H72	2.03	0.41
2:A:259:ARG:N	2:A:260:PRO:CD	2.83	0.41
1:D:5:DA:H2'	2:B:527:THR:HG21	2.01	0.41
1:D:6:DC:H4'	1:D:7:DT:H71	2.03	0.41
2:B:174:ILE:O	2:B:178:ARG:HG3	2.21	0.41
2:A:100:HIS:ND1	2:A:100:HIS:C	2.79	0.40
2:A:157:PRO:HG3	2:A:584:LYS:HG3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	630/659 (96%)	600 (95%)	27 (4%)	3 (0%)	24	43
2	B	627/659 (95%)	606 (97%)	18 (3%)	3 (0%)	24	43
All	All	1257/1318 (95%)	1206 (96%)	45 (4%)	6 (0%)	24	43

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	247	GLU
2	B	612	ARG
2	B	613	GLN
2	B	232	ALA
2	A	260	PRO
2	A	120	GLN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	538/555 (97%)	521 (97%)	17 (3%)	34	62
2	B	536/555 (97%)	522 (97%)	14 (3%)	40	68
All	All	1074/1110 (97%)	1043 (97%)	31 (3%)	37	65

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	17	ASP
2	A	27	ASP
2	A	48	GLN
2	A	100	HIS
2	A	212	GLU
2	A	219	LYS
2	A	229	VAL
2	A	233	ASP
2	A	246	CYS
2	A	283	ARG
2	A	367	ASP
2	A	404	ASN
2	A	454	GLU
2	A	462	THR
2	A	563	GLU
2	A	564	MET
2	A	611	VAL
2	B	18	VAL
2	B	27	ASP
2	B	76	GLN
2	B	80	SER
2	B	247	GLU
2	B	266	VAL
2	B	317	GLN
2	B	378	VAL
2	B	397	GLU
2	B	443	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	497	LEU
2	B	498	SER
2	B	546	ARG
2	B	584	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	48	GLN
2	A	135	ASN
2	A	190	GLN
2	A	215	ASN
2	A	345	HIS
2	A	486	GLN
2	A	593	GLN
2	A	659	HIS
2	B	264	HIS
2	B	323	HIS
2	B	361	GLN
2	B	486	GLN
2	B	573	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TDR	D	9	-	9,9,9	1.95	3 (33%)	12,12,12	3.02	8 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TDR	D	9	-	-	-	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	9	TDR	C6-C5	3.65	1.39	1.34
3	D	9	TDR	C4-C5	3.06	1.49	1.44
3	D	9	TDR	C4-N3	-2.31	1.34	1.38

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	9	TDR	N1-C2-N3	5.77	121.25	115.17
3	D	9	TDR	C4-N3-C2	-3.75	121.20	126.37
3	D	9	TDR	CM5-C5-C4	3.65	122.68	118.78
3	D	9	TDR	C5-C4-N3	3.56	118.42	115.32
3	D	9	TDR	O2-C2-N1	-3.18	119.51	122.79
3	D	9	TDR	CM5-C5-C6	-3.13	120.00	123.23
3	D	9	TDR	C5-C6-N1	-2.86	119.34	122.42
3	D	9	TDR	O4-C4-C5	-2.39	122.18	124.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	C	7/8 (87%)	-0.70	0 100 100	39, 40, 50, 68	0
1	D	7/8 (87%)	-0.63	0 100 100	40, 44, 61, 78	0
2	A	634/659 (96%)	0.02	17 (2%) 56 51	34, 56, 109, 152	0
2	B	631/659 (95%)	-0.01	19 (3%) 52 48	35, 61, 102, 128	0
All	All	1279/1334 (95%)	-0.00	36 (2%) 55 50	34, 58, 105, 152	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	658	HIS	5.4
2	A	656	HIS	5.1
2	B	618	VAL	5.1
2	A	264	HIS	4.4
2	B	611	VAL	4.3
2	A	478	GLY	3.7
2	A	657	HIS	3.5
2	B	617	ILE	3.4
2	A	245	ALA	3.4
2	A	261	LEU	3.3
2	B	614	PHE	3.0
2	B	651	ALA	2.9
2	B	515	ILE	2.9
2	A	455	ARG	2.9
2	B	379	ASP	2.8
2	B	271	GLY	2.7
2	B	519	THR	2.7
2	B	657	HIS	2.7
2	B	654	HIS	2.7
2	B	93	HIS	2.6
2	A	217	VAL	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	A	617	ILE	2.5
2	A	258	HIS	2.5
2	A	655	HIS	2.5
2	A	659	HIS	2.5
2	A	268	ARG	2.5
2	A	647	VAL	2.3
2	B	231	PRO	2.3
2	A	654	HIS	2.3
2	B	518	ALA	2.3
2	B	517	ARG	2.3
2	B	656	HIS	2.2
2	A	508	GLN	2.2
2	A	423	CYS	2.2
2	B	646	GLU	2.0
2	B	652	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	TDR	D	9	9/9	0.74	0.17	107,108,108,108	0
4	CL	A	801	1/1	0.96	0.06	52,52,52,52	0
4	CL	A	800	1/1	0.99	0.04	49,49,49,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.