



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 04:20 PM UTC

PDB ID : 2OUV / pdb_00002ouv
Title : crystal structure of pde10a2 mutant of D564N
Authors : Wang, H.C.; Liu, Y.D.; Hou, J.; Zheng, M.Y.; Robinson, H.
Deposited on : 2007-02-12
Resolution : 1.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	<div> <div>11%</div> <div>82%</div> <div>15%</div> <div>..</div> </div>
1	B	331	<div> <div>12%</div> <div>83%</div> <div>15%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ZN	B	777	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5752 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	325	Total	C	N	O	S	0	0	0
			2636	1682	449	480	25			
1	B	328	Total	C	N	O	S	0	0	0
			2654	1692	452	485	25			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	446	SER	-	cloning artifact	UNP Q9Y233
A	447	HIS	-	cloning artifact	UNP Q9Y233
A	448	MET	-	cloning artifact	UNP Q9Y233
A	564	ASN	ASP	engineered mutation	UNP Q9Y233
B	446	SER	-	cloning artifact	UNP Q9Y233
B	447	HIS	-	cloning artifact	UNP Q9Y233
B	448	MET	-	cloning artifact	UNP Q9Y233
B	564	ASN	ASP	engineered mutation	UNP Q9Y233

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	252	Total 252	O 252	0	0
4	B	206	Total 206	O 206	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.44Å 82.20Å 155.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.56 30.00 – 1.56	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-1.56) 94.8 (30.00-1.56)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 1.56Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.209 , 0.228 0.209 , 0.229	Depositor DCC
R_{free} test set	9155 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	16.1	Xtriage
Anisotropy	0.522	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 31.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5752	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/2701	0.82	7/3657 (0.2%)
1	B	0.35	0/2719	0.80	6/3682 (0.2%)
All	All	0.34	0/5420	0.81	13/7339 (0.2%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	563	HIS	N-CA-C	7.07	121.01	112.38
1	B	676	CYS	N-CA-C	6.47	120.63	112.87
1	B	545	HIS	N-CA-C	6.16	118.80	111.71
1	B	563	HIS	N-CA-C	6.13	119.86	112.38
1	B	734	ALA	N-CA-C	6.12	117.62	111.07
1	A	734	ALA	N-CA-C	5.95	117.45	110.97
1	B	657	ASN	N-CA-C	-5.84	106.15	113.28
1	A	483	GLU	N-CA-C	5.70	117.95	111.11
1	A	545	HIS	N-CA-C	5.49	120.36	111.37
1	B	746	PRO	N-CA-C	5.47	117.37	110.70
1	A	519	TYR	N-CA-C	-5.37	100.96	109.76
1	A	676	CYS	N-CA-C	5.31	119.24	112.87
1	A	590	THR	N-CA-C	5.30	117.49	111.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2636	0	2597	30	0
1	B	2654	0	2614	37	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	252	0	0	0	0
4	B	206	0	0	3	0
All	All	5752	0	5211	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:646:GLU:HG2	1:B:650:GLN:HE21	1.36	0.89
1:B:709:GLN:HE21	1:B:709:GLN:HA	1.41	0.84
1:B:460:MET:HE3	1:B:505:LEU:HD23	1.64	0.79
1:B:641:ASN:HD22	1:B:664:ARG:HH11	1.40	0.68
1:B:644:GLN:O	1:B:648:MET:HG3	1.94	0.68
1:A:655:ASN:ND2	1:A:657:ASN:OD1	2.31	0.63
1:B:646:GLU:HG2	1:B:650:GLN:NE2	2.12	0.63
1:A:658:ASN:HD22	1:A:658:ASN:C	2.08	0.62
1:B:689:ALA:HB2	1:B:773:THR:HG21	1.80	0.62
1:B:724:GLN:HG3	4:B:895:HOH:O	2.00	0.60
1:A:578:PHE:CD2	1:A:706:LEU:HD13	2.37	0.59
1:A:709:GLN:NE2	1:A:710:PRO:HD2	2.17	0.59
1:A:453:SER:HA	1:A:456:TRP:NE1	2.17	0.58
1:B:483:GLU:HA	1:B:486:TRP:CE2	2.38	0.57
1:B:617:SER:O	1:B:621:GLN:HG3	2.05	0.57
1:B:655:ASN:ND2	1:B:657:ASN:H	2.02	0.57
1:A:573:SER:O	1:A:577:LYS:HG3	2.05	0.56
1:A:616:SER:O	1:A:620:GLU:HG3	2.06	0.56
1:A:483:GLU:HA	1:A:486:TRP:CE2	2.42	0.55
1:A:448:MET:HE2	1:A:602:ILE:HG13	1.89	0.53
1:B:739:THR:O	1:B:743:GLN:HG3	2.08	0.53
1:B:483:GLU:HG3	1:B:486:TRP:CZ3	2.44	0.53
1:B:689:ALA:CB	1:B:773:THR:HG21	2.39	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:468:LEU:O	1:A:472:ILE:HG13	2.09	0.52
1:B:448:MET:HE2	1:B:602:ILE:HG13	1.92	0.52
1:B:659:GLN:HG3	4:B:853:HOH:O	2.09	0.51
1:B:460:MET:HE3	1:B:505:LEU:CD2	2.38	0.51
1:B:643:LYS:O	1:B:647:GLU:HG3	2.11	0.51
1:B:735:ILE:HB	1:B:736:PRO:HD3	1.94	0.50
1:B:549:THR:OG1	1:B:552:GLU:HG3	2.11	0.50
1:B:532:THR:HG22	1:B:676:CYS:SG	2.52	0.50
1:B:453:SER:HA	1:B:456:TRP:NE1	2.27	0.50
1:A:456:TRP:CZ3	1:A:506:GLU:HG3	2.45	0.49
1:B:655:ASN:C	1:B:655:ASN:HD22	2.19	0.49
1:A:459:LEU:HD13	1:A:487:PRO:HB2	1.95	0.49
1:B:655:ASN:HD22	1:B:656:LEU:N	2.10	0.49
1:A:600:VAL:O	1:A:604:GLN:HG3	2.12	0.49
1:A:644:GLN:O	1:A:648:MET:HG3	2.14	0.48
1:B:624:GLU:HG2	1:B:628:LYS:HE3	1.96	0.47
1:B:502:CYS:SG	1:B:554:LYS:HE2	2.54	0.47
1:A:453:SER:O	1:A:457:GLN:HG3	2.14	0.47
1:A:682:TRP:HB3	1:A:683:PRO:HD3	1.97	0.46
1:A:735:ILE:HB	1:A:736:PRO:HD3	1.98	0.46
1:B:529:HIS:HD2	4:B:810:HOH:O	1.98	0.45
1:B:709:GLN:HA	1:B:709:GLN:NE2	2.21	0.45
1:A:483:GLU:HG3	1:A:486:TRP:CZ3	2.53	0.44
1:A:484:ASN:HD22	1:A:484:ASN:H	1.65	0.44
1:B:460:MET:HG2	1:B:505:LEU:HG	1.99	0.44
1:A:696:PHE:CE2	1:A:713:MET:HE3	2.53	0.43
1:B:624:GLU:CG	1:B:628:LYS:HE3	2.49	0.43
1:A:453:SER:HB2	1:A:457:GLN:HE21	1.83	0.43
1:A:739:THR:O	1:A:743:GLN:HG3	2.18	0.43
1:A:573:SER:HB3	1:A:577:LYS:HE3	2.00	0.43
1:B:453:SER:HA	1:B:456:TRP:CD1	2.54	0.43
1:A:549:THR:OG1	1:A:552:GLU:HG3	2.19	0.43
1:B:619:TYR:O	1:B:622:VAL:HG12	2.18	0.42
1:A:771:THR:HA	1:B:713:MET:HB3	2.00	0.42
1:B:658:ASN:OD1	1:B:660:SER:HB3	2.19	0.42
1:A:675:LEU:O	1:A:678:VAL:HG22	2.20	0.42
1:A:456:TRP:O	1:A:460:MET:HG3	2.20	0.41
1:B:700:GLY:HA3	1:B:714:MET:O	2.20	0.41
1:A:692:ILE:HG22	1:A:696:PHE:CE2	2.55	0.41
1:A:764:LYS:HB3	1:A:764:LYS:NZ	2.35	0.41
1:B:655:ASN:ND2	1:B:655:ASN:C	2.78	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:462:PHE:CD2	1:A:488:GLY:HA3	2.57	0.40
1:B:462:PHE:CD2	1:B:488:GLY:HA3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	323/331 (98%)	316 (98%)	7 (2%)	0	100	100
1	B	326/331 (98%)	321 (98%)	5 (2%)	0	100	100
All	All	649/662 (98%)	637 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/299 (98%)	291 (99%)	3 (1%)	68	47
1	B	296/299 (99%)	294 (99%)	2 (1%)	76	59
All	All	590/598 (99%)	585 (99%)	5 (1%)	73	56

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	484	ASN
1	A	564	ASN
1	A	658	ASN
1	B	547	LEU
1	B	709	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	457	GLN
1	A	484	ASN
1	A	604	GLN
1	A	655	ASN
1	A	658	ASN
1	A	690	ASN
1	A	709	GLN
1	B	461	GLN
1	B	576	GLN
1	B	593	GLN
1	B	609	ASN
1	B	641	ASN
1	B	650	GLN
1	B	655	ASN
1	B	690	ASN
1	B	709	GLN
1	B	743	GLN
1	B	761	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	325/331 (98%)	0.57	36 (11%)	10 12	11, 17, 32, 36	0
1	B	328/331 (99%)	0.76	40 (12%)	8 10	11, 19, 32, 36	0
All	All	653/662 (98%)	0.66	76 (11%)	9 11	11, 18, 32, 36	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	773	THR	5.2
1	B	502	CYS	4.7
1	B	772	ALA	4.6
1	B	456	TRP	4.5
1	A	657	ASN	4.4
1	A	771	THR	4.3
1	A	462	PHE	4.0
1	A	577	LYS	3.9
1	A	502	CYS	3.9
1	A	447	HIS	3.8
1	B	770	GLU	3.7
1	A	501	SER	3.7
1	A	471	GLU	3.6
1	A	463	THR	3.6
1	B	585	LEU	3.5
1	A	466	VAL	3.5
1	A	521	ARG	3.4
1	A	456	TRP	3.4
1	A	647	GLU	3.3
1	B	613	THR	3.2
1	B	588	THR	3.2
1	B	584	ALA	3.2
1	B	501	SER	3.1
1	A	461	GLN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	650	GLN	3.0
1	A	504	GLU	3.0
1	A	459	LEU	3.0
1	A	464	LEU	2.9
1	A	606	GLU	2.9
1	A	770	GLU	2.9
1	B	649	TYR	2.9
1	B	647	GLU	2.9
1	B	586	TYR	2.9
1	B	498	CYS	2.9
1	B	579	ASP	2.8
1	B	721	GLU	2.8
1	A	578	PHE	2.8
1	A	460	MET	2.8
1	A	469	CYS	2.8
1	B	617	SER	2.7
1	A	470	LYS	2.7
1	B	616	SER	2.7
1	B	461	GLN	2.7
1	A	454	GLU	2.6
1	B	749	GLU	2.6
1	B	621	GLN	2.6
1	B	521	ARG	2.6
1	B	551	LEU	2.6
1	B	618	GLU	2.6
1	B	659	GLN	2.6
1	A	458	GLY	2.6
1	A	550	ASP	2.6
1	A	467	ARG	2.6
1	A	656	LEU	2.5
1	A	651	THR	2.5
1	A	472	ILE	2.5
1	B	578	PHE	2.5
1	A	453	SER	2.5
1	B	470	LYS	2.5
1	B	553	ARG	2.5
1	B	767	ARG	2.5
1	B	545	HIS	2.4
1	A	649	TYR	2.4
1	B	658	ASN	2.4
1	B	446	SER	2.3
1	B	652	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	650	GLN	2.3
1	A	506	GLU	2.2
1	B	453	SER	2.2
1	B	651	THR	2.2
1	B	484	ASN	2.2
1	B	609	ASN	2.2
1	A	500	THR	2.1
1	A	709	GLN	2.0
1	B	504	GLU	2.0
1	B	606	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	A	778	1/1	0.73	0.32	24,24,24,24	0
2	ZN	B	777	1/1	0.76	0.59	30,30,30,30	0
2	ZN	A	777	1/1	0.99	0.28	22,22,22,22	0
3	MG	B	778	1/1	0.99	0.03	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.