



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:26 PM UTC

PDB ID : 2QK9 / pdb_00002qk9
Title : Human RNase H catalytic domain mutant D210N in complex with 18-mer RNA/DNA hybrid
Authors : Nowotny, M.; Gaidamakov, S.A.; Ghirlando, R.; Cerritelli, S.M.; Crouch, R.J.; Yang, W.
Deposited on : 2007-07-10
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)
RNA backbone	3983	1112 (2.80-2.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	18	 78% 17% 6%
2	C	18	 39% 44% 17%
3	A	154	 5% 70% 24% ...

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 2112 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*AP*GP*UP*GP*CP*GP*AP*CP*AP*CP*CP*UP*GP*AP*UP*UP*CP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	18	Total	C	N	O	P	0	0	0
			377	170	66	124	17			

- Molecule 2 is a DNA chain called 5'-D(*GP*GP*AP*AP*TP*CP*AP*GP*GP*TP*GP*TP*CP*GP*CP*AP*CP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	18	Total	C	N	O	P	0	0	0
			369	176	70	106	17			

- Molecule 3 is a protein called Ribonuclease H1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	153	Total	C	N	O	S	0	0	0
			1188	739	224	218	7			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	133	GLY	-	expression tag	UNP O60930
A	134	SER	-	expression tag	UNP O60930
A	135	HIS	-	expression tag	UNP O60930
A	210	ASN	ASP	engineered mutation	UNP O60930

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is HEXANE-1,6-DIAMINE (CCD ID: 16D) (formula: C₆H₁₆N₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	N	0	0
			8	6	2		

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Na	0	0
			1	1		

- Molecule 7 is CITRATE ANION (CCD ID: FLC) (formula: C₆H₅O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	22	Total	O	0	0
			22	22		
9	C	26	Total	O	0	0
			26	26		
9	A	61	Total	O	0	0
			61	61		


● Molecule 1: 5'-R(*AP*GP*UP*GP*CP*GP*AP*CP*AP*CP*CP*UP*GP*AP*UP*UP*CP*C)-3'



● Molecule 2: 5'-D(*GP*GP*AP*AP*TP*CP*AP*GP*GP*TP*GP*TP*CP*GP*CP*AP*CP*T)-3'



- Molecule 3: Ribonuclease H1



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	158.58Å 158.58Å 142.06Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.55 30.00 – 2.55	Depositor EDS
% Data completeness (in resolution range)	94.0 (30.00-2.55) 93.9 (30.00-2.55)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.29Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.190 , 0.216 0.186 , 0.214	Depositor DCC
R_{free} test set	2085 reflections (7.77%)	wwPDB-VP
Wilson B-factor (Å ²)	41.1	Xtriage
Anisotropy	0.537	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2112	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 16D, GOL, NA, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.60	0/420	0.96	1/652 (0.2%)
2	C	0.50	0/414	1.02	3/638 (0.5%)
3	A	0.88	1/1213 (0.1%)	1.18	9/1636 (0.6%)
All	All	0.77	1/2047 (0.0%)	1.10	13/2926 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	136	MET	SD-CE	5.68	1.93	1.79

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	29	DG	C5'-C4'-O4'	-8.47	96.70	109.40
3	A	239	ILE	N-CA-C	7.97	118.75	110.62
3	A	135	HIS	N-CA-C	6.75	125.19	110.80
2	C	25	DA	C5'-C4'-O4'	-5.87	100.60	109.40
1	B	14	A	C5'-C4'-C3'	-5.70	107.45	116.00
2	C	32	DG	C2'-C3'-O3'	-5.69	102.97	111.50
3	A	154	ARG	N-CA-C	5.63	117.42	111.28
3	A	180	GLN	N-CA-C	5.63	117.91	107.99
3	A	178	GLY	N-CA-C	5.55	122.29	112.53
3	A	261	VAL	CA-C-N	5.18	125.47	119.93
3	A	261	VAL	C-N-CA	5.18	125.47	119.93
3	A	173	GLY	N-CA-C	-5.14	100.99	113.18
3	A	210	ASN	N-CA-C	-5.09	106.91	113.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	377	0	196	4	0
2	C	369	0	204	20	0
3	A	1188	0	1154	37	0
4	A	20	0	0	0	0
4	B	10	0	0	0	0
4	C	5	0	0	0	0
5	C	8	0	16	5	0
6	A	1	0	0	0	0
7	A	13	0	5	1	0
8	A	12	0	16	0	0
9	A	61	0	0	6	0
9	B	22	0	0	1	0
9	C	26	0	0	1	0
All	All	2112	0	1591	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:220:ASN:C	3:A:220:ASN:HD22	1.63	1.03
2:C:25:DA:H8	2:C:25:DA:H5'	1.26	0.98
3:A:186:GLU:HG2	3:A:211:SER:HB2	1.50	0.93
2:C:24:DC:C2'	2:C:25:DA:H5''	2.01	0.90
3:A:232:THR:CG2	3:A:234:ALA:H	1.86	0.89
3:A:186:GLU:CG	3:A:211:SER:HB2	2.04	0.87
2:C:25:DA:H5'	2:C:25:DA:C8	2.13	0.83
2:C:35:DC:H5'	2:C:35:DC:H6	1.47	0.79
2:C:24:DC:H2''	2:C:25:DA:H5''	1.65	0.77
3:A:230:TRP:CD1	3:A:241:LYS:HE3	2.21	0.75
3:A:136:MET:HE3	3:A:137:GLY:N	2.02	0.75
3:A:136:MET:HE3	3:A:137:GLY:H	1.52	0.74
7:A:1001:FLC:OA1	9:A:9021:HOH:O	2.06	0.74
3:A:232:THR:HG21	9:A:9057:HOH:O	1.88	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:220:ASN:C	3:A:220:ASN:ND2	2.36	0.72
3:A:232:THR:HG23	3:A:234:ALA:H	1.52	0.72
3:A:232:THR:HG22	3:A:234:ALA:H	1.53	0.72
2:C:24:DC:H2'	2:C:25:DA:H5''	1.74	0.70
1:B:1:A:H2'	1:B:2:G:C8	2.29	0.68
1:B:7:A:N6	5:C:1004:16D:H41	2.10	0.67
5:C:1004:16D:H52	9:C:1033:HOH:O	1.95	0.66
2:C:28:DT:H2''	2:C:29:DG:H5'	1.78	0.66
3:A:217:GLY:O	3:A:222:VAL:HG23	1.96	0.65
2:C:35:DC:H5'	2:C:35:DC:C6	2.31	0.65
2:C:35:DC:H2'	2:C:36:DT:H71	1.78	0.63
3:A:186:GLU:HG2	3:A:211:SER:CB	2.28	0.61
2:C:24:DC:H2'	2:C:25:DA:C5'	2.31	0.60
3:A:271:GLU:HG2	9:A:9053:HOH:O	2.03	0.58
2:C:24:DC:C2'	2:C:25:DA:C5'	2.79	0.58
1:B:7:A:H62	5:C:1004:16D:H21	1.69	0.58
1:B:1:A:H2'	1:B:2:G:H8	1.72	0.55
3:A:232:THR:HG22	3:A:234:ALA:N	2.20	0.54
3:A:186:GLU:HG3	3:A:211:SER:HB2	1.88	0.51
3:A:232:THR:HB	3:A:236:LYS:O	2.10	0.51
3:A:172:VAL:HB	9:A:9021:HOH:O	2.12	0.48
2:C:19:DG:H5''	2:C:19:DG:C8	2.49	0.48
3:A:207:LEU:C	3:A:207:LEU:HD23	2.39	0.48
9:B:1027:HOH:O	5:C:1004:16D:H12	2.15	0.47
2:C:31:DC:H2'	2:C:32:DG:C8	2.50	0.47
3:A:232:THR:HG23	3:A:233:SER:N	2.29	0.46
3:A:168:HIS:CE1	3:A:170:LEU:HB2	2.51	0.46
3:A:135:HIS:O	3:A:136:MET:HB3	2.16	0.46
2:C:28:DT:C4	5:C:1004:16D:H62	2.51	0.45
2:C:32:DG:H2''	2:C:33:DC:H5'	1.98	0.44
3:A:156:PRO:HG2	3:A:180:GLN:HB2	1.99	0.44
3:A:274:ASP:CG	3:A:278:ARG:HH21	2.25	0.44
2:C:28:DT:H2''	2:C:29:DG:C5'	2.45	0.43
3:A:177:PRO:HA	3:A:285:GLU:OE2	2.19	0.43
3:A:140:VAL:HG11	3:A:202:ILE:HD13	2.01	0.43
3:A:174:ILE:HG22	3:A:175:ARG:O	2.19	0.42
3:A:174:ILE:HG22	3:A:175:ARG:N	2.33	0.42
3:A:172:VAL:CB	9:A:9021:HOH:O	2.67	0.42
3:A:169:PRO:HD3	9:A:9004:HOH:O	2.18	0.42
2:C:32:DG:H2''	2:C:33:DC:C5'	2.50	0.42
3:A:136:MET:CE	3:A:137:GLY:N	2.80	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:230:TRP:CG	3:A:241:LYS:HE3	2.54	0.41
2:C:25:DA:C8	2:C:25:DA:C5'	2.94	0.41
3:A:182:ASN:OD1	3:A:182:ASN:N	2.53	0.41
2:C:32:DG:H2'	2:C:33:DC:C6	2.56	0.41
2:C:25:DA:H2'	2:C:26:DG:O4'	2.20	0.41
3:A:217:GLY:HA3	3:A:244:PHE:CE1	2.56	0.40
3:A:285:GLU:HG3	3:A:286:ASP:H	1.86	0.40
3:A:203:ASN:ND2	3:A:203:ASN:C	2.80	0.40
3:A:261:VAL:HA	3:A:262:PRO:HD3	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	151/154 (98%)	145 (96%)	3 (2%)	3 (2%)	6 6

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	135	HIS
3	A	136	MET
3	A	220	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	121/124 (98%)	113 (93%)	8 (7%)	15 21

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	138	ASP
3	A	140	VAL
3	A	150	SER
3	A	170	LEU
3	A	199	THR
3	A	220	ASN
3	A	232	THR
3	A	276	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	203	ASN
3	A	220	ASN
3	A	223	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	B	17/18 (94%)	1 (5%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	B	14	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	1006	-	4,4,4	0.40	0	6,6,6	0.30	0
8	GOL	A	1002	-	5,5,5	0.46	0	5,5,5	0.44	0
7	FLC	A	1001	-	12,12,12	1.51	2 (16%)	17,17,17	1.51	1 (5%)
4	SO4	A	1011	-	4,4,4	0.43	0	6,6,6	0.28	0
4	SO4	C	1009	-	4,4,4	0.48	0	6,6,6	0.18	0
8	GOL	A	1003	-	5,5,5	0.56	0	5,5,5	0.43	0
4	SO4	A	1010	-	4,4,4	0.47	0	6,6,6	0.17	0
4	SO4	B	1008	-	4,4,4	0.41	0	6,6,6	0.10	0
4	SO4	B	1007	-	4,4,4	0.37	0	6,6,6	0.14	0
4	SO4	A	1005	-	4,4,4	0.43	0	6,6,6	0.45	0
5	16D	C	1004	-	7,7,7	0.62	0	6,6,6	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	A	1003	-	-	0/4/4/4	-
8	GOL	A	1002	-	-	0/4/4/4	-
5	16D	C	1004	-	-	0/5/5/5	-
7	FLC	A	1001	-	-	0/16/16/16	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1001	FLC	OA2-CAC	-2.89	1.21	1.30
7	A	1001	FLC	OG2-CGC	-2.74	1.21	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	1001	FLC	OB2-CBC-CB	4.39	121.57	113.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1001	FLC	1	0
5	C	1004	16D	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	B	18/18 (100%)	-0.66	0	100	100	27, 39, 53, 55	0
2	C	18/18 (100%)	-0.52	0	100	100	33, 39, 59, 64	0
3	A	153/154 (99%)	0.01	7 (4%)	37	38	25, 39, 62, 99	5 (3%)
All	All	189/190 (99%)	-0.11	7 (3%)	45	46	25, 39, 59, 99	5 (2%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	134	SER	10.6
3	A	135	HIS	10.1
3	A	136	MET	7.4
3	A	286	ASP	4.4
3	A	227	LYS	2.2
3	A	223	GLN	2.2
3	A	137	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	A	1010	5/5	0.72	0.25	126,127,127,128	0
5	16D	C	1004	8/8	0.81	0.36	67,72,73,73	0
4	SO4	B	1008	5/5	0.82	0.18	140,140,140,141	0
4	SO4	C	1009	5/5	0.82	0.29	121,122,122,122	0
8	GOL	A	1002	6/6	0.84	0.23	72,74,78,81	0
8	GOL	A	1003	6/6	0.84	0.21	70,72,74,74	0
4	SO4	B	1007	5/5	0.88	0.28	134,134,134,134	0
4	SO4	A	1005	5/5	0.88	0.13	81,82,83,83	0
7	FLC	A	1001	13/13	0.92	0.20	72,75,76,78	13
4	SO4	A	1011	5/5	0.93	0.10	73,74,76,76	0
6	NA	A	9001	1/1	0.94	0.12	50,50,50,50	0
4	SO4	A	1006	5/5	0.94	0.14	70,70,71,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.