



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:08 AM UTC

PDB ID : 2V52 / pdb_00002v52
Title : Structure of MAL-RPEL2 complexed to G-actin
Authors : Mouilleron, S.; Guettler, S.; Langer, C.A.; Treisman, R.; McDonald, N.Q.
Deposited on : 2008-10-01
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	377	
2	M	32	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3738 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ACTIN, ALPHA SKELETAL MUSCLE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	360	Total	C	N	O	S	0	35	0
			3003	1921	486	572	24			

- Molecule 2 is a protein called MKL/MYOCARDIN-LIKE PROTEIN 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	30	Total	C	N	O	S	0	7	0
			305	189	69	46	1			

- Molecule 3 is LATRUNCULIN B (CCD ID: LAB) (formula: C₂₀H₂₉NO₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			27	20	1	5	1		

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Mg 1 1	0	0

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	334	Total	O	0	2
			334	334		
7	M	25	Total	O	0	0
			25	25		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ACTIN, ALPHA SKELETAL MUSCLE



• Molecule 2: MKL/MYOCARDIN-LIKE PROTEIN 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.75Å 55.44Å 138.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.35 – 1.45 29.35 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.3 (29.35-1.45) 99.3 (29.35-1.45)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 1.45Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.147 , 0.188 0.143 , 0.183	Depositor DCC
R_{free} test set	3769 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	17.8	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 58.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.024 for k,h,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	3738	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LAB, MG, GOL, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	1.64	79/3138 (2.5%)	1.13	18/4245 (0.4%)
2	M	1.66	9/319 (2.8%)	0.99	1/420 (0.2%)
All	All	1.64	88/3457 (2.5%)	1.12	19/4665 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	1	2

All (88) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	149	THR	C-O	-17.05	1.04	1.24
1	B	148	THR	C-O	-12.98	1.06	1.24
1	B	192	ILE	C-O	-11.96	1.08	1.24
1	B	197	GLY	C-O	-11.49	1.08	1.24
1	B	371	HIS	C-O	-11.24	1.09	1.24
1	B	87	HIS	C-O	-10.78	1.11	1.24
1	B	230	ALA	CA-CB	-10.65	1.36	1.53
1	B	100	GLU	C-O	-10.43	1.10	1.24
1	B	4	GLU	C-O	-10.38	1.11	1.24
1	B	373	LYS	C-O	-9.80	1.11	1.24
1	B	147	ARG	C-O	-9.62	1.11	1.23
1	B	53	TYR	C-O	-9.52	1.13	1.24
1	B	234	SER	C-O	-9.45	1.12	1.24
2	M	134	MET	SD-CE	-9.20	1.56	1.79
1	B	193	LEU	C-O	-9.05	1.12	1.24
1	B	234	SER	CA-C	-9.04	1.40	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	235	SER	N-CA	-8.95	1.34	1.46
1	B	370	VAL	C-O	-8.85	1.13	1.24
1	B	232	SER	C-O	-8.83	1.12	1.24
1	B	194	THR	C-O	-8.80	1.13	1.24
1	B	99	GLU	C-O	-8.44	1.13	1.24
1	B	198	TYR	CD1-CE1	-8.36	1.13	1.38
1	B	178[A]	LEU	C-O	-8.34	1.14	1.24
1	B	178[B]	LEU	C-O	-8.34	1.14	1.24
1	B	197	GLY	N-CA	-8.31	1.32	1.45
1	B	236	LEU	C-O	-8.18	1.14	1.24
1	B	57	GLU	C-O	-8.12	1.13	1.24
1	B	196	ARG	NE-CZ	-7.89	1.24	1.33
1	B	56	ASP	C-O	-7.87	1.13	1.24
1	B	59	GLN	C-O	-7.85	1.14	1.24
1	B	196	ARG	CD-NE	-7.71	1.35	1.46
2	M	130[A]	GLU	C-O	-7.61	1.15	1.24
2	M	130[B]	GLU	C-O	-7.61	1.15	1.24
1	B	231	ALA	N-CA	-7.49	1.36	1.46
1	B	195	GLU	CD-OE1	-7.47	1.11	1.25
1	B	198	TYR	C-O	-7.45	1.14	1.24
1	B	372	ARG	N-CA	-7.27	1.36	1.46
1	B	235	SER	C-O	-7.24	1.14	1.24
2	M	134	MET	C-O	-7.14	1.14	1.24
1	B	372	ARG	C-O	-7.05	1.14	1.24
1	B	99	GLU	CB-CG	-7.04	1.31	1.52
1	B	58	ALA	CA-CB	-6.90	1.41	1.53
1	B	235	SER	CA-C	-6.84	1.42	1.52
1	B	194	THR	CA-CB	-6.82	1.41	1.53
2	M	135	HIS	C-O	-6.74	1.14	1.23
1	B	270	GLU	C-O	-6.72	1.15	1.24
1	B	234	SER	C-N	-6.70	1.24	1.33
1	B	372	ARG	CB-CG	-6.66	1.32	1.52
1	B	270	GLU	C-N	-6.57	1.25	1.33
1	B	234	SER	N-CA	-6.34	1.38	1.46
1	B	232	SER	CA-C	-6.27	1.44	1.52
1	B	237	GLU	CD-OE1	-6.02	1.14	1.25
2	M	121[A]	LYS	C-O	-6.00	1.17	1.24
2	M	121[B]	LYS	C-O	-6.00	1.17	1.24
1	B	196	ARG	CZ-NH1	-5.97	1.24	1.32
1	B	198	TYR	CA-C	-5.96	1.45	1.52
1	B	99	GLU	CD-OE1	-5.86	1.14	1.25
1	B	145[A]	SER	C-O	-5.79	1.16	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	145[B]	SER	C-O	-5.79	1.16	1.24
1	B	230	ALA	N-CA	-5.78	1.39	1.46
2	M	127	GLU	C-O	-5.78	1.16	1.23
1	B	230	ALA	CA-C	-5.74	1.45	1.52
1	B	234	SER	CB-OG	-5.69	1.30	1.42
1	B	3	ASP	N-CA	-5.62	1.35	1.46
1	B	99	GLU	CD-OE2	-5.59	1.14	1.25
1	B	50	LYS	CA-CB	-5.57	1.42	1.53
1	B	196	ARG	CZ-NH2	-5.56	1.26	1.33
1	B	194	THR	CB-CG2	-5.55	1.34	1.52
1	B	53	TYR	N-CA	-5.53	1.39	1.46
1	B	99	GLU	C-N	-5.48	1.25	1.33
1	B	196	ARG	CA-C	-5.40	1.43	1.52
1	B	147	ARG	CZ-NH2	-5.39	1.26	1.33
1	B	58	ALA	C-O	-5.37	1.17	1.24
1	B	194	THR	N-CA	5.36	1.53	1.46
1	B	147	ARG	NE-CZ	-5.36	1.27	1.33
1	B	230	ALA	C-O	-5.28	1.17	1.24
1	B	198	TYR	CD2-CE2	-5.21	1.23	1.38
1	B	237	GLU	CA-C	-5.21	1.45	1.52
1	B	100	GLU	CD-OE2	-5.21	1.15	1.25
1	B	51	ASP	C-O	-5.19	1.17	1.24
1	B	59	GLN	CA-C	-5.19	1.46	1.52
1	B	87	HIS	CA-C	-5.09	1.46	1.52
1	B	231	ALA	C-O	-5.09	1.17	1.24
2	M	134	MET	C-N	-5.07	1.26	1.33
1	B	100	GLU	N-CA	-5.03	1.39	1.46
1	B	196	ARG	C-N	-5.02	1.26	1.33
1	B	52	SER	C-O	-5.01	1.17	1.23
1	B	51	ASP	CA-C	5.01	1.59	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	235	SER	N-CA-C	-10.29	100.27	112.92
1	B	196	ARG	NE-CZ-NH1	-9.55	111.95	121.50
1	B	230	ALA	N-CA-C	-8.42	101.33	111.69
1	B	195	GLU	N-CA-C	7.95	119.58	111.07
1	B	236	LEU	N-CA-C	7.14	119.15	111.36
1	B	59	GLN	N-CA-C	-6.75	103.16	111.40
1	B	194	THR	OG1-CB-CG2	6.71	122.72	109.30
1	B	196	ARG	CA-C-N	-6.39	112.32	122.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	196	ARG	C-N-CA	-6.39	112.32	122.10
2	M	114	THR	CB-CA-C	-6.27	96.28	110.07
1	B	372	ARG	NE-CZ-NH1	6.11	127.61	121.50
1	B	196	ARG	NE-CZ-NH2	6.00	124.60	119.20
1	B	196	ARG	CD-NE-CZ	5.93	132.71	124.40
1	B	57	GLU	CG-CD-OE1	5.73	131.57	118.40
1	B	230	ALA	CA-C-N	5.30	131.67	121.54
1	B	230	ALA	C-N-CA	5.30	131.67	121.54
1	B	4	GLU	N-CA-C	5.18	117.67	111.71
1	B	235	SER	CA-C-N	-5.14	112.99	120.29
1	B	235	SER	C-N-CA	-5.14	112.99	120.29

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	194	THR	CB

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	233	SER	Peptide
1	B	234	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3003	0	3032	49	0
2	M	305	0	332	13	0
3	B	27	0	29	1	0
4	B	31	0	12	0	0
5	B	1	0	0	0	0
6	B	12	0	16	0	0
7	B	334	0	0	11	1
7	M	25	0	0	1	0
All	All	3738	0	3421	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:111:ASN:HD21	1:B:119[B]:MET:HE1	1.01	1.17
2:M:123[B]:ARG:NH2	7:M:201:HOH:O	1.79	1.15
2:M:111[B]:ARG:HG2	2:M:111[B]:ARG:HH11	1.07	1.13
1:B:111:ASN:ND2	1:B:119[B]:MET:HE1	1.77	0.99
1:B:299:MET:CE	1:B:313[B]:MET:HG3	1.96	0.94
2:M:117:TYR:HD2	2:M:120[A]:ARG:HH22	1.05	0.93
1:B:299:MET:HE1	1:B:313[B]:MET:HG3	1.56	0.87
1:B:111:ASN:HD21	1:B:119[B]:MET:CE	1.88	0.85
2:M:111[B]:ARG:HG2	2:M:111[B]:ARG:NH1	1.83	0.85
2:M:117:TYR:HD2	2:M:120[A]:ARG:NH2	1.79	0.80
2:M:117:TYR:O	2:M:121[B]:LYS:HG2	1.83	0.79
2:M:111[B]:ARG:HH11	2:M:111[B]:ARG:CG	1.89	0.78
1:B:234:SER:HA	1:B:236:LEU:H	1.47	0.76
1:B:336[B]:LYS:HE2	7:B:633:HOH:O	1.85	0.75
1:B:68[B]:LYS:HD2	7:B:600:HOH:O	1.85	0.75
1:B:56:ASP:OD2	7:B:501:HOH:O	2.05	0.74
1:B:191:LYS:O	1:B:194:THR:HG22	1.90	0.71
2:M:119[B]:LYS:HD3	2:M:123[B]:ARG:NH2	2.07	0.68
1:B:276[B]:GLU:OE2	7:B:502:HOH:O	2.13	0.67
1:B:191:LYS:NZ	7:B:506:HOH:O	2.35	0.59
1:B:313[B]:MET:CE	7:B:578:HOH:O	2.49	0.59
1:B:276[A]:GLU:CD	7:B:529:HOH:O	2.45	0.58
1:B:230:ALA:C	1:B:232:SER:H	2.12	0.58
1:B:299:MET:HE2	1:B:313[B]:MET:HG3	1.84	0.58
1:B:252:ASN:C	1:B:252:ASN:HD22	2.12	0.57
1:B:115:ASN:C	1:B:119[B]:MET:HE3	2.29	0.57
1:B:272:ALA:HB1	1:B:276[B]:GLU:HB3	1.86	0.57
1:B:105[B]:LEU:HD13	1:B:132[B]:MET:HE2	1.87	0.55
1:B:252:ASN:HD22	1:B:253:GLU:N	2.06	0.53
1:B:56:ASP:OD1	1:B:59:GLN:OE1	2.29	0.51
1:B:180[B]:LEU:HD13	1:B:267:ILE:HD11	1.92	0.51
1:B:361:GLU:HB3	1:B:369:ILE:CD1	2.40	0.51
1:B:115:ASN:O	1:B:119[B]:MET:HE3	2.11	0.50
2:M:130[B]:GLU:HG2	2:M:134:MET:HE3	1.92	0.50
1:B:336[B]:LYS:HG2	7:B:633:HOH:O	2.11	0.50
1:B:234:SER:CA	1:B:236:LEU:H	2.18	0.50
1:B:291[A]:LYS:HE2	7:B:773:HOH:O	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:170:ALA:O	1:B:172:PRO:HD3	2.13	0.48
1:B:252:ASN:HD21	1:B:256:ARG:HH11	1.60	0.48
2:M:111[B]:ARG:NH1	2:M:111[B]:ARG:CG	2.58	0.48
1:B:124:PHE:CZ	1:B:132[A]:MET:HG3	2.49	0.48
1:B:351:THR:OG1	2:M:119[B]:LYS:HG3	2.13	0.48
1:B:109:PRO:HD3	7:B:539:HOH:O	2.13	0.48
1:B:176:MET:HE1	1:B:284:LYS:NZ	2.28	0.48
1:B:252:ASN:ND2	1:B:256:ARG:HH11	2.12	0.47
3:B:401:LAB:O2	3:B:401:LAB:H42	2.15	0.46
1:B:119[B]:MET:CE	7:B:784:HOH:O	2.64	0.45
1:B:250:ILE:HG13	1:B:253:GLU:HB2	2.00	0.44
2:M:120[A]:ARG:HB2	2:M:120[A]:ARG:CZ	2.48	0.44
1:B:8:LEU:HG	1:B:101:HIS:HB3	2.00	0.43
1:B:299:MET:HE1	1:B:313[B]:MET:CG	2.39	0.43
1:B:335:ARG:HA	1:B:338[A]:SER:OG	2.18	0.43
1:B:229:THR:O	1:B:232:SER:HB3	2.18	0.43
2:M:119[B]:LYS:HD3	2:M:123[B]:ARG:CZ	2.48	0.43
1:B:230:ALA:HB1	1:B:236:LEU:CD2	2.50	0.41
1:B:120:THR:HA	1:B:132[A]:MET:HE3	2.02	0.41
1:B:178[B]:LEU:HA	1:B:178[B]:LEU:HD23	1.68	0.41
1:B:131:ALA:HA	1:B:357:ILE:O	2.21	0.41
1:B:272:ALA:HB1	1:B:276[A]:GLU:HB2	2.03	0.41
1:B:305[B]:MET:O	1:B:306:TYR:C	2.64	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:765:HOH:O	7:B:766:HOH:O[3_445]	2.11	0.09

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	389/377 (103%)	379 (97%)	7 (2%)	3 (1%)	16	3
2	M	34/32 (106%)	34 (100%)	0	0	100	100
All	All	423/409 (103%)	413 (98%)	7 (2%)	3 (1%)	16	4

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	233	SER
1	B	234	SER
1	B	231	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	337/320 (105%)	332 (98%)	5 (2%)	57	25
2	M	31/29 (107%)	28 (90%)	3 (10%)	8	0
All	All	368/349 (105%)	360 (98%)	8 (2%)	51	13

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	233	SER
1	B	249[A]	THR
1	B	249[B]	THR
1	B	252	ASN
1	B	372	ARG
2	M	111[A]	ARG
2	M	111[B]	ARG
2	M	114	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	111	ASN
1	B	252	ASN
1	B	280	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	B	404	-	5,5,5	0.23	0	5,5,5	0.23	0
6	GOL	B	405	-	5,5,5	0.98	1 (20%)	5,5,5	1.87	2 (40%)
3	LAB	B	401	-	28,29,29	1.32	3 (10%)	29,41,41	1.92	6 (20%)
4	ATP	B	402	5	32,33,33	1.40	5 (15%)	48,52,52	1.54	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	B	404	-	-	0/4/4/4	-
6	GOL	B	405	-	-	1/4/4/4	-
3	LAB	B	401	-	-	3/21/49/49	0/2/3/3
4	ATP	B	402	5	-	3/22/38/38	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	LAB	O2-C1	4.05	1.43	1.34
4	B	402	ATP	C5-C4	3.58	1.45	1.39
4	B	402	ATP	C4-N9	-3.10	1.31	1.37
4	B	402	ATP	PB-O3A	2.55	1.62	1.59
4	B	402	ATP	C5-N7	-2.25	1.35	1.39
3	B	401	LAB	C18-S1	-2.25	1.73	1.77
3	B	401	LAB	C14-C15	2.16	1.55	1.52
6	B	405	GOL	O2-C2	-2.06	1.37	1.43
4	B	402	ATP	C8-N7	2.05	1.35	1.31

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	LAB	C17-S1-C18	4.80	94.82	92.04
4	B	402	ATP	C5-C4-N3	-4.56	120.44	126.72
3	B	401	LAB	C4-C5-C6	-3.65	104.64	112.57
4	B	402	ATP	N3-C4-N9	3.40	132.94	127.17
4	B	402	ATP	N3-C2-N1	-3.18	123.76	128.58
3	B	401	LAB	C19-C3-C2	-3.10	113.73	122.90
4	B	402	ATP	C2-N3-C4	3.06	119.29	111.83
3	B	401	LAB	C19-C3-C4	2.98	120.41	115.23
3	B	401	LAB	O2-C1-C2	2.94	118.01	111.20
6	B	405	GOL	C3-C2-C1	2.57	121.22	111.80
4	B	402	ATP	C4-N9-C8	2.56	108.42	105.74
6	B	405	GOL	O2-C2-C3	-2.41	99.22	109.18
4	B	402	ATP	N9-C8-N7	-2.34	110.61	113.94
3	B	401	LAB	C9-C10-C11	-2.24	109.98	114.06
4	B	402	ATP	C4-C5-N7	-2.23	108.03	110.58
4	B	402	ATP	C2-N1-C6	2.17	122.29	118.73
4	B	402	ATP	O4'-C1'-N9	2.16	112.23	108.09
4	B	402	ATP	O3G-PG-O2G	2.05	115.50	107.80

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	401	LAB	O3-C15-C16-C17
4	B	402	ATP	PG-O3B-PB-O1B
3	B	401	LAB	O2-C1-C2-C3
6	B	405	GOL	O1-C1-C2-O2
4	B	402	ATP	PB-O3B-PG-O2G
3	B	401	LAB	O1-C1-C2-C3
4	B	402	ATP	PG-O3B-PB-O2B


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	LAB	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand LAB B 401





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	M	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	M	129:ALA	C	130[B]:GLU	N	1.19

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	360/377 (95%)	-0.27	13 (3%) 46 48	6, 19, 42, 62	42 (11%)
2	M	30/32 (93%)	0.42	2 (6%) 24 23	12, 23, 45, 58	10 (33%)
All	All	390/409 (95%)	-0.22	15 (3%) 44 47	6, 19, 42, 62	52 (13%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	232	SER	7.2
1	B	233	SER	6.2
1	B	234	SER	4.8
1	B	64	ILE	4.2
1	B	228	ALA	3.7
1	B	231	ALA	3.5
1	B	60	SER	2.8
1	B	53	TYR	2.5
2	M	117	TYR	2.4
1	B	229	THR	2.3
1	B	51	ASP	2.3
1	B	235	SER	2.3
2	M	140	THR	2.1
1	B	243	PRO	2.1
1	B	59	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	B	405	6/6	0.76	0.14	46,49,51,51	0
6	GOL	B	404	6/6	0.91	0.09	34,35,37,37	0
3	LAB	B	401	27/27	0.99	0.04	14,17,20,23	0
4	ATP	B	402	31/31	0.99	0.03	11,13,15,16	4
5	MG	B	403	1/1	1.00	0.01	13,13,13,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LAB B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.