



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 22, 2026 – 08:24 PM UTC

PDB ID : 3AHG / pdb_00003ahg
Title : H64A mutant of Phosphoketolase from Bifidobacterium Breve complexed with a tricyclic ring form of thiamine diphosphate
Authors : Suzuki, R.; Katayama, T.; Kim, B.-J.; Wakagi, T.; Shoun, H.; Ashida, H.; Yamamoto, K.; Fushinobu, S.
Deposited on : 2010-04-22
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	FAILED
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	FAILED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	845	 82% 11% • 5%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7265 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xylulose 5-phosphate/fructose 6-phosphate phosphoketolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	802	Total	C	N	O	S	0	0	0
			6374	4050	1083	1217	24			


There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP D6PAH1
A	-18	GLY	-	expression tag	UNP D6PAH1
A	-17	SER	-	expression tag	UNP D6PAH1
A	-16	SER	-	expression tag	UNP D6PAH1
A	-15	HIS	-	expression tag	UNP D6PAH1
A	-14	HIS	-	expression tag	UNP D6PAH1
A	-13	HIS	-	expression tag	UNP D6PAH1
A	-12	HIS	-	expression tag	UNP D6PAH1
A	-11	HIS	-	expression tag	UNP D6PAH1
A	-10	HIS	-	expression tag	UNP D6PAH1
A	-9	SER	-	expression tag	UNP D6PAH1
A	-8	SER	-	expression tag	UNP D6PAH1
A	-7	GLY	-	expression tag	UNP D6PAH1
A	-6	LEU	-	expression tag	UNP D6PAH1
A	-5	VAL	-	expression tag	UNP D6PAH1
A	-4	PRO	-	expression tag	UNP D6PAH1
A	-3	ARG	-	expression tag	UNP D6PAH1
A	-2	GLY	-	expression tag	UNP D6PAH1
A	-1	SER	-	expression tag	UNP D6PAH1
A	0	HIS	-	expression tag	UNP D6PAH1
A	64	ALA	HIS	engineered mutation	UNP D6PAH1

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 2-[(9aR)-2,7-dimethyl-9a,10-dihydro-5H-pyrimido[4,5-d][1,3]thiazolo[3,2-a]pyrimidin-8-yl]ethyl trihydrogen diphosphate (CCD ID: HTD) (formula: C₁₂H₁₈N₄O₇P₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	819	Total	O	0	0
			819	819		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylulose 5-phosphate/fructose 6-phosphate phosphoketolase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	175.03Å 175.03Å 163.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.50 – 1.90 33.50 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.1 (33.50-1.90) 98.1 (33.50-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.35 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.163 , 0.205 0.164 , 0.206	Depositor DCC
R_{free} test set	4849 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	24.4	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7265	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MG, NA, HTD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.70	45/6547 (0.7%)	1.32	24/8901 (0.3%)

All (45) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	180	ILE	CA-CB	-9.67	1.43	1.54
1	A	701	ALA	CA-CB	9.56	1.69	1.53
1	A	749	VAL	CA-CB	8.25	1.64	1.54
1	A	373	ALA	CA-CB	8.08	1.66	1.53
1	A	754	ARG	CD-NE	-7.80	1.35	1.46
1	A	488	ALA	CA-CB	7.77	1.65	1.53
1	A	779	ALA	CA-CB	7.68	1.65	1.53
1	A	587	ILE	CA-C	7.48	1.62	1.52
1	A	185	ALA	CA-CB	7.21	1.66	1.53
1	A	417	ALA	CA-CB	7.02	1.64	1.53
1	A	552	SER	CB-OG	6.77	1.55	1.42
1	A	754	ARG	N-CA	6.76	1.54	1.46
1	A	322	VAL	CA-CB	6.37	1.60	1.54
1	A	66	GLY	N-CA	6.27	1.53	1.45
1	A	707	PHE	CA-C	6.24	1.60	1.52
1	A	494	ARG	CD-NE	6.03	1.54	1.46
1	A	650	VAL	CA-C	6.02	1.59	1.52
1	A	584	LEU	N-CA	5.94	1.53	1.46
1	A	363	PHE	N-CA	5.79	1.53	1.46
1	A	700	THR	N-CA	5.71	1.53	1.46
1	A	68	THR	C-O	-5.64	1.19	1.24
1	A	5	VAL	N-CA	5.52	1.56	1.46
1	A	732	VAL	CA-C	5.52	1.59	1.52
1	A	183	GLY	N-CA	5.38	1.52	1.45
1	A	692	ASP	N-CA	5.36	1.52	1.46
1	A	505	VAL	N-CA	-5.33	1.39	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	61	LEU	N-CA	5.31	1.53	1.46
1	A	742	THR	CA-CB	5.30	1.61	1.53
1	A	164	ALA	C-O	5.23	1.30	1.24
1	A	735	LYS	CA-CB	5.23	1.60	1.53
1	A	759	ALA	CA-CB	5.22	1.61	1.53
1	A	787	PHE	N-CA	-5.18	1.40	1.46
1	A	563	ILE	CA-CB	5.17	1.63	1.54
1	A	553	HIS	CA-C	5.14	1.59	1.52
1	A	570	ASP	CA-C	-5.13	1.46	1.52
1	A	291	TYR	C-O	-5.13	1.17	1.24
1	A	438	THR	C-O	5.12	1.29	1.24
1	A	606	GLN	CA-C	5.12	1.59	1.53
1	A	648	GLY	C-O	5.11	1.30	1.23
1	A	659	SER	N-CA	5.11	1.52	1.46
1	A	703	LYS	CA-C	5.08	1.59	1.53
1	A	411	ALA	CA-CB	-5.05	1.48	1.53
1	A	765	ILE	CA-CB	-5.04	1.48	1.54
1	A	426	ASN	C-O	-5.01	1.18	1.23
1	A	720	ILE	CA-CB	-5.01	1.47	1.54

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	754	ARG	NE-CZ-NH2	-8.95	111.14	119.20
1	A	43	ARG	NE-CZ-NH2	-7.43	112.52	119.20
1	A	180	ILE	CA-C-N	-7.21	113.49	120.34
1	A	180	ILE	C-N-CA	-7.21	113.49	120.34
1	A	382	ILE	CB-CA-C	-6.91	101.90	111.65
1	A	557	GLY	N-CA-C	6.54	123.42	115.31
1	A	732	VAL	N-CA-C	-6.21	99.47	108.17
1	A	105	GLN	N-CA-C	-6.20	104.43	111.07
1	A	43	ARG	CD-NE-CZ	6.07	132.89	124.40
1	A	761	ALA	N-CA-C	-5.98	104.68	111.14
1	A	759	ALA	N-CA-C	5.94	118.24	111.11
1	A	754	ARG	CB-CG-CD	-5.69	98.22	111.30
1	A	250	GLY	N-CA-C	-5.68	107.26	115.27
1	A	165	TYR	CA-C-N	-5.60	113.84	120.00
1	A	165	TYR	C-N-CA	-5.60	113.84	120.00
1	A	36	SER	N-CA-C	-5.59	105.27	111.36
1	A	43	ARG	NE-CZ-NH1	5.53	127.03	121.50
1	A	546	GLN	CB-CA-C	5.40	119.98	111.72
1	A	674	VAL	N-CA-C	-5.34	99.95	107.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	740	THR	OG1-CB-CG2	5.15	119.60	109.30
1	A	82	ILE	CB-CA-C	-5.10	105.26	112.14
1	A	269	THR	CA-C-O	5.07	125.92	120.55
1	A	485	PHE	CB-CA-C	-5.04	102.96	110.88
1	A	137	GLY	N-CA-C	-5.00	108.70	115.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6374	0	6083	33	1
2	A	1	0	0	0	0
3	A	26	0	14	0	0
4	A	1	0	0	0	0
5	A	44	0	66	4	0
6	A	819	0	0	6	1
All	All	7265	0	6163	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:549:ASN:O	1:A:740:THR:HB	1.89	0.72
1:A:654:GLU:OE1	1:A:754:ARG:HD3	1.91	0.70
1:A:43:ARG:HD3	6:A:1504:HOH:O	2.01	0.59
1:A:432:ILE:CD1	1:A:446:THR:HG21	2.31	0.59
1:A:797:GLU:OE1	5:A:837:EDO:O1	2.21	0.59
1:A:654:GLU:OE1	1:A:754:ARG:CD	2.52	0.57
1:A:27:LYS:HE2	6:A:950:HOH:O	2.04	0.57
1:A:620:GLU:HG2	1:A:626:ALA:CB	2.34	0.57
1:A:597:LYS:HE3	6:A:1222:HOH:O	2.06	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:525:GLU:HB3	6:A:1387:HOH:O	2.06	0.54
1:A:702:ASP:OD1	5:A:829:EDO:O1	2.25	0.53
1:A:170:ASN:HD21	1:A:431:ARG:HH11	1.55	0.53
1:A:635:GLU:HG2	1:A:639:GLU:OE2	2.08	0.53
1:A:554:GLN:HE22	1:A:738:GLY:HA3	1.74	0.52
1:A:716:VAL:O	1:A:720:ILE:HG12	2.10	0.52
1:A:627:GLU:OE1	5:A:830:EDO:H11	2.12	0.49
1:A:546:GLN:OE1	1:A:553:HIS:HB3	2.12	0.49
1:A:583:MET:HA	1:A:617:ALA:HB1	1.95	0.48
1:A:756:ALA:HB2	1:A:781:ARG:CZ	2.44	0.48
1:A:300:LYS:HE2	1:A:320:HIS:HA	1.95	0.47
1:A:435:PRO:HA	1:A:476:GLN:O	2.16	0.46
1:A:135:TYR:CG	1:A:136:PRO:HD2	2.51	0.45
1:A:96:GLY:HA3	1:A:153:GLU:O	2.18	0.44
1:A:432:ILE:HD13	1:A:446:THR:HG21	1.99	0.44
1:A:498:TRP:CE2	1:A:537:LEU:HD13	2.53	0.43
1:A:411:ALA:HB3	1:A:412:PRO:HD3	2.02	0.42
1:A:454:TRP:CE2	1:A:456:ASN:HB3	2.54	0.42
1:A:434:GLY:O	1:A:475:GLU:HA	2.20	0.42
5:A:831:EDO:C1	6:A:1505:HOH:O	2.68	0.42
1:A:763:LYS:HB2	1:A:763:LYS:HE2	1.44	0.41
1:A:635:GLU:HA	1:A:635:GLU:OE1	2.21	0.41
1:A:543:VAL:HG23	1:A:554:GLN:HB3	2.02	0.40
1:A:190:LEU:O	1:A:190:LEU:HD13	2.21	0.40
1:A:622:GLU:HG2	6:A:1494:HOH:O	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:MET:CE	6:A:1074:HOH:O[6_565]	2.15	0.05

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	800/845 (95%)	769 (96%)	31 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	676/711 (95%)	668 (99%)	8 (1%)	63	63

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	ARG
1	A	189	PRO
1	A	391	LEU
1	A	552	SER
1	A	561	LEU
1	A	740	THR
1	A	754	ARG
1	A	763	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	105	GLN
1	A	118	ASN
1	A	127	GLN
1	A	170	ASN
1	A	441	ASN
1	A	472	GLN
1	A	554	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	574	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

Mogul failed to run properly - this section is therefore empty.

5.5 Carbohydrates [i](#)

Mogul failed to run properly - this section is therefore empty.

5.6 Ligand geometry [i](#)

Mogul failed to run properly - this section is therefore empty.

5.7 Other polymers [i](#)

Mogul failed to run properly - this section is therefore empty.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	802/845 (94%)	-0.32	4 (0%) 87 89	17, 25, 37, 57	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	546	GLN	2.8
1	A	806	VAL	2.7
1	A	5	VAL	2.4
1	A	640	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	A	836	4/4	0.84	0.18	55,56,57,62	0
5	EDO	A	831	4/4	0.88	0.15	57,60,62,62	0
5	EDO	A	832	4/4	0.90	0.15	52,59,60,60	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	EDO	A	837	4/4	0.90	0.12	37,38,41,50	0
5	EDO	A	835	4/4	0.91	0.15	36,38,41,43	0
5	EDO	A	833	4/4	0.92	0.11	35,40,42,42	0
5	EDO	A	834	4/4	0.93	0.14	33,34,38,39	0
5	EDO	A	830	4/4	0.93	0.12	54,54,56,57	0
5	EDO	A	839	4/4	0.94	0.10	37,41,43,45	0
5	EDO	A	838	4/4	0.95	0.08	21,22,22,28	0
5	EDO	A	829	4/4	0.95	0.11	38,39,42,45	0
4	NA	A	828	1/1	0.98	0.04	57,57,57,57	1
3	HTD	A	827	26/26	0.99	0.04	17,20,27,29	0
2	MG	A	826	1/1	1.00	0.03	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.