



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 02:35 PM UTC

PDB ID : 3BLJ / pdb\_00003blj  
Title : Crystal structure of human poly(ADP-ribose) polymerase 15, catalytic fragment  
Authors : Karlberg, T.; Lehtio, L.; Arrowsmith, C.H.; Berglund, H.; Busam, R.D.; Collins, R.; Dahlgren, L.G.; Edwards, A.M.; Flodin, S.; Flores, A.; Graslund, S.; Hammarstrom, M.; Johansson, I.; Kallas, A.; Kotenyova, T.; Moche, M.; Nilsson, M.E.; Nordlund, P.; Nyman, T.; Persson, C.; Sagemark, J.; Svensson, L.; Thorsell, A.G.; Tresaugues, L.; Van Den Berg, S.; Welin, M.; Weigelt, J.; Structural Genomics Consortium (SGC)  
Deposited on : 2007-12-11  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	: 4-5-2 with Phenix2.0
Mogul	: 2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	: 2.0
EDS	: 3.0
Percentile statistics	: 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	: 9.0.010 (Gargrove)
Density-Fitness	: 1.0.12


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	
1	B	221	

Ideal geometry (proteins) : Engh & Huber (2001)  
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.49

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3461 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly(ADP-ribose) polymerase 15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	192	Total	C	N	O	S	0	0	0
			1562	991	272	292	7			
1	B	198	Total	C	N	O	S	0	0	0
			1596	1012	277	300	7			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	436	MET	-	expression tag	UNP Q460N3
A	437	HIS	-	expression tag	UNP Q460N3
A	438	HIS	-	expression tag	UNP Q460N3
A	439	HIS	-	expression tag	UNP Q460N3
A	440	HIS	-	expression tag	UNP Q460N3
A	441	HIS	-	expression tag	UNP Q460N3
A	442	HIS	-	expression tag	UNP Q460N3
A	443	SER	-	expression tag	UNP Q460N3
A	444	SER	-	expression tag	UNP Q460N3
A	445	GLY	-	expression tag	UNP Q460N3
A	446	VAL	-	expression tag	UNP Q460N3
A	447	ASP	-	expression tag	UNP Q460N3
A	448	LEU	-	expression tag	UNP Q460N3
A	449	GLY	-	expression tag	UNP Q460N3
A	450	THR	-	expression tag	UNP Q460N3
A	451	GLU	-	expression tag	UNP Q460N3
A	452	ASN	-	expression tag	UNP Q460N3
A	453	LEU	-	expression tag	UNP Q460N3
A	454	TYR	-	expression tag	UNP Q460N3
A	455	PHE	-	expression tag	UNP Q460N3
A	456	GLN	-	expression tag	UNP Q460N3
A	457	SER	-	expression tag	UNP Q460N3
A	458	MET	-	expression tag	UNP Q460N3
B	436	MET	-	expression tag	UNP Q460N3
B	437	HIS	-	expression tag	UNP Q460N3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	438	HIS	-	expression tag	UNP Q460N3
B	439	HIS	-	expression tag	UNP Q460N3
B	440	HIS	-	expression tag	UNP Q460N3
B	441	HIS	-	expression tag	UNP Q460N3
B	442	HIS	-	expression tag	UNP Q460N3
B	443	SER	-	expression tag	UNP Q460N3
B	444	SER	-	expression tag	UNP Q460N3
B	445	GLY	-	expression tag	UNP Q460N3
B	446	VAL	-	expression tag	UNP Q460N3
B	447	ASP	-	expression tag	UNP Q460N3
B	448	LEU	-	expression tag	UNP Q460N3
B	449	GLY	-	expression tag	UNP Q460N3
B	450	THR	-	expression tag	UNP Q460N3
B	451	GLU	-	expression tag	UNP Q460N3
B	452	ASN	-	expression tag	UNP Q460N3
B	453	LEU	-	expression tag	UNP Q460N3
B	454	TYR	-	expression tag	UNP Q460N3
B	455	PHE	-	expression tag	UNP Q460N3
B	456	GLN	-	expression tag	UNP Q460N3
B	457	SER	-	expression tag	UNP Q460N3
B	458	MET	-	expression tag	UNP Q460N3

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0
2	B	1	Total Na 1 1	0	0

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	133	Total	O	0	0
			133	133		
5	B	155	Total	O	0	0
			155	155		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly(ADP-ribose) polymerase 15



- Molecule 1: Poly(ADP-ribose) polymerase 15



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.21Å 68.12Å 158.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.78 – 2.20 19.78 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.78-2.20) 99.2 (19.78-2.20)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	0.19	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.15 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.3.0040	Depositor
R, $R_{free}$	0.179 , 0.243 0.179 , 0.241	Depositor DCC
$R_{free}$ test set	1273 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.4	Xtriage
Anisotropy	0.362	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 44.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3461	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.92	1/1605 (0.1%)	0.96	2/2174 (0.1%)
1	B	0.96	1/1640 (0.1%)	0.97	1/2223 (0.0%)
All	All	0.94	2/3245 (0.1%)	0.97	3/4397 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	544	VAL	CA-CB	9.52	1.59	1.54
1	A	544	VAL	CA-CB	8.14	1.58	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	612	THR	CA-C-N	-6.08	114.11	120.38
1	A	612	THR	C-N-CA	-6.08	114.11	120.38
1	B	611	VAL	CB-CA-C	5.08	116.42	110.88

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1562	0	1495	9	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1596	0	1528	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
4	A	6	0	8	1	0
4	B	6	0	8	2	0
5	A	133	0	0	0	0
5	B	155	0	0	1	0
All	All	3461	0	3039	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:488:LYS:HG3	1:A:500:ILE:HD12	1.52	0.88
1:A:537:HIS:HD2	1:A:539:THR:HG22	1.50	0.75
1:A:517:VAL:HG21	1:B:517:VAL:HG21	1.75	0.67
1:B:642:PHE:CE2	4:B:702:GOL:H12	2.36	0.61
1:A:537:HIS:CD2	1:A:539:THR:HG22	2.35	0.59
1:A:461:PRO:HG3	1:A:512:TRP:CZ2	2.43	0.54
1:A:488:LYS:HG3	1:A:500:ILE:CD1	2.33	0.53
1:A:467:MET:HG2	1:A:473:CYS:SG	2.50	0.52
1:B:525:LYS:NZ	1:B:626:ASP:OD1	2.43	0.51
1:B:552:PHE:HB3	1:B:568:THR:HG21	1.95	0.48
1:B:520:ARG:HG3	5:B:879:HOH:O	2.13	0.48
1:B:539:THR:HG21	1:B:594:TYR:HE2	1.78	0.48
1:A:522:MET:HE1	1:A:599:LEU:HB3	1.97	0.47
1:B:479:PRO:HA	1:B:484:TYR:CD2	2.53	0.44
1:B:613:PRO:HG2	4:B:702:GOL:O1	2.18	0.43
4:A:702:GOL:H12	1:B:518:LYS:NZ	2.35	0.41
1:B:595:VAL:HB	1:B:650:TYR:HB2	2.03	0.40
1:B:599:LEU:HB2	1:B:645:GLN:HG2	2.04	0.40
1:A:616:LYS:HA	1:A:625:PHE:CE2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/221 (85%)	182 (97%)	6 (3%)	0	100	100
1	B	196/221 (89%)	191 (97%)	5 (3%)	0	100	100
All	All	384/442 (87%)	373 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/198 (87%)	168 (98%)	4 (2%)	44	59
1	B	175/198 (88%)	171 (98%)	4 (2%)	44	59
All	All	347/396 (88%)	339 (98%)	8 (2%)	44	59

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	493	ARG
1	A	534	LEU
1	A	539	THR
1	A	587	SER
1	B	465	THR
1	B	466	ASP
1	B	497	SER
1	B	530	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	481	GLN
1	A	485	ASN
1	A	516	GLN
1	A	588	ASN
1	B	470	GLN
1	B	485	ASN
1	B	528	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	702	-	5,5,5	0.50	0	5,5,5	0.65	0
4	GOL	B	702	-	5,5,5	0.41	0	5,5,5	1.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	702	-	-	1/4/4/4	-
4	GOL	B	702	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	702	GOL	C1-C2-C3-O3
4	B	702	GOL	O2-C2-C3-O3
4	B	702	GOL	O1-C1-C2-O2
4	A	702	GOL	C1-C2-C3-O3
4	B	702	GOL	O1-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	702	GOL	1	0
4	B	702	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	192/221 (86%)	-0.42	2 (1%) 79 77	12, 22, 46, 59	0
1	B	198/221 (89%)	-0.49	2 (1%) 79 77	12, 20, 37, 58	0
All	All	390/442 (88%)	-0.46	4 (1%) 79 77	12, 21, 42, 59	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	557	ALA	3.4
1	B	656	ALA	2.3
1	B	459	ASN	2.2
1	A	619	HIS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NA	A	701	1/1	0.66	0.12	55,55,55,55	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	B	702	6/6	0.83	0.15	40,43,43,44	0
4	GOL	A	702	6/6	0.84	0.12	35,36,38,41	0
2	NA	B	701	1/1	0.87	0.07	40,40,40,40	0
3	CL	A	703	1/1	0.98	0.03	28,28,28,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.