



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:48 AM UTC

PDB ID : 3BOF / pdb_00003bof
Title : Cobalamin-dependent methionine synthase (1-566) from *Thermotoga maritima* complexed with Zn²⁺ and Homocysteine
Authors : Koutmos, M.; Smith, J.L.; Ludwig, M.L.
Deposited on : 2007-12-17
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	566	<div> <div>5%</div> <div>88%</div> <div>10% ..</div> </div>
1	B	566	<div> <div>15%</div> <div>86%</div> <div>11% ..</div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9195 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5-methyltetrahydrofolate S-homocysteine methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	560	Total	C	N	O	S	0	0	0
			4402	2826	734	829	13			
1	B	556	Total	C	N	O	S	0	0	0
			4361	2800	727	821	13			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		


- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	K	0	0
			1	1		
3	B	1	Total	K	0	0
			1	1		

- Molecule 4 is YTTRIUM (III) ION (CCD ID: YT3) (formula: Y).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Y	0	0
			1	1		

- Molecule 5 is 2-AMINO-4-MERCAPTO-BUTYRIC ACID (CCD ID: HCS) (formula: C₄H₉NO₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			8	4	1	2	1		
5	B	1	Total	C	N	O	S	0	0
			8	4	1	2	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	261	Total	O	0	0
			261	261		
6	B	150	Total	O	0	0
			150	150		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5-methyltetrahydrofolate S-homocysteine methyltransferase



- Molecule 1: 5-methyltetrahydrofolate S-homocysteine methyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.07Å 86.31Å 125.88Å 90.00° 100.03° 90.00°	Depositor
Resolution (Å)	46.68 – 1.70 46.68 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.68-1.70) 99.9 (46.68-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.53 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0019, CNS 1.2 / REFMAC 5.2.0019	Depositor
R, R_{free}	0.195 , 0.222 (Not available) , 0.231	Depositor DCC
R_{free} test set	6933 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.098	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9195	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, HCS, YT3, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/4484	0.94	2/6060 (0.0%)
1	B	0.66	0/4442	0.90	4/6004 (0.1%)
All	All	0.73	0/8926	0.92	6/12064 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
All	All	0	4

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	210	GLY	CA-C-N	7.85	127.13	118.97
1	B	210	GLY	C-N-CA	7.85	127.13	118.97
1	B	378	THR	N-CA-C	5.77	117.37	111.14
1	A	558	LYS	N-CA-C	5.47	116.46	108.14
1	B	304	SER	N-CA-C	5.36	117.33	109.24
1	A	559	GLU	N-CA-C	5.07	121.60	110.80

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	411	ASN	Mainchain,Peptide
1	A	558	LYS	Peptide
1	B	411	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4402	0	4463	49	0
1	B	4361	0	4402	39	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
5	A	8	0	7	0	0
5	B	8	0	7	0	0
6	A	261	0	0	4	0
6	B	150	0	0	1	0
All	All	9195	0	8879	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:372:VAL:HG11	1:A:396:LEU:HD21	1.51	0.90
1:A:27:MET:HE3	1:A:32:ASP:HB3	1.52	0.90
1:A:200:ILE:CD1	1:A:203:LEU:HD21	2.02	0.90
1:A:14:ARG:NH1	1:A:289:ASN:OD1	2.06	0.89
1:B:200:ILE:HD11	1:B:203:LEU:HD21	1.53	0.88
1:B:394:VAL:HB	1:B:422:MET:HE1	1.57	0.85
1:B:200:ILE:CD1	1:B:203:LEU:HD21	2.09	0.82
1:A:200:ILE:HD11	1:A:203:LEU:HD21	1.62	0.81
1:A:296:LYS:O	1:A:297:LYS:HB3	1.83	0.77
1:A:402:ARG:HG2	1:A:430:TYR:CE1	2.23	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:ASN:C	1:B:3:ASN:HD22	1.96	0.73
1:A:3:ASN:C	1:A:3:ASN:HD22	1.97	0.72
1:A:200:ILE:CD1	1:A:203:LEU:CD2	2.69	0.69
1:A:237:LYS:HG2	1:A:238:PRO:HD2	1.75	0.67
1:A:335:GLN:NE2	1:A:366:GLN:HG2	2.10	0.67
1:A:301:PHE:CZ	1:A:471:ILE:HG13	2.30	0.66
1:A:394:VAL:HG21	1:A:422:MET:HE3	1.78	0.66
1:A:372:VAL:HG11	1:A:396:LEU:CD2	2.25	0.65
1:A:27:MET:HB2	6:A:1002:HOH:O	1.95	0.65
1:A:508:ASN:HA	1:A:511:PHE:CD2	2.32	0.65
1:B:291:LYS:HG2	1:B:292:PRO:HD2	1.80	0.63
1:A:252:HIS:O	1:A:256:VAL:HG13	1.98	0.63
1:B:369:VAL:HG11	1:B:399:ARG:HD2	1.80	0.63
1:A:200:ILE:HD12	1:A:203:LEU:CD2	2.29	0.63
1:B:390:ASP:HA	1:B:411:ASN:HB3	1.81	0.62
1:A:70:ARG:HD2	1:A:79:GLU:OE2	2.01	0.61
1:A:397:THR:HG23	1:A:410:PHE:CE1	2.36	0.60
1:A:414:LYS:NZ	1:A:439:MET:SD	2.75	0.60
1:B:200:ILE:CD1	1:B:203:LEU:CD2	2.77	0.60
1:B:3:ASN:ND2	1:B:6:GLU:H	2.00	0.58
1:A:213:GLU:HG2	1:A:237:LYS:NZ	2.20	0.57
1:A:200:ILE:HD12	1:A:203:LEU:HD23	1.85	0.57
1:B:4:ARG:HD3	1:B:201:ASP:OD2	2.05	0.57
1:A:188:PRO:HG3	1:A:217:ILE:HG23	1.86	0.57
1:A:122:GLU:HG2	6:A:824:HOH:O	2.05	0.56
1:A:296:LYS:HG3	1:A:297:LYS:H	1.71	0.55
1:A:192:ALA:HB2	1:A:221:LEU:HD12	1.88	0.54
1:B:2:ARG:HD3	1:B:101:LEU:CD2	2.37	0.54
1:B:200:ILE:HD12	1:B:203:LEU:CD2	2.38	0.54
1:B:192:ALA:HB2	1:B:221:LEU:HD12	1.90	0.54
1:B:369:VAL:HG11	1:B:399:ARG:CD	2.37	0.54
1:B:333:GLU:HB3	1:B:338:ASN:HB3	1.89	0.53
1:B:171:ILE:HG12	1:B:202:ALA:HB3	1.90	0.53
1:B:477:LEU:HB2	1:B:483:GLY:HA2	1.89	0.53
1:A:213:GLU:HG2	1:A:237:LYS:HZ1	1.72	0.53
1:B:3:ASN:C	1:B:3:ASN:ND2	2.68	0.52
1:B:394:VAL:CB	1:B:422:MET:HE1	2.37	0.51
1:A:122:GLU:HG2	6:A:742:HOH:O	2.11	0.51
1:B:458:LEU:HD21	1:B:500:PHE:CZ	2.47	0.50
1:A:335:GLN:HE21	1:A:366:GLN:HG2	1.75	0.50
1:A:335:GLN:HE21	1:A:366:GLN:CG	2.24	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:ILE:HG12	1:A:202:ALA:HB3	1.94	0.49
1:B:188:PRO:HG3	1:B:217:ILE:HG23	1.95	0.49
1:A:188:PRO:CG	1:A:217:ILE:HG23	2.43	0.48
1:B:200:ILE:HD12	1:B:203:LEU:HD23	1.94	0.48
1:B:321:ARG:O	1:B:327:ARG:NH1	2.47	0.48
1:A:333:GLU:HA	1:A:336:LYS:HG2	1.96	0.47
1:B:419:GLU:O	1:B:423:LYS:HG2	2.15	0.47
1:A:559:GLU:HB3	1:A:560:LEU:H	1.48	0.47
1:A:2:ARG:HG3	1:A:6:GLU:OE2	2.15	0.47
1:A:533:SER:HB2	6:A:919:HOH:O	2.15	0.47
1:A:335:GLN:NE2	1:A:366:GLN:CG	2.78	0.46
1:B:115:PRO:HG3	1:B:378:THR:HG23	1.96	0.46
1:A:335:GLN:NE2	1:A:366:GLN:HE21	2.14	0.46
1:A:399:ARG:HA	1:A:399:ARG:HD2	1.77	0.46
1:B:414:LYS:HD3	1:B:439:MET:HG3	1.98	0.45
1:B:188:PRO:CG	1:B:217:ILE:HG23	2.46	0.45
1:B:162:ARG:HD2	1:B:166:ARG:HA	1.98	0.45
1:A:417:GLU:OE2	1:A:463:ARG:NH1	2.51	0.44
1:B:17:LEU:HD11	1:B:283:PHE:HB3	2.00	0.43
1:A:162:ARG:HD3	1:A:162:ARG:HA	1.72	0.43
1:B:209:LEU:HB2	1:B:214:ILE:HG13	2.01	0.42
1:A:558:LYS:HD3	1:A:559:GLU:HG3	2.01	0.42
1:B:451:LYS:HG3	1:B:494:PHE:CE2	2.54	0.42
1:B:4:ARG:CD	1:B:201:ASP:OD2	2.68	0.42
1:B:31:TYR:CZ	1:B:45:VAL:HG21	2.54	0.42
1:A:396:LEU:O	1:A:396:LEU:HD23	2.20	0.42
1:A:209:LEU:HB3	1:A:213:GLU:HB2	2.02	0.42
1:B:296:LYS:HG3	1:B:296:LYS:O	2.19	0.42
1:A:394:VAL:HG11	1:A:422:MET:HG2	2.01	0.41
1:B:253:ASP:O	1:B:256:VAL:HG22	2.20	0.41
1:B:324:PRO:HB2	1:B:331:TRP:HB2	2.03	0.41
1:B:392:GLN:NE2	6:B:1101:HOH:O	2.52	0.41
1:A:272:CYS:O	1:A:273:CYS:C	2.63	0.41
1:A:301:PHE:CZ	1:A:471:ILE:CG1	3.03	0.41
1:A:399:ARG:HD2	1:A:399:ARG:N	2.34	0.41
1:B:14:ARG:NH2	1:B:289:ASN:OD1	2.54	0.41
1:B:145:PHE:CE2	1:B:157:ALA:HB1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	558/566 (99%)	537 (96%)	19 (3%)	2 (0%)	30	16
1	B	552/566 (98%)	537 (97%)	12 (2%)	3 (0%)	24	12
All	All	1110/1132 (98%)	1074 (97%)	31 (3%)	5 (0%)	24	12

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	297	LYS
1	A	412	SER
1	B	246	VAL
1	B	245	THR
1	B	412	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	482/494 (98%)	471 (98%)	11 (2%)	44	27
1	B	475/494 (96%)	458 (96%)	17 (4%)	31	14
All	All	957/988 (97%)	929 (97%)	28 (3%)	37	20

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	3	ASN
1	A	18	LEU
1	A	27	MET
1	A	147	THR
1	A	163	GLU
1	A	293	LEU
1	A	295	ARG
1	A	296	LYS
1	A	399	ARG
1	A	509	LEU
1	B	3	ASN
1	B	5	ARG
1	B	17	LEU
1	B	18	LEU
1	B	126	GLU
1	B	143	ILE
1	B	162	ARG
1	B	166	ARG
1	B	293	LEU
1	B	296	LYS
1	B	298	LYS
1	B	396	LEU
1	B	422	MET
1	B	455	GLU
1	B	459	LYS
1	B	463	ARG
1	B	509	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	76	HIS
1	A	252	HIS
1	A	335	GLN
1	A	366	GLN
1	B	3	ASN
1	B	76	HIS
1	B	335	GLN
1	B	366	GLN
1	B	549	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	HCS	A	711	2	6,7,7	1.51	1 (16%)	5,8,8	1.94	1 (20%)
5	HCS	B	712	2	6,7,7	0.90	0	5,8,8	1.85	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HCS	A	711	2	-	0/7/7/7	-
5	HCS	B	712	2	-	1/7/7/7	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	711	HCS	CB-CG	3.01	1.56	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	711	HCS	CB-CG-SD	-3.91	109.67	113.74
5	B	712	HCS	CB-CG-SD	-3.65	109.94	113.74

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	712	HCS	OXT-C-CA-N

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	560/566 (98%)	0.17	29 (5%)	33 36	11, 20, 38, 56	0
1	B	556/566 (98%)	0.79	85 (15%)	5 4	14, 28, 53, 60	0
All	All	1116/1132 (98%)	0.48	114 (10%)	12 12	11, 24, 48, 60	0

All (114) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	560	LEU	8.6
1	A	511	PHE	6.1
1	B	331	TRP	5.6
1	B	29	TYR	5.5
1	A	240	VAL	5.3
1	B	363	ILE	5.2
1	B	245	THR	5.2
1	B	330	LEU	4.7
1	A	243	GLY	4.3
1	B	239	ILE	4.3
1	B	244	LYS	4.0
1	B	483	GLY	4.0
1	B	460	ILE	3.9
1	B	332	ALA	3.9
1	B	246	VAL	3.8
1	B	511	PHE	3.8
1	B	369	VAL	3.8
1	B	443	VAL	3.7
1	A	512	GLY	3.6
1	A	1	MET	3.6
1	B	31	TYR	3.6
1	B	337	GLY	3.5
1	B	166	ARG	3.5
1	B	445	LYS	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	365	SER	3.4
1	B	439	MET	3.4
1	B	447	PHE	3.3
1	B	27	MET	3.2
1	B	324	PRO	3.2
1	A	242	ASN	3.1
1	A	513	LEU	3.0
1	B	367	ILE	3.0
1	B	412	SER	3.0
1	A	289	ASN	3.0
1	A	443	VAL	2.9
1	B	260	SER	2.9
1	A	296	LYS	2.9
1	B	325	ALA	2.9
1	B	560	LEU	2.9
1	B	444	PRO	2.9
1	A	442	ASP	2.9
1	B	422	MET	2.9
1	B	296	LYS	2.9
1	B	464	HIS	2.9
1	B	508	ASN	2.8
1	B	448	GLU	2.8
1	B	486	VAL	2.8
1	B	418	GLU	2.8
1	A	395	ASP	2.8
1	B	170	LEU	2.7
1	B	477	LEU	2.7
1	B	362	GLY	2.7
1	B	481	ALA	2.7
1	B	415	VAL	2.7
1	B	454	PHE	2.6
1	B	516	ARG	2.6
1	B	494	PHE	2.6
1	B	372	VAL	2.6
1	A	239	ILE	2.6
1	B	341	ILE	2.6
1	A	336	LYS	2.6
1	B	514	PRO	2.6
1	B	479	LEU	2.6
1	B	371	TYR	2.6
1	B	1	MET	2.6
1	B	289	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	33	ASP	2.5
1	B	426	LEU	2.5
1	B	461	LEU	2.5
1	A	299	ARG	2.5
1	A	483	GLY	2.5
1	B	476	VAL	2.5
1	B	399	ARG	2.5
1	B	334	MET	2.4
1	B	446	SER	2.4
1	B	336	LYS	2.4
1	B	392	GLN	2.4
1	A	300	ILE	2.4
1	B	343	ILE	2.4
1	B	478	PRO	2.3
1	A	481	ALA	2.3
1	B	442	ASP	2.3
1	B	480	GLY	2.3
1	B	559	GLU	2.3
1	B	449	GLU	2.3
1	B	396	LEU	2.3
1	A	241	GLU	2.3
1	B	370	ARG	2.2
1	B	513	LEU	2.2
1	B	329	LYS	2.2
1	A	514	PRO	2.2
1	B	263	GLU	2.2
1	B	453	TYR	2.2
1	B	484	LYS	2.2
1	B	400	ALA	2.1
1	A	293	LEU	2.1
1	A	325	ALA	2.1
1	B	342	VAL	2.1
1	B	252	HIS	2.1
1	B	326	GLY	2.1
1	A	441	LYS	2.1
1	B	338	ASN	2.1
1	B	438	LEU	2.1
1	B	247	TYR	2.1
1	A	480	GLY	2.1
1	A	27	MET	2.1
1	A	166	ARG	2.1
1	B	491	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	237	LYS	2.1
1	B	335	GLN	2.0
1	B	2	ARG	2.0
1	A	516	ARG	2.0
1	B	300	ILE	2.0
1	A	558	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	YT3	A	705	1/1	0.91	0.14	41,41,41,41	0
5	HCS	A	711	8/8	0.98	0.06	11,13,16,16	0
5	HCS	B	712	8/8	0.98	0.05	17,18,19,20	0
3	K	B	704	1/1	0.99	0.03	17,17,17,17	0
2	ZN	A	701	1/1	0.99	0.02	14,14,14,14	0
2	ZN	B	702	1/1	0.99	0.02	20,20,20,20	0
3	K	A	703	1/1	0.99	0.03	12,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.