



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:50 PM UTC

PDB ID : 3GZ9 / pdb_00003gz9
Title : Crystal Structure of Peroxisome Proliferator-Activated Receptor Delta (PPARd) in Complex with a Full Agonist
Authors : Wang, Z.; Sudom, A.; Walker, N.P.
Deposited on : 2009-04-06
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peroxisome proliferator-activated receptor delta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	262	2115	1371	358	376	10	0	3	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	307	GLY	SER	engineered mutation	UNP Q03181
A	333	LEU	ILE	engineered mutation	UNP Q03181

- Molecule 2 is heptyl beta-D-glucopyranoside (CCD ID: B7G) (formula: C₁₃H₂₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	19	13	6	0	0

- Molecule 3 is (2,3-dimethyl-4-{[2-(prop-2-yn-1-yloxy)-4-{[4-(trifluoromethyl)phenoxy]methyl}phenyl]sulfanyl}phenoxy)acetic acid (CCD ID: D32) (formula: C₂₇H₂₃F₃O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	O	S	0	0
			36	27	3	5	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	172	Total O 172 172	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor delta



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	94.58Å 92.13Å 39.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	94.49 – 2.00 65.99 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.7 (94.49-2.00) 99.8 (65.99-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 1.99Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.189 , 0.245 0.188 , 0.245	Depositor DCC
R_{free} test set	1230 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtriage
Anisotropy	0.591	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.036 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2342	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: D32, B7G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.51	20/2174 (0.9%)	1.40	19/2935 (0.6%)

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	390	ALA	CA-CB	9.13	1.67	1.53
1	A	331	ALA	C-O	7.67	1.33	1.24
1	A	319	LYS	C-O	7.32	1.32	1.24
1	A	392	ILE	N-CA	6.88	1.54	1.46
1	A	323	HIS	N-CA	6.73	1.54	1.46
1	A	370	PHE	C-O	6.64	1.31	1.24
1	A	325	ALA	CA-CB	6.26	1.63	1.53
1	A	320	TYR	C-O	6.07	1.32	1.24
1	A	329	MET	C-O	-6.05	1.16	1.24
1	A	432	PHE	C-O	-5.90	1.19	1.24
1	A	236	ILE	CA-CB	-5.80	1.48	1.54
1	A	252	ILE	N-CA	5.76	1.53	1.46
1	A	214	PHE	C-O	-5.53	1.17	1.24
1	A	372[A]	VAL	CA-CB	5.48	1.61	1.54
1	A	372[B]	VAL	CA-CB	5.48	1.61	1.54
1	A	438	LYS	C-O	-5.41	1.17	1.24
1	A	320	TYR	N-CA	5.23	1.53	1.46
1	A	277	ILE	CA-CB	5.17	1.62	1.54
1	A	385	ALA	C-O	-5.16	1.18	1.24
1	A	321	GLY	N-CA	5.07	1.52	1.45

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	391	ILE	N-CA-C	-7.65	102.99	110.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	289	THR	N-CA-C	-7.34	103.36	111.36
1	A	214	PHE	N-CA-C	-7.05	103.01	111.69
1	A	396	ASP	N-CA-C	6.53	120.16	112.72
1	A	271	LEU	CA-C-N	-6.31	113.88	120.38
1	A	271	LEU	C-N-CA	-6.31	113.88	120.38
1	A	395	GLY	CA-C-N	-6.09	112.93	122.73
1	A	395	GLY	C-N-CA	-6.09	112.93	122.73
1	A	342	ALA	CA-C-N	-5.95	110.18	121.54
1	A	342	ALA	C-N-CA	-5.95	110.18	121.54
1	A	318	LEU	N-CA-C	5.93	117.42	111.07
1	A	387	PHE	N-CA-C	-5.90	104.92	111.36
1	A	344	GLY	N-CA-C	-5.82	107.06	115.27
1	A	231	LYS	N-CA-C	-5.77	104.91	111.14
1	A	472	ILE	N-CA-C	-5.50	105.01	110.62
1	A	425	HIS	CA-C-N	-5.45	114.00	119.56
1	A	425	HIS	C-N-CA	-5.45	114.00	119.56
1	A	361	SER	N-CA-C	-5.18	106.12	112.90
1	A	309	LEU	N-CA-C	-5.01	104.25	110.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2115	0	2167	52	0
2	A	19	0	26	0	0
3	A	36	0	22	2	0
4	A	172	0	0	10	0
All	All	2342	0	2215	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:425:HIS:HD2	4:A:47:HOH:O	1.62	0.80
1:A:257:GLN:CD	4:A:114:HOH:O	2.30	0.74
1:A:207:GLN:O	1:A:210:ASP:HB2	1.94	0.68
1:A:207:GLN:HA	1:A:207:GLN:OE1	1.93	0.67
1:A:237:LEU:HG	1:A:340:LEU:HD11	1.77	0.66
1:A:257:GLN:CG	4:A:114:HOH:O	2.43	0.66
1:A:234:ARG:NH1	1:A:332:SER:O	2.32	0.62
1:A:257:GLN:HG3	4:A:114:HOH:O	2.00	0.61
1:A:260:LYS:O	1:A:260:LYS:HG3	2.01	0.60
1:A:227:ASN:HB3	1:A:343:ASN:OD1	2.03	0.59
1:A:474:LYS:O	1:A:475:ASP:HB2	2.03	0.59
1:A:383:ASP:OD2	1:A:425:HIS:HE1	1.85	0.58
1:A:369:GLU:CD	4:A:96:HOH:O	2.46	0.57
1:A:228:MET:HG2	1:A:333:LEU:HD21	1.86	0.57
1:A:250:HIS:HD2	4:A:54:HOH:O	1.88	0.56
1:A:396:ASP:C	1:A:396:ASP:OD2	2.46	0.56
1:A:421:LEU:HD22	1:A:431:LEU:HD23	1.89	0.55
1:A:312:ASN:ND2	1:A:312:ASN:H	2.05	0.54
1:A:384:LEU:O	1:A:388:ILE:HG12	2.08	0.54
1:A:467:PRO:O	1:A:470:GLN:HB2	2.09	0.53
1:A:322:VAL:O	1:A:326:ILE:HG13	2.10	0.52
1:A:261:GLY:HA2	1:A:264:TRP:O	2.10	0.52
1:A:358:LYS:HE3	1:A:362:ASP:OD2	2.11	0.50
1:A:469:LEU:HD13	3:A:1:D32:H14	1.94	0.49
1:A:250:HIS:HE1	4:A:122:HOH:O	1.96	0.49
1:A:318:LEU:O	1:A:319:LYS:C	2.56	0.47
1:A:421:LEU:HD11	1:A:435:LEU:HD12	1.97	0.47
1:A:289:THR:O	1:A:293:VAL:HG23	2.16	0.46
1:A:327:PHE:CE2	1:A:446:VAL:HG22	2.51	0.46
1:A:403:VAL:HB	1:A:404:PRO:HD3	1.98	0.46
1:A:474:LYS:O	1:A:475:ASP:CB	2.64	0.46
1:A:253:GLU:OE2	1:A:257:GLN:NE2	2.49	0.45
1:A:304:PRO:O	1:A:305:SER:OG	2.26	0.45
1:A:367:LYS:N	1:A:367:LYS:HD2	2.31	0.45
3:A:1:D32:H18	3:A:1:D32:H7A	1.69	0.45
1:A:300:ALA:C	1:A:302:SER:N	2.73	0.45
1:A:373:LYS:HE2	4:A:90:HOH:O	2.17	0.45
1:A:367:LYS:N	1:A:367:LYS:CD	2.80	0.44
1:A:300:ALA:O	1:A:302:SER:N	2.51	0.44
1:A:455:ARG:HA	1:A:455:ARG:HD2	1.58	0.44
1:A:336:LYS:HG2	4:A:157:HOH:O	2.17	0.44
1:A:435:LEU:HD23	1:A:435:LEU:HA	1.73	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:471:GLU:O	1:A:472:ILE:C	2.63	0.41
1:A:414:LEU:HA	1:A:414:LEU:HD23	1.87	0.41
1:A:432:PHE:HB3	1:A:433:PRO:HD3	2.02	0.41
1:A:253:GLU:O	1:A:257:GLN:HG2	2.20	0.41
1:A:265:LYS:HG2	4:A:123:HOH:O	2.21	0.41
1:A:279:VAL:O	1:A:282:PHE:HB3	2.21	0.41
1:A:308:SER:OG	1:A:405[B]:ARG:NH1	2.54	0.41
1:A:319:LYS:HE2	1:A:320:TYR:OH	2.20	0.41
1:A:300:ALA:O	1:A:301:LYS:C	2.64	0.40
1:A:383:ASP:OD2	1:A:425:HIS:CE1	2.71	0.40
1:A:275:LYS:HE2	1:A:462:GLU:CD	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/269 (96%)	247 (95%)	12 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/236 (98%)	219 (94%)	13 (6%)	19	16

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	216	LYS
1	A	234	ARG
1	A	240	LYS
1	A	277	ILE
1	A	294	ARG
1	A	304	PRO
1	A	310	PHE
1	A	312	ASN
1	A	339	LEU
1	A	345	SER
1	A	418	GLU
1	A	461	THR
1	A	462	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	250	HIS
1	A	312	ASN
1	A	425	HIS
1	A	454	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	D32	A	1	-	38,38,38	1.71	7 (18%)	51,53,53	1.75	10 (19%)
2	B7G	A	570	-	19,19,19	0.66	0	24,24,24	1.17	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	D32	A	1	-	-	5/24/24/24	0/3/3/3
2	B7G	A	570	-	-	7/10/30/30	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1	D32	C8-C10	4.85	1.46	1.38
3	A	1	D32	C13-S7	-4.12	1.72	1.78
3	A	1	D32	C11-C19	3.91	1.29	1.18
3	A	1	D32	C3-S7	-3.28	1.73	1.78
3	A	1	D32	C15-C13	-2.83	1.36	1.39
3	A	1	D32	C20-C16	2.73	1.43	1.38
3	A	1	D32	C18-C15	2.42	1.43	1.38

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1	D32	C3-C1-C2	6.39	122.84	118.87
3	A	1	D32	C10-C5-C2	-3.28	116.89	121.65
3	A	1	D32	C24-O25-C26	3.23	125.32	117.62
3	A	1	D32	C5-C2-C1	3.04	121.52	118.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1	D32	C7-O1-C15	-3.01	110.72	117.45
2	A	570	B7G	O5-C5-C4	2.97	115.05	109.70
3	A	1	D32	O22-C17-O21	-2.81	116.11	123.33
3	A	1	D32	C4-C1-C2	-2.46	115.25	119.88
3	A	1	D32	C15-C18-C23	2.36	123.53	120.04
2	A	570	B7G	O6-C6-C5	-2.32	103.44	111.33
3	A	1	D32	C18-C15-C13	-2.21	118.31	120.78
3	A	1	D32	O22-C17-C14	2.04	123.27	113.66

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	570	B7G	C8-C7-O1-C1
2	A	570	B7G	C4-C5-C6-O6
2	A	570	B7G	O5-C5-C6-O6
2	A	570	B7G	C11-C10-C9-C8
2	A	570	B7G	O1-C7-C8-C9
3	A	1	D32	C28-C26-O25-C24
3	A	1	D32	C15-C13-S7-C3
3	A	1	D32	C27-C26-O25-C24
2	A	570	B7G	C7-C8-C9-C10
2	A	570	B7G	C9-C10-C11-C12
3	A	1	D32	C23-C24-O25-C26
3	A	1	D32	C19-C11-C7-O1


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1	D32	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/269 (97%)	-0.42	3 (1%) 78 77	13, 29, 50, 74	3 (1%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	241	ALA	4.9
1	A	310	PHE	2.3
1	A	208	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	B7G	A	570	19/19	0.84	0.14	48,64,70,71	0
3	D32	A	1	36/36	0.97	0.05	15,24,29,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.