



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:30 AM UTC

PDB ID : 3KDA / pdb\_00003kda  
Title : Crystal structure of the CFTR inhibitory factor Cif with the H269A mutation  
Authors : Bahl, C.D.; Madden, D.R.  
Deposited on : 2009-10-22  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	<div> <div>4%</div> <div> <div></div> <div>91%</div> <div>8%</div> </div> </div>
1	B	301	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>8%</div> </div> </div>
1	C	301	<div> <div>3%</div> <div> <div></div> <div>94%</div> <div>5%</div> </div> </div>
1	D	301	<div> <div>2%</div> <div> <div></div> <div>91%</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CFTR inhibitory factor (Cif).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	298	Total	C	N	O	S	0	1	0
			2383	1531	414	427	11			
1	C	298	Total	C	N	O	S	0	0	0
			2378	1528	413	426	11			
1	A	298	Total	C	N	O	S	0	4	0
			2397	1542	416	428	11			
1	B	298	Total	C	N	O	S	0	4	0
			2403	1544	419	429	11			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	269	ALA	HIS	engineered mutation	UNP Q02P97
D	320	HIS	-	expression tag	UNP Q02P97
D	321	HIS	-	expression tag	UNP Q02P97
D	322	HIS	-	expression tag	UNP Q02P97
D	323	HIS	-	expression tag	UNP Q02P97
D	324	HIS	-	expression tag	UNP Q02P97
D	325	HIS	-	expression tag	UNP Q02P97
C	269	ALA	HIS	engineered mutation	UNP Q02P97
C	320	HIS	-	expression tag	UNP Q02P97
C	321	HIS	-	expression tag	UNP Q02P97
C	322	HIS	-	expression tag	UNP Q02P97
C	323	HIS	-	expression tag	UNP Q02P97
C	324	HIS	-	expression tag	UNP Q02P97
C	325	HIS	-	expression tag	UNP Q02P97
A	269	ALA	HIS	engineered mutation	UNP Q02P97
A	320	HIS	-	expression tag	UNP Q02P97
A	321	HIS	-	expression tag	UNP Q02P97
A	322	HIS	-	expression tag	UNP Q02P97
A	323	HIS	-	expression tag	UNP Q02P97
A	324	HIS	-	expression tag	UNP Q02P97
A	325	HIS	-	expression tag	UNP Q02P97

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	269	ALA	HIS	engineered mutation	UNP Q02P97
B	320	HIS	-	expression tag	UNP Q02P97
B	321	HIS	-	expression tag	UNP Q02P97
B	322	HIS	-	expression tag	UNP Q02P97
B	323	HIS	-	expression tag	UNP Q02P97
B	324	HIS	-	expression tag	UNP Q02P97
B	325	HIS	-	expression tag	UNP Q02P97


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	314	Total 314	O 314	0	0
2	C	269	Total 269	O 269	0	0
2	A	288	Total 288	O 288	0	0
2	B	240	Total 240	O 240	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CFTR inhibitory factor (Cif)




- Molecule 1: CFTR inhibitory factor (Cif)



- Molecule 1: CFTR inhibitory factor (Cif)



- Molecule 1: CFTR inhibitory factor (Cif)



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	167.96Å 83.64Å 88.93Å 90.00° 100.49° 90.00°	Depositor
Resolution (Å)	30.72 – 1.50 30.72 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.72-1.50) 99.8 (30.72-1.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.00 (at 1.50Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.169 , 0.183 0.164 , 0.176	Depositor DCC
$R_{free}$ test set	9636 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.4	Xtriage
Anisotropy	0.278	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 45.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10672	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.38	0/2482	0.68	0/3368
1	B	0.38	0/2488	0.68	0/3375
1	C	0.38	0/2451	0.66	0/3326
1	D	0.39	0/2459	0.67	0/3337
All	All	0.38	0/9880	0.67	0/13406

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2397	0	2323	16	0
1	B	2403	0	2326	16	0
1	C	2378	0	2289	9	0
1	D	2383	0	2295	14	0
2	A	288	0	0	3	0
2	B	240	0	0	1	0
2	C	269	0	0	1	0
2	D	314	0	0	2	0
All	All	10672	0	9233	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:287:VAL:HG12	2:A:664:HOH:O	1.58	1.01
1:D:256:ARG:HE	1:D:285:GLU:HG2	1.35	0.91
1:C:129:ASP:HA	1:C:154:ALA:HB2	1.71	0.72
1:B:129:ASP:HA	1:B:154:ALA:HB2	1.71	0.72
1:D:256:ARG:NE	1:D:285:GLU:HG2	2.06	0.70
1:A:129:ASP:HA	1:A:154:ALA:HB2	1.77	0.67
1:A:261[B]:THR:HG21	1:A:280:MET:HE2	1.77	0.65
1:D:129:ASP:HA	1:D:154:ALA:HB2	1.77	0.65
1:D:156:ILE:HD11	1:D:280:MET:HE1	1.79	0.65
1:B:80:ARG:HD3	1:B:320:HIS:CD2	2.33	0.64
1:A:261[B]:THR:HG21	1:A:280:MET:CE	2.30	0.62
1:D:39:ARG:HH21	1:D:116:GLN:HE21	1.48	0.62
1:D:39:ARG:HH21	1:D:116:GLN:NE2	2.01	0.57
1:A:57[B]:VAL:HG22	1:A:124:ASP:HB2	1.87	0.57
1:B:53:GLN:NE2	1:B:321:HIS:CE1	2.73	0.56
1:C:62:GLY:HA3	1:C:129:ASP:HB3	1.89	0.55
1:C:251:GLU:O	1:C:254:LYS:HG2	2.06	0.55
1:B:210:ASN:HB3	2:B:786:HOH:O	2.08	0.53
1:C:98:LYS:HE3	2:C:850:HOH:O	2.09	0.52
1:A:62:GLY:HA3	1:A:129:ASP:HB3	1.91	0.52
1:B:132:ILE:HD12	1:B:154:ALA:HB3	1.92	0.52
1:D:62:GLY:HA3	1:D:129:ASP:HB3	1.93	0.51
1:A:217[B]:ARG:NH1	2:A:426:HOH:O	2.34	0.51
1:A:107:ALA:HB1	1:A:138:MET:HB3	1.95	0.48
1:B:259:MET:O	1:B:261[B]:THR:HG23	2.13	0.48
1:B:62:GLY:HA3	1:B:129:ASP:HB3	1.96	0.47
1:B:251:GLU:OE1	1:B:254:LYS:HE3	2.15	0.47
1:C:288:GLU:OE1	1:C:317:ARG:NH1	2.47	0.46
1:B:57:VAL:HG22	1:B:124:ASP:HB2	1.98	0.46
1:A:184:ASP:O	1:A:185:ASP:HB2	2.16	0.46
1:D:59:LEU:O	1:D:86:PRO:HD2	2.16	0.45
1:A:140:VAL:HG11	1:A:252:LEU:HB3	1.99	0.45
1:B:149:LEU:HB3	1:B:261[B]:THR:HG22	1.99	0.45
1:D:240:ARG:HD3	2:D:785:HOH:O	2.16	0.45
1:A:217[B]:ARG:NH1	2:A:591:HOH:O	2.48	0.45
1:D:184:ASP:HA	2:D:785:HOH:O	2.18	0.44
1:C:172:GLU:HG2	1:A:212:GLU:OE1	2.17	0.44
1:B:251:GLU:CD	1:B:254:LYS:HE3	2.43	0.44
1:B:59:LEU:O	1:B:86:PRO:HD2	2.18	0.44
1:C:292:LEU:HD12	1:C:292:LEU:N	2.33	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:ARG:NH1	1:A:317:ARG:HD2	2.33	0.43
1:A:309:ARG:HD2	1:A:313:ASP:OD2	2.19	0.43
1:D:55:PRO:HG2	1:D:319:ARG:O	2.19	0.42
1:A:179:SER:HB3	1:B:176:TRP:CD1	2.55	0.42
1:A:149:LEU:HD11	1:A:151:TYR:CE1	2.55	0.41
1:D:57:VAL:HG22	1:D:124:ASP:HB2	2.02	0.41
1:D:262:MET:HA	1:D:288:GLU:O	2.21	0.41
1:C:140:VAL:HG11	1:C:252:LEU:HB3	2.03	0.41
1:D:158:ASP:O	1:D:161:ILE:HG12	2.22	0.40
1:C:107:ALA:HB1	1:C:138:MET:HB3	2.03	0.40
1:B:115[A]:ARG:HA	1:B:115[A]:ARG:HD2	1.87	0.40
1:B:262:MET:HA	1:B:288:GLU:O	2.21	0.40
1:B:275:PHE:CD1	1:B:275:PHE:C	2.99	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	300/301 (100%)	292 (97%)	8 (3%)	0	100	100
1	B	300/301 (100%)	292 (97%)	8 (3%)	0	100	100
1	C	296/301 (98%)	288 (97%)	8 (3%)	0	100	100
1	D	297/301 (99%)	290 (98%)	7 (2%)	0	100	100
All	All	1193/1204 (99%)	1162 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/247 (100%)	247 (100%)	1 (0%)	84	71
1	B	248/247 (100%)	247 (100%)	1 (0%)	84	71
1	C	244/247 (99%)	243 (100%)	1 (0%)	84	71
1	D	245/247 (99%)	243 (99%)	2 (1%)	73	54
All	All	985/988 (100%)	980 (100%)	5 (0%)	81	66

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	226	TYR
1	D	303	CYS
1	C	226	TYR
1	A	226	TYR
1	B	226	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	53	GLN
1	D	116	GLN
1	D	243	ASN
1	D	322	HIS
1	C	243	ASN
1	A	170	GLN
1	A	243	ASN
1	B	53	GLN
1	B	243	ASN
1	B	321	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	298/301 (99%)	-0.24	11 (3%) 45 49	5, 11, 25, 47	4 (1%)
1	B	298/301 (99%)	-0.19	8 (2%) 56 60	6, 12, 31, 45	4 (1%)
1	C	298/301 (99%)	-0.17	8 (2%) 56 60	6, 12, 29, 47	0
1	D	298/301 (99%)	-0.37	6 (2%) 65 68	5, 11, 24, 42	1 (0%)
All	All	1192/1204 (99%)	-0.25	33 (2%) 55 59	5, 11, 27, 47	9 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	25	ALA	5.8
1	B	321	HIS	4.3
1	B	25	ALA	3.9
1	C	319	ARG	3.7
1	A	322	HIS	3.6
1	B	322	HIS	3.5
1	B	319	ARG	3.5
1	B	318	GLY	3.4
1	C	322	HIS	3.4
1	D	319	ARG	3.3
1	A	319	ARG	3.1
1	A	321	HIS	3.0
1	A	255	THR	3.0
1	A	318	GLY	3.0
1	D	321	HIS	2.9
1	D	25	ALA	2.9
1	C	321	HIS	2.7
1	A	25	ALA	2.6
1	C	170	GLN	2.6
1	A	121	ARG	2.4
1	C	53	GLN	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	212	GLU	2.3
1	B	210	ASN	2.3
1	C	318	GLY	2.3
1	D	322	HIS	2.2
1	D	318	GLY	2.2
1	B	32	ASN	2.2
1	A	170	GLN	2.2
1	A	256	ARG	2.2
1	C	255	THR	2.1
1	B	213	VAL	2.1
1	A	32	ASN	2.1
1	D	53	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.