



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 05:24 AM UTC

PDB ID : 3UO0 / pdb\_00003uo0  
Title : phosphorylated Bacillus cereus phosphopentomutase soaked with glucose 1,6-bisphosphate  
Authors : Iverson, T.M.; Birmingham, W.R.; Panosian, T.D.; Nannemann, D.P.; Bachmann, B.O.  
Deposited on : 2011-11-16  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	399	<div> <div> <div></div> <div>91%</div> <div>7%</div> <div></div> </div> <div></div> </div>
1	B	399	<div> <div> <div></div> <div>88%</div> <div>9%</div> <div></div> </div> <div></div> </div>
1	C	399	<div> <div> <div>2%</div> <div>90%</div> <div>6%</div> <div></div> </div> <div></div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphopentomutase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	391	Total	C	N	O	P	S	8	0	0
			3068	1939	505	606	1	17			
1	B	390	Total	C	N	O	P	S	13	0	0
			3060	1935	503	604	1	17			
1	C	390	Total	C	N	O	P	S	29	0	0
			3060	1935	503	604	1	17			


There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q818Z9
A	-3	SER	-	expression tag	UNP Q818Z9
A	-2	HIS	-	expression tag	UNP Q818Z9
A	-1	MET	-	expression tag	UNP Q818Z9
A	0	ALA	-	expression tag	UNP Q818Z9
A	1	SER	-	expression tag	UNP Q818Z9
B	-4	GLY	-	expression tag	UNP Q818Z9
B	-3	SER	-	expression tag	UNP Q818Z9
B	-2	HIS	-	expression tag	UNP Q818Z9
B	-1	MET	-	expression tag	UNP Q818Z9
B	0	ALA	-	expression tag	UNP Q818Z9
B	1	SER	-	expression tag	UNP Q818Z9
C	-4	GLY	-	expression tag	UNP Q818Z9
C	-3	SER	-	expression tag	UNP Q818Z9
C	-2	HIS	-	expression tag	UNP Q818Z9
C	-1	MET	-	expression tag	UNP Q818Z9
C	0	ALA	-	expression tag	UNP Q818Z9
C	1	SER	-	expression tag	UNP Q818Z9

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Mn	0	0
			3	3		
2	B	2	Total	Mn	0	0
			2	2		
2	C	3	Total	Mn	0	0
			3	3		

- Molecule 3 is 1,6-di-O-phosphono-alpha-D-glucopyranose (CCD ID: G16) (formula:  $C_6H_{13}O_{12}P_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			20	6	12	2		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula:  $C_4H_{12}NO_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	179	Total	O	0	0
			179	179		
5	B	196	Total	O	0	0
			196	196		
5	C	84	Total	O	0	0
			84	84		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phosphopentomutase



- Molecule 1: Phosphopentomutase



- Molecule 1: Phosphopentomutase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.82Å 77.08Å 108.33Å 90.00° 108.94° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 20.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-2.30) 99.5 (20.00-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.97 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.5.0109, CNS	Depositor
R, $R_{free}$	0.182 , 0.216 0.184 , 0.217	Depositor DCC
$R_{free}$ test set	7460 reflections (11.71%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.2	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 43.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9683	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, MN, G16, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	0/3117	0.76	1/4208 (0.0%)
1	B	0.44	0/3109	0.76	0/4197
1	C	0.83	6/3109 (0.2%)	0.92	9/4197 (0.2%)
All	All	0.60	6/9335 (0.1%)	0.82	10/12602 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	257	LYS	CD-CE	24.96	2.27	1.52
1	C	65	LYS	CD-CE	-17.09	1.01	1.52
1	C	202	GLU	CG-CD	16.47	1.93	1.52
1	C	67	GLU	CB-CG	16.42	2.01	1.52
1	C	208	ARG	CB-CG	-7.56	1.29	1.52
1	C	182	GLU	CG-CD	6.95	1.69	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	65	LYS	CG-CD-CE	21.76	161.36	111.30
1	C	202	GLU	CB-CG-CD	-14.55	87.87	112.60
1	C	257	LYS	CG-CD-CE	-13.75	79.68	111.30
1	C	67	GLU	CA-CB-CG	-10.61	92.87	114.10
1	C	182	GLU	CB-CG-CD	-6.73	101.16	112.60
1	C	167	VAL	N-CA-C	6.28	116.91	110.82
1	C	67	GLU	CB-CG-CD	6.14	123.04	112.60
1	C	166	GLU	CB-CA-C	5.24	119.76	110.85
1	C	208	ARG	CA-CB-CG	5.17	124.44	114.10
1	A	289	PHE	N-CA-C	5.11	119.99	113.55

There are no chirality outliers.



There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3068	0	3007	12	0
1	B	3060	0	3001	20	0
1	C	3060	0	3001	14	0
2	A	3	0	0	0	0
2	B	2	0	0	0	0
2	C	3	0	0	0	0
3	A	20	0	10	0	0
4	A	8	0	12	0	0
5	A	179	0	0	0	0
5	B	196	0	0	0	0
5	C	84	0	0	0	0
All	All	9683	0	9031	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:4:TYR:H	1:B:376:ASN:HD21	1.34	0.73
1:A:164:HIS:HD2	1:A:166:GLU:H	1.44	0.65
1:A:4:TYR:H	1:A:376:ASN:HD21	1.45	0.62
1:C:4:TYR:H	1:C:376:ASN:HD21	1.47	0.61
1:A:56:ASN:HD21	1:A:66:VAL:H	1.51	0.59
1:C:164:HIS:HD2	1:C:166:GLU:H	1.50	0.59
1:B:240:LYS:HE3	1:B:244:ILE:HD11	1.84	0.59
1:B:164:HIS:HD2	1:B:166:GLU:H	1.51	0.57
1:C:331:ASP:HB3	1:C:334:HIS:HB2	1.86	0.57
1:B:56:ASN:HD21	1:B:66:VAL:H	1.51	0.56
1:B:93:MET:O	1:B:224:VAL:HG23	2.05	0.56
1:A:14:SER:HB2	1:A:330:ASN:HB2	1.87	0.55
1:A:45:GLN:H	1:A:303:GLN:HE21	1.54	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:LEU:HD22	1:B:124:ILE:HG21	1.87	0.55
1:B:45:GLN:H	1:B:303:GLN:HE21	1.54	0.55
1:B:227:GLU:HG3	1:B:379:VAL:HB	1.89	0.54
1:C:256:THR:HG21	1:C:262:GLY:HA2	1.92	0.52
1:A:240:LYS:HE3	1:A:244:ILE:HD11	1.92	0.51
1:A:256:THR:HG21	1:A:262:GLY:HA2	1.92	0.51
1:B:331:ASP:HB3	1:B:334:HIS:HB2	1.93	0.51
1:C:45:GLN:H	1:C:303:GLN:HE21	1.59	0.51
1:C:113:LEU:HD22	1:C:124:ILE:HG21	1.92	0.51
1:C:14:SER:HB2	1:C:330:ASN:HB2	1.93	0.50
1:A:46:MET:HE2	1:A:49:MET:HG3	1.95	0.49
1:C:267:VAL:HA	1:C:270:LEU:HD12	1.95	0.49
1:A:267:VAL:HA	1:A:270:LEU:HD12	1.95	0.48
1:C:56:ASN:HD21	1:C:66:VAL:H	1.62	0.48
1:A:113:LEU:HD22	1:A:124:ILE:HG21	1.96	0.47
1:C:184:THR:O	1:C:190:MET:HA	2.14	0.47
1:B:46:MET:HE2	1:B:49:MET:HG3	1.97	0.45
1:B:14:SER:HB2	1:B:330:ASN:HB2	1.98	0.44
1:C:202:GLU:HG2	1:C:205:ASN:HB3	2.00	0.44
1:B:256:THR:HG21	1:B:262:GLY:HA2	2.00	0.43
1:A:4:TYR:N	1:A:376:ASN:HD21	2.15	0.43
1:A:288:LEU:O	1:A:292:ARG:HD2	2.20	0.42
1:B:150:ILE:HB	1:B:162:ALA:HB3	2.01	0.42
1:B:164:HIS:HE1	1:B:201:GLY:O	2.02	0.42
1:B:164:HIS:CD2	1:B:166:GLU:H	2.33	0.41
1:C:14:SER:HB2	1:C:330:ASN:CB	2.50	0.41
1:B:44:LEU:HD23	1:B:63:ILE:HD11	2.01	0.41
1:C:44:LEU:HD23	1:C:63:ILE:HD11	2.02	0.41
1:B:14:SER:HA	1:B:328:HIS:O	2.22	0.40
1:B:170:LEU:HD22	1:B:174:TYR:CE2	2.57	0.40
1:B:213:HIS:HD2	1:C:234:ASP:OD1	2.04	0.40
1:B:267:VAL:HA	1:B:270:LEU:HD12	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/399 (97%)	372 (96%)	16 (4%)	0	100	100
1	B	387/399 (97%)	374 (97%)	13 (3%)	0	100	100
1	C	387/399 (97%)	371 (96%)	16 (4%)	0	100	100
All	All	1162/1197 (97%)	1117 (96%)	45 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	332/338 (98%)	327 (98%)	5 (2%)	57	75
1	B	331/338 (98%)	328 (99%)	3 (1%)	70	84
1	C	331/338 (98%)	325 (98%)	6 (2%)	51	70
All	All	994/1014 (98%)	980 (99%)	14 (1%)	59	76

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	GLU
1	A	189	TYR
1	A	212	ARG
1	A	281	ASN
1	A	360	LEU
1	B	170	LEU
1	B	189	TYR
1	B	360	LEU
1	C	65	LYS
1	C	67	GLU
1	C	124	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	202	GLU
1	C	208	ARG
1	C	257	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (35) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	48	ASN
1	A	56	ASN
1	A	77	GLN
1	A	102	GLN
1	A	164	HIS
1	A	205	ASN
1	A	211	ASN
1	A	259	ASN
1	A	358	GLN
1	A	364	GLN
1	A	376	ASN
1	A	390	ASN
1	B	25	GLN
1	B	48	ASN
1	B	56	ASN
1	B	77	GLN
1	B	89	HIS
1	B	164	HIS
1	B	213	HIS
1	B	259	ASN
1	B	281	ASN
1	B	296	GLN
1	B	303	GLN
1	B	376	ASN
1	C	56	ASN
1	C	77	GLN
1	C	89	HIS
1	C	164	HIS
1	C	211	ASN
1	C	213	HIS
1	C	271	ASN
1	C	281	ASN
1	C	303	GLN
1	C	376	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	TPO	B	85	2,1	8,10,11	0.74	0	10,14,16	1.15	0
1	TPO	A	85	2,1	8,10,11	0.79	0	10,14,16	1.12	0
1	TPO	C	85	2,1	8,10,11	0.79	0	10,14,16	1.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	B	85	2,1	-	3/9/11/13	-
1	TPO	A	85	2,1	-	3/9/11/13	-
1	TPO	C	85	2,1	-	1/9/11/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	85	TPO	O-C-CA-CB
1	B	85	TPO	O-C-CA-CB
1	A	85	TPO	CG2-CB-OG1-P
1	B	85	TPO	CG2-CB-OG1-P

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
1	A	85	TPO	CB-OG1-P-O1P
1	B	85	TPO	CB-OG1-P-O1P
1	C	85	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 8 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	G16	A	398	-	19,20,20	1.31	3 (15%)	30,31,31	0.91	1 (3%)
4	TRS	A	399	-	7,7,7	0.28	0	9,9,9	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	G16	A	398	-	-	8/11/31/31	0/1/1/1
4	TRS	A	399	-	-	0/9/9/9	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	398	G16	P-O1P	3.79	1.62	1.50
3	A	398	G16	P'-O1X	2.03	1.62	1.54
3	A	398	G16	P'-O3X	2.02	1.62	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	398	G16	O2P-P-O6	2.54	113.30	106.67

There are no chirality outliers.


All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	398	G16	C4-C5-C6-O6
3	A	398	G16	O5-C5-C6-O6
3	A	398	G16	C1-O1-P'-O2X
3	A	398	G16	C6-O6-P-O2P
3	A	398	G16	C6-O6-P-O1P
3	A	398	G16	O5-C1-O1-P'
3	A	398	G16	C6-O6-P-O3P
3	A	398	G16	C1-O1-P'-O1X

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	390/399 (97%)	-0.21	3 (0%) 82 83	16, 31, 46, 63	17 (4%)
1	B	389/399 (97%)	-0.25	0 100 100	15, 30, 42, 52	14 (3%)
1	C	389/399 (97%)	0.15	8 (2%) 63 65	16, 42, 68, 85	24 (6%)
All	All	1168/1197 (97%)	-0.10	11 (0%) 81 82	15, 34, 59, 85	55 (4%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	99	THR	3.1
1	C	189	TYR	2.8
1	A	273	ASP	2.7
1	A	171	ASP	2.5
1	C	186	ASP	2.4
1	C	155	ALA	2.2
1	C	110	LYS	2.1
1	C	67	GLU	2.1
1	C	355	GLU	2.0
1	C	102	GLN	2.0
1	A	2	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	C	85	11/12	0.93	0.09	34,35,37,38	0
1	TPO	B	85	11/12	0.98	0.07	26,27,30,31	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	TPO	A	85	11/12	0.98	0.07	28,29,33,33	0

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	G16	A	398	20/20	0.79	0.26	54,55,55,55	20
4	TRS	A	399	8/8	0.81	0.14	56,56,56,56	0
2	MN	B	396	1/1	0.99	0.02	26,26,26,26	0
2	MN	C	395	1/1	0.99	0.02	39,39,39,39	0
2	MN	C	396	1/1	0.99	0.03	33,33,33,33	0
2	MN	C	397	1/1	0.99	0.03	50,50,50,50	0
2	MN	A	396	1/1	0.99	0.03	24,24,24,24	0
2	MN	A	397	1/1	0.99	0.03	30,30,30,30	1
2	MN	A	395	1/1	1.00	0.01	25,25,25,25	0
2	MN	B	395	1/1	1.00	0.02	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around G16 A 398:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.