



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:18 PM UTC

PDB ID : 3WDT / pdb\_00003wdt  
Title : The apo-form structure of PtLic16A from Paecilomyces thermophila  
Authors : Cheng, Y.S.; Huang, C.H.; Chen, C.C.; Huang, T.Y.; Ko, T.P.; Huang, J.W.;  
Wu, T.H.; Liu, J.R.; Guo, R.T.  
Deposited on : 2013-06-25  
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 14%, green 85%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>85%</span> <span>14%</span> <span>.</span> </div> </div>
1	B	298	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 89%, yellow 11%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>89%</span> <span>11%</span> <span></span> </div> </div>
1	C	298	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 12%, green 87%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>87%</span> <span>12%</span> <span>.</span> </div> </div>
1	D	298	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 90%, yellow 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>90%</span> <span>9%</span> <span>.</span> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9772 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-1,3-1,4-glucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	0	0	0
			2276	1434	388	443	11			
1	B	297	Total	C	N	O	S	0	0	0
			2267	1429	387	440	11			
1	C	297	Total	C	N	O	S	0	0	0
			2267	1429	387	440	11			
1	D	297	Total	C	N	O	S	0	0	0
			2267	1429	387	440	11			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLU	-	expression tag	UNP E0XN39
A	0	PHE	-	expression tag	UNP E0XN39
B	-1	GLU	-	expression tag	UNP E0XN39
B	0	PHE	-	expression tag	UNP E0XN39
C	-1	GLU	-	expression tag	UNP E0XN39
C	0	PHE	-	expression tag	UNP E0XN39
D	-1	GLU	-	expression tag	UNP E0XN39
D	0	PHE	-	expression tag	UNP E0XN39

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	186	Total	O	0	0
			186	186		
3	B	187	Total	O	0	0
			187	187		
3	C	151	Total	O	0	0
			151	151		
3	D	166	Total	O	0	0
			166	166		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Beta-1,3-1,4-glucanase




#### • Molecule 1: Beta-1,3-1,4-glucanase



#### • Molecule 1: Beta-1,3-1,4-glucanase



#### • Molecule 1: Beta-1,3-1,4-glucanase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.34Å 172.93Å 167.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.98 25.00 – 1.98	Depositor EDS
% Data completeness (in resolution range)	96.9 (25.00-1.98) 96.8 (25.00-1.98)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.12 (at 1.98Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.171 , 0.208 0.171 , 0.207	Depositor DCC
$R_{free}$ test set	4368 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.0	Xtriage
Anisotropy	0.353	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 37.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9772	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.67	0/2346	1.07	11/3196 (0.3%)
1	B	0.66	0/2337	1.06	12/3184 (0.4%)
1	C	0.64	0/2337	1.04	16/3184 (0.5%)
1	D	0.62	0/2337	1.04	12/3184 (0.4%)
All	All	0.65	0/9357	1.05	51/12748 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	60	PRO	N-CA-C	-8.01	98.65	111.14
1	A	100	PHE	N-CA-C	-7.21	96.66	108.41
1	D	100	PHE	N-CA-C	-7.10	96.84	108.41
1	C	100	PHE	N-CA-C	-7.00	97.00	108.41
1	B	60	PRO	N-CA-C	-6.98	100.26	111.14
1	C	281	PHE	N-CA-C	6.89	121.39	112.92
1	C	276	ASP	N-CA-C	6.76	121.52	113.28
1	C	59	ASN	N-CA-C	6.73	119.14	110.39
1	C	6	ASP	N-CA-C	-6.63	97.71	108.52
1	D	276	ASP	N-CA-C	6.53	121.26	113.16
1	B	276	ASP	N-CA-C	6.49	121.35	113.17
1	C	288	VAL	N-CA-C	6.49	117.92	108.58

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	59	ASN	N-CA-C	6.45	118.78	110.39
1	B	100	PHE	N-CA-C	-6.44	97.92	108.41
1	C	60	PRO	N-CA-C	-6.30	101.20	111.21
1	A	252	ASP	N-CA-C	6.25	119.75	111.75
1	B	59	ASN	N-CA-C	6.24	118.50	110.39
1	A	276	ASP	N-CA-C	6.24	120.89	113.16
1	D	281	PHE	N-CA-C	6.16	119.74	112.72
1	D	60	PRO	N-CA-C	-6.12	101.64	111.38
1	D	252	ASP	N-CA-C	6.05	118.65	111.33
1	D	288	VAL	N-CA-C	5.97	116.71	107.99
1	A	110	TYR	N-CA-C	5.88	119.55	112.38
1	B	281	PHE	N-CA-C	5.75	120.00	112.92
1	B	252	ASP	N-CA-C	5.74	119.10	111.75
1	A	87	ALA	N-CA-C	-5.73	106.12	113.23
1	D	59	ASN	N-CA-C	5.68	117.77	110.39
1	C	252	ASP	N-CA-C	5.65	118.98	111.75
1	C	228	ALA	N-CA-C	5.53	115.85	108.38
1	B	186	ILE	N-CA-C	5.50	115.80	108.11
1	A	281	PHE	N-CA-C	5.38	118.85	112.72
1	C	110	TYR	N-CA-C	5.37	117.89	111.71
1	C	18	ASN	N-CA-C	-5.34	101.57	110.17
1	A	259	GLY	N-CA-C	5.33	120.62	114.16
1	B	234	TRP	N-CA-C	5.25	117.68	111.33
1	B	288	VAL	N-CA-C	5.24	116.24	108.23
1	C	156	ALA	CA-C-N	5.22	126.37	119.84
1	C	156	ALA	C-N-CA	5.22	126.37	119.84
1	C	289	ASN	N-CA-C	-5.22	105.27	111.69
1	B	206	GLU	N-CA-C	5.19	116.42	109.83
1	A	289	ASN	N-CA-C	-5.18	105.08	111.40
1	C	138	ILE	N-CA-C	5.18	116.82	108.85
1	D	89	MET	CA-C-N	-5.13	114.44	119.93
1	D	89	MET	C-N-CA	-5.13	114.44	119.93
1	B	18	ASN	N-CA-C	-5.11	101.36	109.59
1	C	108	TRP	N-CA-C	-5.10	103.36	109.83
1	D	289	ASN	N-CA-C	-5.09	105.42	111.69
1	B	133	SER	N-CA-C	-5.07	103.36	110.35
1	A	133	SER	N-CA-C	-5.03	103.41	110.35
1	D	156	ALA	CA-C-N	-5.02	115.06	120.03
1	D	156	ALA	C-N-CA	-5.02	115.06	120.03

There are no chirality outliers.

All (1) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	D	110	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2276	0	2058	28	0
1	B	2267	0	2052	12	0
1	C	2267	0	2052	16	0
1	D	2267	0	2052	13	1
2	D	5	0	0	0	0
3	A	186	0	0	5	0
3	B	187	0	0	0	0
3	C	151	0	0	1	0
3	D	166	0	0	2	0
All	All	9772	0	8214	65	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:THR:HB	1:A:198:ILE:HG12	1.60	0.84
1:B:170:THR:HG21	1:C:140:ASN:HD22	1.54	0.71
1:D:151:ASN:H	1:D:159:GLN:HE22	1.40	0.69
1:A:-1:GLU:O	1:A:0:PHE:HB2	1.93	0.66
1:C:226:LYS:HE2	3:C:421:HOH:O	1.96	0.65
1:B:170:THR:HG21	1:C:140:ASN:ND2	2.11	0.64
1:D:189:THR:HB	1:D:198:ILE:HG12	1.80	0.64
1:C:151:ASN:H	1:C:159:GLN:HE22	1.44	0.64
1:A:79:HIS:HE1	1:A:235:GLU:OE1	1.84	0.61
1:C:207:PRO:HG2	1:C:210:VAL:HB	1.83	0.60
1:D:189:THR:HG23	3:D:482:HOH:O	2.00	0.60
1:A:-1:GLU:O	1:A:0:PHE:CB	2.52	0.57
1:A:189:THR:HG23	3:A:407:HOH:O	2.05	0.57
1:A:116:ILE:HD13	1:A:189:THR:HG21	1.85	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:ASN:H	1:A:159:GLN:HE22	1.52	0.56
1:A:116:ILE:CD1	1:A:189:THR:HG21	2.36	0.55
1:B:151:ASN:H	1:B:159:GLN:HE22	1.55	0.55
1:D:148:LYS:HA	1:D:148:LYS:HE2	1.88	0.54
1:A:84:ILE:HG12	1:A:86:LEU:HD13	1.91	0.53
1:A:105:THR:HG23	1:A:240:ALA:O	2.10	0.51
1:B:79:HIS:CE1	1:B:235:GLU:OE1	2.65	0.50
1:A:9:ARG:NH1	1:A:46:GLU:OE2	2.45	0.50
1:A:210:VAL:HG13	1:A:211:LEU:HG	1.94	0.49
1:C:7:TYR:CD2	1:C:291:LEU:HD13	2.48	0.49
1:C:79:HIS:CE1	1:C:235:GLU:OE1	2.66	0.48
1:A:206:GLU:HB2	1:A:210:VAL:HG11	1.96	0.48
1:A:79:HIS:CE1	1:A:235:GLU:OE1	2.65	0.47
1:B:71:LEU:O	1:B:244:ILE:HA	2.15	0.47
1:D:140:ASN:O	1:D:140:ASN:CG	2.56	0.47
1:C:13:PHE:CD2	1:C:51:THR:HG21	2.49	0.47
1:A:232:CYS:HB3	1:A:234:TRP:CE2	2.50	0.46
1:A:189:THR:CG2	3:A:407:HOH:O	2.62	0.46
1:A:141:ASP:HA	3:A:442:HOH:O	2.15	0.46
1:A:272:ASP:OD1	1:D:141:ASP:OD2	2.34	0.46
1:B:79:HIS:HE1	1:B:235:GLU:OE1	1.98	0.46
1:C:232:CYS:HB3	1:C:234:TRP:CE2	2.51	0.45
1:A:217:PRO:HA	1:A:220:TRP:CD2	2.52	0.45
1:A:147:LEU:HD21	1:A:150:SER:OG	2.16	0.45
1:A:206:GLU:HA	1:A:207:PRO:HD3	1.84	0.44
1:B:140:ASN:O	1:B:140:ASN:CG	2.59	0.44
1:D:89:MET:HG2	1:D:179:PHE:CD1	2.52	0.44
1:B:246:ASP:OD1	1:B:246:ASP:C	2.61	0.44
1:D:239:ASN:HB2	3:D:515:HOH:O	2.18	0.44
1:D:287:LEU:HD12	1:D:287:LEU:HA	1.88	0.44
1:C:4:VAL:HG22	1:C:293:VAL:O	2.17	0.43
1:C:71:LEU:O	1:C:244:ILE:HA	2.19	0.43
1:C:130:LEU:HB2	1:C:165:CYS:SG	2.58	0.43
1:D:79:HIS:HE1	1:D:235:GLU:OE1	2.02	0.43
1:A:256:ASN:ND2	3:A:433:HOH:O	2.50	0.42
1:D:89:MET:HE1	1:D:175:TYR:CE1	2.55	0.42
1:B:113:GLU:HB3	1:B:131:HIS:HB2	2.00	0.42
1:A:246:ASP:OD1	1:A:246:ASP:C	2.62	0.42
1:C:153:TYR:CE2	1:C:155:TYR:HB2	2.55	0.41
1:A:158:GLY:HA3	1:B:256:ASN:HD21	1.86	0.41
1:B:235:GLU:OE1	1:B:235:GLU:HA	2.20	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:79:HIS:CE1	1:D:235:GLU:OE1	2.74	0.41
1:C:140:ASN:O	1:C:140:ASN:OD1	2.39	0.41
1:A:235:GLU:OE1	1:A:235:GLU:HA	2.21	0.40
1:C:232:CYS:HB3	1:C:234:TRP:CZ2	2.56	0.40
1:A:78:GLU:HG2	1:A:79:HIS:CD2	2.57	0.40
1:A:127:HIS:HB2	3:A:477:HOH:O	2.21	0.40
1:B:217:PRO:HA	1:B:220:TRP:CD2	2.57	0.40
1:A:116:ILE:HD13	1:A:189:THR:CG2	2.49	0.40
1:C:159:GLN:O	1:C:160:ASP:C	2.63	0.40
1:D:50:ARG:HE	1:D:289:ASN:HD21	1.69	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:160:ASP:OD2	1:D:160:ASP:OD2[4_555]	2.11	0.09

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/298 (99%)	290 (98%)	5 (2%)	1 (0%)	36	27
1	B	295/298 (99%)	291 (99%)	4 (1%)	0	100	100
1	C	295/298 (99%)	289 (98%)	5 (2%)	1 (0%)	36	27
1	D	295/298 (99%)	288 (98%)	7 (2%)	0	100	100
All	All	1181/1192 (99%)	1158 (98%)	21 (2%)	2 (0%)	43	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	0	PHE
1	C	157	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	226 (98%)	5 (2%)	45	38
1	B	230/231 (100%)	224 (97%)	6 (3%)	40	32
1	C	230/231 (100%)	227 (99%)	3 (1%)	61	57
1	D	230/231 (100%)	226 (98%)	4 (2%)	53	48
All	All	921/924 (100%)	903 (98%)	18 (2%)	48	42

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	LEU
1	A	109	PRO
1	A	157	PRO
1	A	189	THR
1	A	291	LEU
1	B	5	ASP
1	B	86	LEU
1	B	90	PRO
1	B	109	PRO
1	B	177	LYS
1	B	291	LEU
1	C	157	PRO
1	C	226	LYS
1	C	291	LEU
1	D	5	ASP
1	D	109	PRO
1	D	189	THR
1	D	246	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	ASN
1	A	79	HIS
1	A	146	ASN
1	A	159	GLN
1	A	256	ASN
1	B	2	HIS
1	B	75	ASN
1	B	146	ASN
1	B	159	GLN
1	B	194	ASN
1	B	256	ASN
1	C	2	HIS
1	C	75	ASN
1	C	79	HIS
1	C	140	ASN
1	C	159	GLN
1	C	194	ASN
1	D	75	ASN
1	D	79	HIS
1	D	159	GLN
1	D	269	ASN
1	D	289	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	D	301	-	4,4,4	0.34	0	6,6,6	0.22	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	298/298 (100%)	-0.42	3 (1%) 79 86	11, 18, 28, 53	0
1	B	297/298 (99%)	-0.35	1 (0%) 90 93	11, 19, 30, 45	0
1	C	297/298 (99%)	-0.22	4 (1%) 75 82	12, 21, 37, 45	0
1	D	297/298 (99%)	-0.31	0 100 100	12, 20, 33, 42	0
All	All	1189/1192 (99%)	-0.33	8 (0%) 84 89	11, 19, 33, 53	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	0	PHE	4.2
1	B	0	PHE	3.3
1	C	157	PRO	3.1
1	C	158	GLY	2.9
1	A	158	GLY	2.7
1	C	110	TYR	2.5
1	C	141	ASP	2.4
1	A	-1	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	D	301	5/5	0.92	0.09	61,61,62,62	0

## 6.5 Other polymers

There are no such residues in this entry.