



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 02:21 AM UTC

PDB ID : 3WII / pdb\_00003wii  
Title : Crystal structure of the Fab fragment of B2212A, a murine monoclonal antibody specific for the third fibronectin domain (Fn3) of human ROBO1.  
Authors : Nakayama, T.; Mizohata, E.; Yamashita, T.; Nagatoishi, S.; Nakakido, M.; Iwanari, H.; Mochizuki, Y.; Kado, Y.; Yokota, Y.; Sato, R.; Tsumoto, K.; Fujitani, H.; Kodama, T.; Hamakubo, T.; Inoue, T.  
Deposited on : 2013-09-12  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	213	<div> <div>4%</div> <div>84%</div> <div>15%</div> <div>.</div> </div>
1	M	213	<div> <div>2%</div> <div>92%</div> <div>7%</div> <div>.</div> </div>
2	H	220	<div> <div>5%</div> <div>92%</div> <div>7%</div> <div>.</div> </div>
2	I	220	<div> <div>5%</div> <div>90%</div> <div>8%</div> <div>.</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called anti-human ROBO1 antibody B2212A Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	213	Total	C	N	O	S	0	1	0
			1658	1030	280	341	7			
1	M	213	Total	C	N	O	S	0	1	0
			1658	1030	280	341	7			

- Molecule 2 is a protein called anti-human ROBO1 antibody B2212A Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	220	Total	C	N	O	S	0	2	0
			1680	1064	270	338	8			
2	I	220	Total	C	N	O	S	0	1	0
			1677	1062	270	338	7			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	122	Total	O	0	0
			122	122		
3	H	174	Total	O	0	0
			174	174		
3	M	149	Total	O	0	0
			149	149		
3	I	218	Total	O	0	0
			218	218		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: anti-human ROBO1 antibody B2212A Fab light chain




- Molecule 1: anti-human ROBO1 antibody B2212A Fab light chain



- Molecule 2: anti-human ROBO1 antibody B2212A Fab heavy chain



- Molecule 2: anti-human ROBO1 antibody B2212A Fab heavy chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.82Å 136.49Å 77.51Å 90.00° 91.86° 90.00°	Depositor
Resolution (Å)	14.91 – 1.60 14.91 – 1.60	Depositor EDS
% Data completeness (in resolution range)	94.1 (14.91-1.60) 94.0 (14.91-1.60)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.09 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.5.0109, PHENIX 1.8.4_1496	Depositor
R, $R_{free}$	0.185 , 0.230 0.190 , 0.235	Depositor DCC
$R_{free}$ test set	5363 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.5	Xtriage
Anisotropy	0.191	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 51.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.035 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7336	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.43	0/1694	0.75	1/2297 (0.0%)
1	M	0.54	0/1694	0.77	0/2297
2	H	0.48	0/1730	0.78	0/2366
2	I	0.53	0/1724	0.83	2/2358 (0.1%)
All	All	0.50	0/6842	0.78	3/9318 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	I	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	I	133	VAL	CA-C-N	6.74	133.84	121.70
2	I	133	VAL	C-N-CA	6.74	133.84	121.70
1	L	212	ASN	N-CA-C	5.23	117.94	109.83

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	I	179	ASP	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1658	0	1600	27	0
1	M	1658	0	1600	14	0
2	H	1680	0	1631	11	0
2	I	1677	0	1627	14	0
3	H	174	0	0	1	0
3	I	218	0	0	3	2
3	L	122	0	0	4	0
3	M	149	0	0	3	2
All	All	7336	0	6458	63	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:149:LYS:NZ	1:L:195:GLU:OE2	2.19	0.76
2:I:176:LEU:HD22	2:I:179:ASP:HA	1.67	0.74
1:L:110:GLU:OE1	3:L:375:HOH:O	2.06	0.72
2:I:133:VAL:HA	2:I:134:CYS:HB2	1.71	0.71
1:M:213:GLU:OE2	3:M:307:HOH:O	2.08	0.71
1:M:138:ASN:HA	1:M:172:THR:HG23	1.73	0.71
1:M:190:ASN:OD1	1:M:213:GLU:HG2	1.91	0.70
2:H:44:SER:OG	3:H:330:HOH:O	2.11	0.68
1:L:195:GLU:HG2	1:L:206:VAL:HG22	1.77	0.66
1:L:149:LYS:HB2	1:L:193:THR:HB	1.79	0.65
1:M:195:GLU:HG2	1:M:206:VAL:HG22	1.77	0.65
1:M:70:ASP:OD2	3:M:395:HOH:O	2.16	0.63
2:H:133:VAL:O	2:H:135:GLY:N	2.32	0.63
1:L:150:ILE:HD11	1:L:179:LEU:HD21	1.79	0.63
2:I:133:VAL:HG23	2:I:136:ASP:HB2	1.87	0.57
1:L:211:ARG:O	1:L:212:ASN:ND2	2.36	0.57
1:M:138:ASN:HA	1:M:172:THR:CG2	2.35	0.56
1:L:190:ASN:ND2	1:L:210:ASN:OD1	2.39	0.56
2:I:133:VAL:CA	2:I:134:CYS:HB2	2.35	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:146:VAL:HB	1:L:175:MET:HE1	1.88	0.56
1:L:24:ARG:NH1	3:L:333:HOH:O	2.36	0.55
1:L:210:ASN:O	1:L:212:ASN:N	2.40	0.55
2:H:212:VAL:HG22	2:I:210:THR:HG22	1.89	0.54
1:L:33:LEU:HD13	1:L:71:PHE:CG	2.43	0.54
1:L:122:SER:O	1:L:126:THR:HG23	2.07	0.53
2:I:65:ARG:NH2	3:I:457:HOH:O	2.24	0.53
1:L:211:ARG:C	1:L:212:ASN:HD22	2.16	0.53
2:H:176:LEU:HG	2:H:179:ASP:HA	1.91	0.53
1:L:119:PRO:HD3	2:H:133:VAL:HG12	1.91	0.52
1:L:37:GLN:HB2	1:L:47:LEU:HD11	1.92	0.52
1:L:33:LEU:HD13	1:L:71:PHE:CD1	2.45	0.51
1:L:211:ARG:HH11	1:L:211:ARG:HG3	1.75	0.51
2:I:20:ILE:HD11	2:I:81:MET:HE2	1.93	0.51
1:M:37:GLN:HB2	1:M:47:LEU:HD11	1.93	0.50
2:H:133:VAL:HG22	2:H:136:ASP:OD2	2.11	0.50
1:M:212:ASN:OD1	1:M:213:GLU:N	2.45	0.49
1:M:16:GLY:HA2	1:M:77:LYS:HG3	1.94	0.48
2:I:195:PRO:HB3	2:I:218:PRO:HG3	1.96	0.48
2:H:194:TRP:CG	2:H:195:PRO:HA	2.48	0.48
1:M:170:ASP:OD1	1:M:172:THR:HB	2.14	0.48
1:L:186:TYR:HE2	1:L:211:ARG:NH1	2.12	0.47
1:L:123:GLU:OE1	3:L:323:HOH:O	2.20	0.46
2:H:120:ALA:HB2	2:H:179:ASP:OD1	2.16	0.45
1:M:160:LEU:HB3	2:I:175:ILE:HD12	1.98	0.44
2:I:177:GLN:NE2	3:I:401:HOH:O	2.50	0.44
3:M:301:HOH:O	2:I:44:SER:HB3	2.16	0.44
2:I:35:ASN:OD1	2:I:50:ASP:HB3	2.17	0.44
1:M:193:THR:HG22	1:M:195:GLU:HG3	2.00	0.44
2:I:135:GLY:HA3	3:I:315:HOH:O	2.18	0.43
1:L:23:CYS:HB2	1:L:35:TRP:CH2	2.53	0.43
1:L:163:TRP:HD1	3:L:396:HOH:O	2.01	0.43
2:I:208:SER:O	2:I:210:THR:HG23	2.19	0.43
2:H:125:PRO:HB3	2:H:151:TYR:HB3	2.01	0.43
1:L:136:LEU:HD22	1:L:175:MET:HE2	1.99	0.43
1:L:110:GLU:HG2	1:L:141:PRO:HD3	2.00	0.42
1:M:184:ASP:O	1:M:188:ARG:HG3	2.19	0.42
1:L:16:GLY:HA2	1:L:77:LYS:HG3	2.00	0.42
1:M:212:ASN:OD1	1:M:212:ASN:C	2.63	0.41
1:L:211:ARG:HG3	1:L:212:ASN:HD22	1.86	0.41
2:H:43:LYS:HB3	2:H:43:LYS:HE2	1.76	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:120:PRO:HD3	1:L:132:VAL:HG22	2.02	0.40
1:L:110:GLU:OE2	1:L:199:LYS:NZ	2.27	0.40
2:H:35:ASN:OD1	2:H:50:ASP:HB3	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:M:411:HOH:O	3:I:475:HOH:O[1_655]	2.14	0.06
3:M:410:HOH:O	3:I:476:HOH:O[1_655]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	212/213 (100%)	204 (96%)	7 (3%)	1 (0%)	24	10
1	M	212/213 (100%)	206 (97%)	5 (2%)	1 (0%)	24	10
2	H	220/220 (100%)	213 (97%)	4 (2%)	3 (1%)	9	1
2	I	219/220 (100%)	209 (95%)	7 (3%)	3 (1%)	9	1
All	All	863/866 (100%)	832 (96%)	23 (3%)	8 (1%)	14	3

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	211	ARG
2	I	134	CYS
2	I	137	THR
2	I	179	ASP
2	H	134	CYS
2	H	135	GLY

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	M	212	ASN
2	H	179	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	192/191 (100%)	190 (99%)	2 (1%)	68	50
1	M	192/191 (100%)	190 (99%)	2 (1%)	68	50
2	H	194/192 (101%)	192 (99%)	2 (1%)	68	50
2	I	193/192 (100%)	191 (99%)	2 (1%)	68	50
All	All	771/766 (101%)	763 (99%)	8 (1%)	70	50

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	194[A]	CYS
1	L	194[B]	CYS
2	H	54	ASN
2	H	99	PHE
1	M	172	THR
1	M	212	ASN
2	I	99	PHE
2	I	183	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	3	GLN
1	L	80	GLN
1	L	212	ASN
2	H	54	ASN
1	M	27	GLN
1	M	92	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	M	145	ASN
2	I	3	GLN
2	I	41	HIS
2	I	177	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	213/213 (100%)	0.46	8 (3%) 44 46	23, 34, 66, 146	3 (1%)
1	M	213/213 (100%)	0.13	5 (2%) 61 64	17, 27, 54, 135	2 (0%)
2	H	220/220 (100%)	0.35	11 (5%) 34 35	15, 28, 72, 143	2 (0%)
2	I	220/220 (100%)	0.14	12 (5%) 30 31	16, 24, 66, 168	1 (0%)
All	All	866/866 (100%)	0.27	36 (4%) 40 42	15, 29, 63, 168	8 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	42	GLY	5.6
2	H	135	GLY	5.6
2	I	135	GLY	5.0
2	H	134	CYS	4.9
2	I	41	HIS	4.8
2	I	137	THR	4.7
2	I	220	ALA	4.5
2	I	134	CYS	4.0
2	H	138	THR	3.9
2	H	41	HIS	3.9
2	I	104	TYR	3.8
2	I	138	THR	3.8
1	M	68	GLY	3.7
2	H	137	THR	3.6
1	M	212	ASN	3.6
2	I	42	GLY	3.5
1	L	212	ASN	3.3
2	I	139	GLY	3.2
1	L	127	SER	3.1
2	H	133	VAL	3.0
2	I	178	SER	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	I	179	ASP	2.9
2	H	179	ASP	2.8
1	L	191	SER	2.8
2	H	175	ILE	2.7
2	H	220	ALA	2.6
1	M	213	GLU	2.5
1	L	149	LYS	2.5
1	L	211	ARG	2.4
1	L	190	ASN	2.3
1	M	67	SER	2.3
1	M	138	ASN	2.2
1	L	175	MET	2.2
2	H	176	LEU	2.1
1	L	153	SER	2.0
2	I	101	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.