Full wwPDB X-ray Structure Validation Report ⓘ

Nov 11, 2026 – 08:13 PM JST

PDB ID	:	47LZ / pdb_000047lz
Title	:	Structure of the Human ROS1 Kinase Domain with the G2032R Solvent-Front Mutation in Complex with Lorlatinib
Authors	:	Zhang, K.; Yang, Y.; Chen, R.; Fu, L.; Cheng, H.; Sun, X.; Zhou, L.; Wei, X.
Deposited on	:	2026-10-26
Resolution	:	3.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1605 (3.30-3.22)
Clashscore	190562	1660 (3.30-3.22)
Ramachandran outliers	187476	1630 (3.30-3.22)
Sidechain outliers	187428	1629 (3.30-3.22)
RSRZ outliers	180081	1605 (3.30-3.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	8% 76% 14% 10%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2437 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase ROS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	299	2406	1550	407	436	13	0	0	0

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1901	MET	-	initiating methionine	UNP P08922
A	1902	GLY	-	expression tag	UNP P08922
A	1903	SER	-	expression tag	UNP P08922
A	1904	SER	-	expression tag	UNP P08922
A	1905	HIS	-	expression tag	UNP P08922
A	1906	HIS	-	expression tag	UNP P08922
A	1907	HIS	-	expression tag	UNP P08922
A	1908	HIS	-	expression tag	UNP P08922
A	1909	HIS	-	expression tag	UNP P08922
A	1910	HIS	-	expression tag	UNP P08922
A	1911	SER	-	expression tag	UNP P08922
A	1912	SER	-	expression tag	UNP P08922
A	1913	GLY	-	expression tag	UNP P08922
A	1914	TRP	-	expression tag	UNP P08922
A	1915	SER	-	expression tag	UNP P08922
A	1916	HIS	-	expression tag	UNP P08922
A	1917	PRO	-	expression tag	UNP P08922
A	1918	GLN	-	expression tag	UNP P08922
A	1919	PHE	-	expression tag	UNP P08922
A	1920	GLU	-	expression tag	UNP P08922
A	1921	LYS	-	expression tag	UNP P08922
A	1922	SER	-	expression tag	UNP P08922
A	1923	ALA	-	expression tag	UNP P08922
A	1924	VAL	-	expression tag	UNP P08922
A	1925	ASP	-	expression tag	UNP P08922
A	1926	GLU	-	expression tag	UNP P08922
A	1927	ASN	-	expression tag	UNP P08922

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1928	LEU	-	expression tag	UNP P08922
A	1929	TYR	-	expression tag	UNP P08922
A	1930	PHE	-	expression tag	UNP P08922
A	1931	GLN	-	expression tag	UNP P08922
A	1932	GLY	-	expression tag	UNP P08922
A	1933	PRO	-	expression tag	UNP P08922
A	2032	ARG	GLY	engineered mutation	UNP P08922

- Molecule 2 is (10R)-7-amino-12-fluoro-2,10,16-trimethyl-15-oxo-10,15,16,17-tetrahydro-2H-8,4-(metheno)pyrazolo[4,3-h][2,5,11]benzoxadiazacyclotetradecine-3-carbonitrile (CCD ID: 5P8) (formula: $C_{21}H_{19}FN_6O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			30	21	1	6	2		


- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Proto-oncogene tyrosine-protein kinase ROS



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α, β, γ	106.68Å 106.68Å 103.26Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.39 – 3.25 47.39 – 3.25	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.39-3.25) 99.8 (47.39-3.25)	Depositor EDS
R_{merge}	0.27	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.86 (at 3.25Å)	Xtriage
Refinement program	PHENIX 2.1_6048	Depositor
R, R_{free}	0.220 , 0.247 0.220 , 0.247	Depositor DCC
R_{free} test set	524 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å^2)	63.7	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 66.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.033 for -h,-k,l	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	2437	wwPDB-VP
Average B, all atoms (Å^2)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5P8, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.13	0/2462	0.33	0/3337

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2406	0	2409	27	0
2	A	30	0	19	0	0
3	A	1	0	0	0	0
All	All	2437	0	2428	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2196:GLN:HG3	1:A:2206:PRO:HD3	1.72	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1982:LEU:HD23	1:A:2022:GLN:HG3	1.79	0.65
1:A:1930:PHE:HB3	1:A:1996:LYS:HB3	1.85	0.59
1:A:1993:GLU:HB3	1:A:2107:ARG:HH12	1.69	0.58
1:A:1982:LEU:HD21	1:A:1991:LYS:HG2	1.88	0.56
1:A:1925:ASP:HB2	1:A:1928:LEU:HG	1.89	0.53
1:A:1988:ASP:O	1:A:1992:ILE:HG12	2.09	0.53
1:A:1951:LEU:HD21	1:A:1961:GLU:HB2	1.94	0.49
1:A:2045:THR:HG22	1:A:2047:TYR:H	1.78	0.48
1:A:1930:PHE:CE1	1:A:2107:ARG:HG2	2.49	0.47
1:A:2191:TRP:HA	1:A:2194:MET:HE3	1.96	0.47
1:A:2166:SER:O	1:A:2170:VAL:HG23	2.15	0.47
1:A:1968:LEU:HG	1:A:1975:ILE:HD13	1.97	0.46
1:A:1964:ALA:HB3	1:A:1967:ILE:CD1	2.46	0.46
1:A:1964:ALA:HB3	1:A:1967:ILE:HD12	1.98	0.46
1:A:2020:GLU:HG2	1:A:2021:PRO:HA	1.97	0.45
1:A:2001:MET:HE1	1:A:2103:PHE:HD1	1.81	0.44
1:A:1994:PHE:CE2	1:A:2024:ILE:HG13	2.53	0.44
1:A:1967:ILE:HG21	1:A:2013:LEU:HB3	1.99	0.43
1:A:2063:ILE:HD11	1:A:2098:VAL:HG22	2.01	0.43
1:A:1943:GLU:H	1:A:1943:GLU:HG2	1.69	0.43
1:A:1983:LYS:O	1:A:1986:SER:HB2	2.19	0.43
1:A:1987:THR:HB	1:A:1990:GLU:HG3	2.01	0.42
1:A:2065:LYS:HG2	1:A:2212:GLN:NE2	2.35	0.42
1:A:2012:GLN:HA	1:A:2026:LEU:HD23	2.01	0.41
1:A:2182:PRO:HB3	1:A:2191:TRP:CD2	2.56	0.41
1:A:2144:VAL:HG13	1:A:2211:ILE:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	295/332 (89%)	283 (96%)	12 (4%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	264/294 (90%)	264 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2167	ASN
1	A	2212	GLN
1	A	2216	GLN
1	A	2224	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	5P8	A	2301	-	32,33,33	0.79	0	39,49,49	1.21	5 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5P8	A	2301	-	-	0/24/26/26	0/3/4/4

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	2301	5P8	C28-N26-C27	3.15	127.94	119.97
2	A	2301	5P8	O7-C8-C10	2.51	112.38	108.00
2	A	2301	5P8	O30-C27-N26	-2.39	119.54	122.41
2	A	2301	5P8	C29-N26-C27	-2.15	116.95	122.96
2	A	2301	5P8	C4-C5-C18	2.04	123.73	120.79

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 5P8 A 2301



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	299/332 (90%)	0.44	28 (9%)	14	11	24, 55, 114, 229	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1956	PHE	5.1
1	A	2109	ILE	4.5
1	A	2110	TYR	4.4
1	A	2227	TYR	4.0
1	A	1926	GLU	3.7
1	A	2121	GLY	3.7
1	A	2113	ASP	3.4
1	A	2122	LEU	3.2
1	A	2175	GLN	3.1
1	A	1949	LEU	2.8
1	A	2112	ASN	2.6
1	A	1927	ASN	2.6
1	A	2120	GLU	2.6
1	A	2226	ILE	2.5
1	A	1954	GLY	2.5
1	A	2111	LYS	2.5
1	A	1958	GLU	2.5
1	A	2105	LEU	2.4
1	A	2119	GLY	2.4
1	A	1928	LEU	2.3
1	A	2045	THR	2.3
1	A	2167	ASN	2.3
1	A	1925	ASP	2.2
1	A	2018	LEU	2.1
1	A	2140	THR	2.1
1	A	1955	ALA	2.1
1	A	1983	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2134	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	5P8	A	2301	30/30	0.95	0.10	32,42,52,65	0
3	MN	A	2302	1/1	0.99	0.04	30,30,30,30	1


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 5P8 A 2301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around MN A 2302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.