



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:29 AM UTC

PDB ID : 4CCG / pdb\_00004ccg  
Title : Structure of an E2-E3 complex  
Authors : Hodson, C.; Purkiss, A.; Walden, H.  
Deposited on : 2013-10-22  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	<div> <div>5%</div> <div>56%</div> <div>18%</div> <div>26%</div> </div>
1	B	212	<div> <div>3%</div> <div>54%</div> <div>15%</div> <div>31%</div> </div>
2	X	88	<div> <div>2%</div> <div>75%</div> <div>10%</div> <div>15%</div> </div>
2	Y	88	<div> <div>16%</div> <div>63%</div> <div>14%</div> <div>24%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EPE	A	1154[A]	-	-	X	-
4	GOL	A	1157	-	-	X	-

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 3744 atoms, of which 40 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UBIQUITIN-CONJUGATING ENZYME E2 T.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	157	Total	C	N	O	S	0	3	0
			1247	805	211	224	7			
1	B	147	Total	C	N	O	S	0	4	0
			1154	751	204	194	5			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	THR	-	expression tag	UNP Q9NW38
A	-13	GLY	-	expression tag	UNP Q9NW38
A	-12	SER	-	expression tag	UNP Q9NW38
A	-11	THR	-	expression tag	UNP Q9NW38
A	-10	GLY	-	expression tag	UNP Q9NW38
A	-9	SER	-	expression tag	UNP Q9NW38
A	-8	THR	-	expression tag	UNP Q9NW38
A	-7	GLY	-	expression tag	UNP Q9NW38
A	-6	GLU	-	expression tag	UNP Q9NW38
A	-5	ASN	-	expression tag	UNP Q9NW38
A	-4	LEU	-	expression tag	UNP Q9NW38
A	-3	TYR	-	expression tag	UNP Q9NW38
A	-2	THR	-	expression tag	UNP Q9NW38
A	-1	GLN	-	expression tag	UNP Q9NW38
A	0	GLY	-	expression tag	UNP Q9NW38
B	-14	THR	-	expression tag	UNP Q9NW38
B	-13	GLY	-	expression tag	UNP Q9NW38
B	-12	SER	-	expression tag	UNP Q9NW38
B	-11	THR	-	expression tag	UNP Q9NW38
B	-10	GLY	-	expression tag	UNP Q9NW38
B	-9	SER	-	expression tag	UNP Q9NW38
B	-8	THR	-	expression tag	UNP Q9NW38
B	-7	GLY	-	expression tag	UNP Q9NW38
B	-6	GLU	-	expression tag	UNP Q9NW38
B	-5	ASN	-	expression tag	UNP Q9NW38

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	LEU	-	expression tag	UNP Q9NW38
B	-3	TYR	-	expression tag	UNP Q9NW38
B	-2	THR	-	expression tag	UNP Q9NW38
B	-1	GLN	-	expression tag	UNP Q9NW38
B	0	GLY	-	expression tag	UNP Q9NW38

- Molecule 2 is a protein called E3 UBIQUITIN-PROTEIN LIGASE FANCL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	75	Total	C	N	O	S	0	4	0
			614	389	100	116	9			
2	Y	67	Total	C	N	O	S	0	0	0
			512	328	80	95	9			

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula:  $C_8H_{18}N_2O_4S$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	1
			15	8	2	4	1		
3	A	1	Total	C	H	N	O	S	0
			11	2	4	1	3	1	

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	A	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	X	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	1
5	Y	1	Total O S 5 4 1	0	0

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Cl 1 1	0	0


- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total Na 1 1	0	0

- Molecule 8 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	X	2	Total Zn 2 2	0	0
8	Y	2	Total Zn 2 2	0	0

- Molecule 9 is AMMONIUM ION (CCD ID: NH4) (formula: H<sub>4</sub>N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	X	1	Total	H	N	0	0
			5	4	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	66	Total	O	0	0
			66	66		
10	B	23	Total	O	0	0
			23	23		
10	X	22	Total	O	0	0
			22	22		
10	Y	3	Total	O	0	0
			3	3		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: UBIQUITIN-CONJUGATING ENZYME E2 T



#### • Molecule 1: UBIQUITIN-CONJUGATING ENZYME E2 T




#### • Molecule 2: E3 UBIQUITIN-PROTEIN LIGASE FANCL



#### • Molecule 2: E3 UBIQUITIN-PROTEIN LIGASE FANCL





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	109.22Å 109.22Å 117.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.82 – 2.40 46.82 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.82-2.40) 94.4 (46.82-2.40)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.72 (at 2.34Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.212 , 0.248 0.227 , 0.260	Depositor DCC
$R_{free}$ test set	1518 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	58.9	Xtriage
Anisotropy	0.264	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 57.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.54$ , $\langle L^2 \rangle = 0.39$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3744	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, ZN, NA, CL, NH4, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/1288	0.73	0/1758
1	B	0.30	0/1199	0.77	0/1638
2	X	0.28	0/639	0.65	0/862
2	Y	0.27	0/524	0.72	0/710
All	All	0.30	0/3650	0.73	0/4968

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1247	0	1228	39	0
1	B	1154	0	1156	26	0
2	X	614	0	580	9	0
2	Y	512	0	467	10	0
3	A	22	4	23	12	0
4	A	12	16	16	5	0
4	B	6	8	7	0	0
4	X	6	8	8	0	0
5	A	5	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	Y	5	0	0	0	0
6	B	1	0	0	0	0
7	B	1	0	0	0	0
8	X	2	0	0	0	0
8	Y	2	0	0	0	0
9	X	1	4	0	0	0
10	A	66	0	0	1	0
10	B	23	0	0	0	0
10	X	22	0	0	0	0
10	Y	3	0	0	0	0
All	All	3704	40	3485	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:ARG:HH21	3:A:1154[A]:EPE:H81	1.46	0.81
2:X:316:ASP:OD2	2:X:373[A]:ARG:NH1	2.18	0.76
1:B:13:MET:HA	1:B:16[B]:THR:HG22	1.68	0.75
1:A:-2:THR:H	1:A:1:MET:HE3	1.55	0.70
1:A:34:LEU:HB2	1:A:55:VAL:HB	1.76	0.67
1:A:139:PHE:HD2	1:A:140[B]:LEU:HD12	1.63	0.64
1:A:9:ARG:HH21	3:A:1154[A]:EPE:C8	2.12	0.63
1:A:136:LYS:HB2	1:A:137:PRO:HD3	1.81	0.62
1:A:140[A]:LEU:HB3	4:A:1157:GOL:H2	1.81	0.62
2:Y:355:ILE:CG2	2:Y:368:LEU:HB3	2.30	0.62
1:B:5:SER:HB2	2:Y:312:ALA:HB2	1.80	0.62
1:A:10:GLU:OE1	1:A:103:ASN:HB2	2.00	0.61
1:A:141:LYS:HA	1:A:144:ARG:NH1	2.16	0.60
2:X:313:TYR:O	2:X:320:PRO:HD3	2.02	0.58
2:Y:364:LYS:HB3	2:Y:365:PRO:HD2	1.85	0.57
1:B:13:MET:HA	1:B:16[B]:THR:CG2	2.33	0.57
1:A:35:ARG:HH12	3:A:1155:EPE:C10	2.18	0.56
1:B:80:ASP:HA	1:B:125:MET:SD	2.46	0.56
1:B:132:PHE:CE1	1:B:136:LYS:HE3	2.41	0.56
1:A:144:ARG:HD3	4:A:1157:GOL:H31	1.88	0.55
1:A:23:THR:HG23	4:A:1156:GOL:H12	1.88	0.55
1:B:86:CYS:HB2	1:B:124:LEU:HG	1.89	0.55
2:Y:309:ILE:HG13	2:Y:333:PHE:CD1	2.41	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:136:LYS:N	1:B:137:PRO:HD2	2.22	0.54
1:A:40:GLY:HA3	1:A:46:TYR:O	2.08	0.54
1:A:-3:TYR:HB2	2:X:316:ASP:OD2	2.08	0.53
1:A:141:LYS:HA	1:A:144:ARG:HH12	1.74	0.53
1:B:14:LEU:HD13	1:B:24:CYS:HB3	1.90	0.53
2:Y:313:TYR:O	2:Y:320:PRO:HD3	2.09	0.52
1:B:26:GLN:HA	1:B:34:LEU:CD2	2.39	0.52
1:B:69:ARG:NH2	1:B:82:ALA:O	2.38	0.52
1:A:140[B]:LEU:HB3	4:A:1157:GOL:H2	1.91	0.52
2:X:355[B]:ILE:HG23	2:X:368:LEU:HB3	1.91	0.52
1:A:9:ARG:NH2	3:A:1154[A]:EPE:H81	2.20	0.52
1:A:70:PHE:CE1	1:A:79:ILE:HD13	2.45	0.52
1:B:7:LEU:O	1:B:11:LEU:HD23	2.10	0.51
1:A:144:ARG:HD3	4:A:1157:GOL:C3	2.41	0.50
2:X:355[B]:ILE:CG2	2:X:368:LEU:HB3	2.41	0.50
1:A:9:ARG:HH22	3:A:1154[A]:EPE:H102	1.76	0.50
1:A:136:LYS:O	1:A:140[B]:LEU:HD13	2.12	0.49
2:Y:355:ILE:HG23	2:Y:368:LEU:HB3	1.94	0.49
1:A:35:ARG:HH12	3:A:1155:EPE:H102	1.76	0.48
1:A:14:LEU:HD13	1:A:24:CYS:HB3	1.95	0.48
1:A:80:ASP:HA	1:A:125:MET:SD	2.54	0.47
1:A:7[A]:LEU:HD21	1:A:57:ILE:HG21	1.94	0.47
1:B:86:CYS:HB2	1:B:124:LEU:CG	2.45	0.47
1:B:140:LEU:O	1:B:144[A]:ARG:HG3	2.15	0.47
1:B:61:TYR:CD1	1:B:62:PRO:HA	2.49	0.47
1:A:43:ASN:HB3	10:A:2023:HOH:O	2.15	0.46
1:A:86:CYS:HB2	1:A:124:LEU:HG	1.98	0.46
1:B:12:HIS:O	1:B:16[B]:THR:HG22	2.16	0.46
1:B:13:MET:HE1	2:X:370:MET:HG3	1.98	0.46
3:A:1154[A]:EPE:H81	3:A:1154[A]:EPE:H31	1.51	0.45
2:X:318:THR:CG2	2:X:373[B]:ARG:HD3	2.46	0.45
1:A:9:ARG:NH2	5:A:1158[B]:SO4:O3	2.49	0.45
1:A:61:TYR:CD1	1:A:62:PRO:HA	2.52	0.45
1:A:44:THR:HB	1:A:45:PRO:HD2	1.99	0.44
1:B:12:HIS:O	1:B:16[A]:THR:HG23	2.17	0.44
1:A:35:ARG:HH12	3:A:1155:EPE:H101	1.82	0.44
1:B:6:ARG:HD2	2:Y:311:TYR:CD2	2.53	0.44
1:A:9:ARG:HH22	3:A:1154[A]:EPE:C10	2.30	0.44
1:B:6:ARG:HD3	2:Y:309:ILE:O	2.18	0.43
1:A:139:PHE:CD2	1:A:140[B]:LEU:HD12	2.48	0.43
1:A:9:ARG:HH21	3:A:1154[A]:EPE:H31	1.83	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:ARG:NH2	3:A:1154[A]:EPE:H31	2.33	0.43
2:Y:341:TRP:NE1	2:Y:345:LEU:HD11	2.33	0.43
1:A:13:MET:HE3	1:A:13:MET:HB3	1.97	0.43
1:B:17:GLU:HG2	2:X:370:MET:CE	2.49	0.42
1:B:90:LEU:HD21	1:B:107[B]:VAL:CG1	2.49	0.42
2:Y:309:ILE:HG13	2:Y:333:PHE:HD1	1.84	0.42
1:B:16[B]:THR:HG23	1:B:17:GLU:N	2.34	0.42
1:B:87:LEU:HG	1:B:89:VAL:HG22	2.02	0.42
1:B:17:GLU:CD	2:X:373[B]:ARG:HH22	2.23	0.42
1:B:34:LEU:HB2	1:B:55:VAL:HB	2.02	0.42
1:B:40:GLY:HA3	1:B:46:TYR:O	2.21	0.41
1:A:86:CYS:SG	1:A:122:ASP:HB3	2.60	0.41
1:A:76:HIS:HA	1:A:77:PRO:HD3	1.95	0.41
3:A:1154[A]:EPE:H101	3:A:1154[A]:EPE:H21	1.71	0.40
1:A:86:CYS:HB2	1:A:124:LEU:CG	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/212 (74%)	154 (98%)	4 (2%)	0	100	100
1	B	147/212 (69%)	146 (99%)	1 (1%)	0	100	100
2	X	76/88 (86%)	72 (95%)	4 (5%)	0	100	100
2	Y	63/88 (72%)	61 (97%)	2 (3%)	0	100	100
All	All	444/600 (74%)	433 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/186 (70%)	129 (98%)	2 (2%)	57	77
1	B	118/186 (63%)	117 (99%)	1 (1%)	73	86
2	X	69/79 (87%)	68 (99%)	1 (1%)	59	79
2	Y	55/79 (70%)	55 (100%)	0	100	100
All	All	373/530 (70%)	369 (99%)	4 (1%)	65	82

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	HIS
1	A	31	MET
1	B	136	LYS
2	X	299	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	GLN
1	A	150	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 6 are monoatomic and 1 is modelled with single atom - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	1157	-	5,5,5	0.36	0	5,5,5	0.21	0
4	GOL	A	1156	-	5,5,5	0.35	0	5,5,5	0.38	0
3	EPE	A	1155	-	6,6,15	3.22	1 (16%)	7,8,20	1.65	1 (14%)
5	SO4	A	1158[B]	-	4,4,4	0.24	0	6,6,6	0.09	0
3	EPE	A	1154[A]	-	15,15,15	2.08	1 (6%)	19,20,20	2.82	12 (63%)
4	GOL	X	1376	-	5,5,5	0.35	0	5,5,5	0.26	0
4	GOL	B	1154	7	5,5,5	0.38	0	5,5,5	0.20	0
5	SO4	Y	1374	-	4,4,4	0.24	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1157	-	-	4/4/4/4	-
4	GOL	A	1156	-	-	2/4/4/4	-
3	EPE	A	1155	-	-	0/4/4/19	-
3	EPE	A	1154[A]	-	-	3/9/19/19	0/1/1/1
4	GOL	X	1376	-	-	2/4/4/4	-
4	GOL	B	1154	7	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1154[A]	EPE	C10-S	-7.74	1.66	1.77
3	A	1155	EPE	C10-S	-7.69	1.66	1.77

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1154[A]	EPE	C6-N1-C2	5.24	120.13	108.84
3	A	1154[A]	EPE	C5-N4-C3	4.88	119.35	108.84
3	A	1154[A]	EPE	C7-N4-C5	3.44	120.40	111.24
3	A	1154[A]	EPE	C7-N4-C3	3.38	120.25	111.24
3	A	1154[A]	EPE	C6-C5-N4	-3.30	104.00	110.65
3	A	1154[A]	EPE	C9-N1-C6	3.28	119.98	111.24
3	A	1154[A]	EPE	C9-N1-C2	3.25	119.89	111.24
3	A	1154[A]	EPE	C2-C3-N4	-2.99	104.63	110.65
3	A	1154[A]	EPE	C5-C6-N1	-2.90	104.80	110.65
3	A	1155	EPE	C10-C9-N1	-2.47	105.97	112.39
3	A	1154[A]	EPE	C3-C2-N1	-2.30	106.02	110.65
3	A	1154[A]	EPE	C8-C7-N4	-2.21	105.69	113.44
3	A	1154[A]	EPE	C10-C9-N1	-2.01	104.77	112.36

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1154[A]	EPE	C10-C9-N1-C2
4	A	1157	GOL	O1-C1-C2-C3
4	B	1154	GOL	C1-C2-C3-O3
4	X	1376	GOL	C1-C2-C3-O3
3	A	1154[A]	EPE	C8-C7-N4-C3
4	A	1156	GOL	C1-C2-C3-O3
4	A	1156	GOL	O2-C2-C3-O3
4	B	1154	GOL	O2-C2-C3-O3
4	X	1376	GOL	O2-C2-C3-O3
4	A	1157	GOL	O1-C1-C2-O2
4	A	1157	GOL	O2-C2-C3-O3
4	A	1157	GOL	C1-C2-C3-O3
3	A	1154[A]	EPE	N4-C7-C8-O8

There are no ring outliers.

5 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1157	GOL	4	0
4	A	1156	GOL	1	0
3	A	1155	EPE	3	0
5	A	1158[B]	SO4	1	0
3	A	1154[A]	EPE	9	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	157/212 (74%)	0.15	10 (6%) 25 22	23, 47, 84, 116	3 (1%)
1	B	147/212 (69%)	0.46	7 (4%) 35 32	32, 65, 86, 114	4 (2%)
2	X	75/88 (85%)	0.11	2 (2%) 56 52	31, 47, 75, 89	4 (5%)
2	Y	67/88 (76%)	1.15	14 (20%) 2 2	52, 72, 101, 124	0
All	All	446/600 (74%)	0.40	33 (7%) 20 17	23, 59, 91, 124	11 (2%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	Y	351	SER	10.0
2	Y	303	PHE	5.1
2	Y	350	GLN	4.4
2	Y	371	SER	4.3
1	B	0	GLY	4.1
1	A	153	GLN	3.9
1	A	32	ASP	3.8
2	Y	304	THR	3.7
1	B	33	ASP	3.7
2	Y	354	ILE	3.6
2	X	299	GLU	3.4
1	A	31	MET	3.3
1	B	121	ASP	3.2
2	Y	316	ASP	3.0
1	A	29	ASP	3.0
2	Y	346	LEU	2.9
1	B	1	MET	2.9
1	A	28	LYS	2.8
1	B	122	ASP	2.7
1	B	120	PRO	2.6
2	Y	370	MET	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	Y	347	THR	2.6
1	A	152	ARG	2.5
2	X	300	LYS	2.5
2	Y	369	LYS	2.5
1	A	122	ASP	2.5
1	B	16[A]	THR	2.4
2	Y	339	TYR	2.4
1	A	121	ASP	2.3
1	A	120	PRO	2.3
2	Y	343	ARG	2.2
2	Y	321	ASP	2.1
1	A	30	GLN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EPE	A	1155	7/15	0.45	0.14	87,110,145,146	0
4	GOL	A	1156	6/6	0.55	0.23	75,90,99,102	0
5	SO4	Y	1374	5/5	0.59	0.12	147,148,149,149	0
4	GOL	A	1157	6/6	0.61	0.27	72,86,99,99	0
4	GOL	B	1154	6/6	0.69	0.17	80,96,100,100	0
3	EPE	A	1154[A]	15/15	0.69	0.28	40,70,96,96	15
4	GOL	X	1376	6/6	0.78	0.17	64,77,90,91	0
9	NH4	X	1377	1/1	0.83	0.13	67,80,80,80	0
5	SO4	A	1158[B]	5/5	0.87	0.17	89,92,93,93	5
7	NA	B	1155	1/1	0.90	0.22	65,65,65,65	0
6	CL	B	1153	1/1	0.97	0.07	72,72,72,72	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
8	ZN	Y	1372	1/1	0.99	0.03	63,63,63,63	0
8	ZN	X	1374	1/1	1.00	0.03	38,38,38,38	0
8	ZN	Y	1373	1/1	1.00	0.02	57,57,57,57	0
8	ZN	X	1375	1/1	1.00	0.04	48,48,48,48	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.