



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:35 PM UTC

PDB ID : 4HX3 / pdb_00004hx3
Title : Crystal structure of Streptomyces caespitosus sermetstatin in complex with S. caespitosus snapalysin
Authors : Trillo-Muyo, S.; Martinez-Rodriguez, S.; Arolas, J.L.; Gomis-Ruth, F.X.
Deposited on : 2012-11-09
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	134	<div> <div>%</div> <div>84% 16%</div> </div>
1	C	134	<div> <div>3%</div> <div>89% 10%</div> </div>
1	E	134	<div> <div>5%</div> <div>91% 7%</div> </div>
1	G	134	<div> <div>2%</div> <div>95%</div> </div>
1	I	134	<div> <div>%</div> <div>92% 8%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	134	<div><div></div><div>10%</div><div>93%</div><div>5% ..</div></div>
2	B	114	<div><div></div><div>2%</div><div>89%</div><div>11% .</div></div>
2	D	114	<div><div></div><div>6%</div><div>84%</div><div>11% . .</div></div>
2	F	114	<div><div></div><div>6%</div><div>88%</div><div>10% ..</div></div>
2	H	114	<div><div></div><div>6%</div><div>84%</div><div>10% . 5%</div></div>
2	J	114	<div><div></div><div>6%</div><div>84%</div><div>13% .</div></div>
2	L	114	<div><div></div><div>2%</div><div>91%</div><div>7% ..</div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11334 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Extracellular small neutral protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	134	Total	C	N	O	S	0	0	0
			1027	627	187	209	4			
1	C	134	Total	C	N	O	S	0	0	0
			1027	627	187	209	4			
1	E	132	Total	C	N	O	S	0	0	0
			1016	620	185	207	4			
1	G	133	Total	C	N	O	S	0	0	0
			1023	625	186	208	4			
1	I	134	Total	C	N	O	S	0	0	0
			1027	627	187	209	4			
1	K	133	Total	C	N	O	S	0	0	0
			1023	625	186	208	4			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P56406
A	-1	PRO	-	expression tag	UNP P56406
A	1	MET	-	expression tag	UNP P56406
C	-2	GLY	-	expression tag	UNP P56406
C	-1	PRO	-	expression tag	UNP P56406
C	1	MET	-	expression tag	UNP P56406
E	-1	GLY	-	expression tag	UNP P56406
E	0	PRO	-	expression tag	UNP P56406
E	1	MET	-	expression tag	UNP P56406
G	-2	GLY	-	expression tag	UNP P56406
G	-1	PRO	-	expression tag	UNP P56406
G	1	MET	-	expression tag	UNP P56406
I	-2	GLY	-	expression tag	UNP P56406
I	-1	PRO	-	expression tag	UNP P56406
I	1	MET	-	expression tag	UNP P56406
K	-2	GLY	-	expression tag	UNP P56406
K	-1	PRO	-	expression tag	UNP P56406

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
K	1	MET	-	expression tag	UNP P56406

- Molecule 2 is a protein called Neutral proteinase inhibitor ScNPI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	113	Total	C	N	O	S	0	0	0
			830	516	144	163	7			
2	D	109	Total	C	N	O	S	0	0	0
			802	499	140	156	7			
2	F	112	Total	C	N	O	S	0	0	0
			824	513	143	161	7			
2	H	108	Total	C	N	O	S	0	0	0
			796	495	139	155	7			
2	J	111	Total	C	N	O	S	0	0	0
			816	507	142	160	7			
2	L	113	Total	C	N	O	S	0	0	0
			830	516	144	163	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLY	-	expression tag	UNP Q9FDS0
D	0	GLY	-	expression tag	UNP Q9FDS0
F	0	GLY	-	expression tag	UNP Q9FDS0
H	-1	GLY	-	expression tag	UNP Q9FDS0
J	0	GLY	-	expression tag	UNP Q9FDS0
L	0	GLY	-	expression tag	UNP Q9FDS0

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	E	1	Total	Zn	0	0
			1	1		
3	G	1	Total	Zn	0	0
			1	1		
3	I	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	K	1	Total	Zn	0	0
			1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			6	3	3		
4	J	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	41	Total	O	0	0
			41	41		
5	B	28	Total	O	0	0
			28	28		
5	C	20	Total	O	0	0
			20	20		
5	D	16	Total	O	0	0
			16	16		
5	E	14	Total	O	0	0
			14	14		
5	F	25	Total	O	0	0
			25	25		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	30	Total 30	O 30	0	0
5	H	23	Total 23	O 23	0	0
5	I	29	Total 29	O 29	0	0
5	J	27	Total 27	O 27	0	0
5	K	2	Total 2	O 2	0	0
5	L	20	Total 20	O 20	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Extracellular small neutral protease



- Molecule 1: Extracellular small neutral protease



- Molecule 1: Extracellular small neutral protease




- Molecule 1: Extracellular small neutral protease




- Molecule 1: Extracellular small neutral protease




- Molecule 1: Extracellular small neutral protease




- Molecule 2: Neutral proteinase inhibitor ScNPI




- Molecule 2: Neutral proteinase inhibitor ScNPI




- Molecule 2: Neutral proteinase inhibitor ScNPI




- Molecule 2: Neutral proteinase inhibitor ScNPI



- Molecule 2: Neutral proteinase inhibitor ScNPI



- Molecule 2: Neutral proteinase inhibitor ScNPI



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	116.54Å 121.81Å 130.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.49 – 2.70 43.49 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.5 (43.49-2.70) 99.5 (43.49-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.69Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.195 , 0.242 0.200 , 0.240	Depositor DCC
R_{free} test set	784 reflections (1.51%)	wwPDB-VP
Wilson B-factor (Å ²)	48.3	Xtriage
Anisotropy	0.239	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 60.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.008 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11334	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.83	0/1055	1.20	3/1437 (0.2%)
1	C	0.82	0/1055	1.26	4/1437 (0.3%)
1	E	0.81	0/1043	1.22	1/1420 (0.1%)
1	G	0.88	0/1051	1.24	1/1431 (0.1%)
1	I	0.85	0/1055	1.21	2/1437 (0.1%)
1	K	0.79	0/1051	1.23	2/1431 (0.1%)
2	B	0.89	0/850	1.31	6/1166 (0.5%)
2	D	0.85	0/819	1.34	6/1120 (0.5%)
2	F	0.91	0/844	1.26	3/1158 (0.3%)
2	H	0.94	1/813 (0.1%)	1.23	0/1111
2	J	0.88	0/835	1.24	4/1144 (0.3%)
2	L	0.85	0/850	1.24	3/1166 (0.3%)
All	All	0.86	1/11321 (0.0%)	1.25	35/15458 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	8	MET	SD-CE	-5.12	1.66	1.79

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	64	ASP	CA-CB-CG	9.51	122.11	112.60
2	B	73	PHE	CA-C-N	8.42	129.69	122.28
2	B	73	PHE	C-N-CA	8.42	129.69	122.28
1	K	58	ASP	CA-CB-CG	7.75	120.35	112.60
2	D	106	ASN	CA-CB-CG	7.20	119.80	112.60
2	J	97	ASN	CA-C-N	6.40	128.85	120.28
2	J	97	ASN	C-N-CA	6.40	128.85	120.28
2	D	67	LEU	CA-C-N	6.29	129.00	120.63

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	67	LEU	C-N-CA	6.29	129.00	120.63
2	F	106	ASN	CA-CB-CG	6.13	118.73	112.60
1	E	58	ASP	CA-CB-CG	6.07	118.67	112.60
1	A	58	ASP	CA-CB-CG	6.05	118.65	112.60
2	D	47	ASP	CA-CB-CG	6.05	118.65	112.60
1	C	39	ASN	CA-C-N	5.90	129.66	120.75
1	C	39	ASN	C-N-CA	5.90	129.66	120.75
1	C	58	ASP	CA-CB-CG	5.77	118.37	112.60
1	G	58	ASP	CA-CB-CG	5.77	118.37	112.60
1	I	58	ASP	CA-CB-CG	5.74	118.34	112.60
2	J	75	PRO	CA-C-N	5.73	130.96	122.71
2	J	75	PRO	C-N-CA	5.73	130.96	122.71
1	C	20	ASN	CA-CB-CG	5.49	118.09	112.60
1	A	42	PHE	CA-CB-CG	5.29	119.09	113.80
2	B	102	SER	CA-C-N	5.29	127.31	120.44
2	B	102	SER	C-N-CA	5.29	127.31	120.44
2	F	60	LEU	CA-C-N	5.27	128.31	120.31
2	F	60	LEU	C-N-CA	5.27	128.31	120.31
1	I	42	PHE	CA-CB-CG	5.17	118.97	113.80
2	B	113	PHE	CA-CB-CG	5.13	118.93	113.80
1	A	38	GLY	N-CA-C	5.12	125.32	113.18
1	K	118	ASN	CA-CB-CG	5.09	117.69	112.60
2	L	97	ASN	CA-C-N	5.09	127.10	120.28
2	L	97	ASN	C-N-CA	5.09	127.10	120.28
2	D	44	ALA	CA-C-N	5.08	127.04	120.44
2	D	44	ALA	C-N-CA	5.08	127.04	120.44
2	B	21	ASP	CA-CB-CG	5.04	117.64	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1027	0	924	10	0
1	C	1027	0	924	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1016	0	914	6	0
1	G	1023	0	921	2	0
1	I	1027	0	924	6	0
1	K	1023	0	921	2	0
2	B	830	0	797	4	0
2	D	802	0	773	3	0
2	F	824	0	789	4	0
2	H	796	0	766	7	0
2	J	816	0	780	6	0
2	L	830	0	797	5	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
3	E	1	0	0	0	0
3	G	1	0	0	0	0
3	I	1	0	0	0	0
3	K	1	0	0	0	0
4	D	6	0	8	1	0
4	J	6	0	8	0	0
5	A	41	0	0	3	0
5	B	28	0	0	0	0
5	C	20	0	0	0	0
5	D	16	0	0	1	0
5	E	14	0	0	0	0
5	F	25	0	0	1	0
5	G	30	0	0	0	0
5	H	23	0	0	0	0
5	I	29	0	0	2	0
5	J	27	0	0	0	0
5	K	2	0	0	0	0
5	L	20	0	0	1	0
All	All	11334	0	10246	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:9:VAL:HG23	2:L:9:VAL:HG23	1.47	0.95
2:H:13:ILE:HG21	2:H:17:GLY:HA2	1.78	0.64
2:D:43:ARG:HB2	4:D:201:GOL:H11	1.82	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:2:VAL:HG23	1:I:132:GLY:HA2	1.85	0.58
2:F:4:GLY:H	2:F:85:ASN:HD22	1.52	0.58
2:H:4:GLY:H	2:H:85:ASN:HD22	1.50	0.57
2:F:62:ALA:O	2:F:106:ASN:HB3	2.04	0.57
2:J:4:GLY:H	2:J:85:ASN:HD22	1.53	0.57
2:J:63:PRO:HB3	2:J:103:ALA:HB1	1.88	0.56
2:B:67:LEU:HD23	2:B:103:ALA:HB2	1.88	0.55
1:A:-1:PRO:HB3	1:A:33:GLN:HG2	1.88	0.55
1:C:124:ARG:O	1:C:128:LEU:HG	2.07	0.55
1:E:2:VAL:HG11	1:E:129:TRP:HE3	1.72	0.54
1:C:8:PRO:HB3	1:C:18:ILE:HG21	1.90	0.54
2:B:74:ASP:O	2:B:97:ASN:HA	2.08	0.54
1:E:8:PRO:HB3	1:E:18:ILE:HG21	1.90	0.54
2:L:4:GLY:H	2:L:85:ASN:HD22	1.55	0.54
1:I:8:PRO:HB3	1:I:18:ILE:HG21	1.91	0.52
1:A:8:PRO:HB3	1:A:18:ILE:HG21	1.92	0.52
1:A:112:CYS:HB3	5:A:326:HOH:O	2.11	0.51
1:A:120:GLN:NE2	5:A:333:HOH:O	2.28	0.51
1:C:62:ARG:NH1	1:C:62:ARG:HB3	2.26	0.50
1:G:8:PRO:HB3	1:G:18:ILE:HG21	1.93	0.50
1:K:8:PRO:HB3	1:K:18:ILE:HG21	1.93	0.50
1:A:95:TYR:O	2:B:43:ARG:HD2	2.12	0.49
2:H:13:ILE:HD11	2:H:23:VAL:HG22	1.94	0.49
1:C:62:ARG:HB3	1:C:62:ARG:HH11	1.78	0.49
2:J:9:VAL:HG23	2:L:9:VAL:CG2	2.33	0.48
1:C:6:TYR:O	1:C:37:GLY:HA2	2.14	0.47
1:I:98:PRO:HB2	5:I:316:HOH:O	2.15	0.47
2:B:40:PRO:HD3	1:C:50:SER:HB3	1.97	0.46
1:I:94:HIS:NE2	5:I:329:HOH:O	2.35	0.46
1:K:17:GLU:H	1:K:17:GLU:CD	2.24	0.46
2:L:89:VAL:HG22	5:L:205:HOH:O	2.16	0.45
2:H:13:ILE:CD1	2:H:23:VAL:HG22	2.45	0.45
1:G:95:TYR:O	2:H:43:ARG:HD2	2.17	0.45
1:I:2:VAL:CG2	1:I:132:GLY:HA2	2.47	0.45
1:A:24:ILE:HD13	1:A:115:PRO:HB2	1.99	0.44
1:E:2:VAL:HG13	1:E:32:VAL:HG23	1.99	0.44
2:H:74:ASP:O	2:H:97:ASN:HA	2.18	0.44
2:F:74:ASP:O	2:F:97:ASN:HA	2.18	0.44
2:H:13:ILE:HD13	2:H:23:VAL:HA	2.01	0.43
1:C:49:ASP:HB2	1:C:66:PHE:CE2	2.54	0.43
1:A:98:PRO:HD2	1:A:101:GLU:HB2	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:95:PHE:CE2	2:L:101:MET:HG3	2.54	0.42
2:D:11:THR:HG23	5:D:311:HOH:O	2.20	0.42
1:E:17:GLU:HG2	1:E:77:SER:HB3	2.01	0.42
1:E:17:GLU:HG2	1:E:77:SER:CB	2.50	0.41
2:D:95:PHE:CE2	2:D:101:MET:HG3	2.56	0.41
1:E:2:VAL:HG11	1:E:129:TRP:CE3	2.53	0.41
2:J:69:CYS:SG	2:J:70:PRO:HD2	2.60	0.41
1:A:2:VAL:HG23	1:A:132:GLY:HA2	2.02	0.41
1:A:27:SER:OG	1:I:122:ARG:NH2	2.54	0.41
1:A:108:PRO:HD2	5:A:326:HOH:O	2.21	0.41
2:F:25:ARG:NE	5:F:211:HOH:O	2.28	0.41
2:J:62:ALA:HA	2:J:63:PRO:HD3	1.96	0.40
1:C:1:MET:HE3	1:C:1:MET:HB3	2.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	132/134 (98%)	127 (96%)	5 (4%)	0	100	100
1	C	132/134 (98%)	128 (97%)	4 (3%)	0	100	100
1	E	130/134 (97%)	126 (97%)	4 (3%)	0	100	100
1	G	131/134 (98%)	128 (98%)	3 (2%)	0	100	100
1	I	132/134 (98%)	128 (97%)	4 (3%)	0	100	100
1	K	131/134 (98%)	128 (98%)	3 (2%)	0	100	100
2	B	111/114 (97%)	110 (99%)	1 (1%)	0	100	100
2	D	105/114 (92%)	103 (98%)	1 (1%)	1 (1%)	12	32
2	F	110/114 (96%)	106 (96%)	4 (4%)	0	100	100
2	H	104/114 (91%)	100 (96%)	3 (3%)	1 (1%)	12	32

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	J	107/114 (94%)	104 (97%)	1 (1%)	2 (2%)	6	17
2	L	111/114 (97%)	110 (99%)	1 (1%)	0	100	100
All	All	1436/1488 (96%)	1398 (97%)	34 (2%)	4 (0%)	36	60

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	J	106	ASN
2	D	32	ALA
2	H	32	ALA
2	J	32	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/107 (100%)	104 (97%)	3 (3%)	38	68
1	C	107/107 (100%)	104 (97%)	3 (3%)	38	68
1	E	106/107 (99%)	104 (98%)	2 (2%)	50	77
1	G	107/107 (100%)	105 (98%)	2 (2%)	50	77
1	I	107/107 (100%)	105 (98%)	2 (2%)	50	77
1	K	107/107 (100%)	103 (96%)	4 (4%)	30	59
2	B	90/90 (100%)	88 (98%)	2 (2%)	45	74
2	D	86/90 (96%)	81 (94%)	5 (6%)	18	42
2	F	89/90 (99%)	85 (96%)	4 (4%)	24	52
2	H	86/90 (96%)	83 (96%)	3 (4%)	32	61
2	J	88/90 (98%)	85 (97%)	3 (3%)	32	62
2	L	90/90 (100%)	88 (98%)	2 (2%)	45	74
All	All	1170/1182 (99%)	1135 (97%)	35 (3%)	36	66

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	GLU
1	A	62	ARG
1	A	124	ARG
2	B	27	THR
2	B	98	THR
1	C	17	GLU
1	C	62	ARG
1	C	124	ARG
2	D	11	THR
2	D	13	ILE
2	D	55	GLU
2	D	87	ARG
2	D	98	THR
1	E	62	ARG
1	E	124	ARG
2	F	27	THR
2	F	71	MET
2	F	98	THR
2	F	105	LEU
1	G	17	GLU
1	G	70	GLN
2	H	27	THR
2	H	43	ARG
2	H	55	GLU
1	I	17	GLU
1	I	23	GLN
2	J	27	THR
2	J	43	ARG
2	J	98	THR
1	K	17	GLU
1	K	50	SER
1	K	62	ARG
1	K	120	GLN
2	L	89	VAL
2	L	98	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
1	A	20	ASN
1	A	23	GLN
1	A	73	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	85	ASN
1	C	16	GLN
1	C	20	ASN
1	C	96	GLN
2	D	85	ASN
2	D	106	ASN
1	E	16	GLN
1	E	20	ASN
2	F	85	ASN
1	G	16	GLN
1	G	20	ASN
2	H	85	ASN
1	I	16	GLN
1	I	20	ASN
1	I	73	GLN
1	I	131	ASN
2	J	85	ASN
1	K	16	GLN
1	K	20	ASN
1	K	70	GLN
1	K	73	GLN
2	L	3	HIS
2	L	85	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	D	201	-	5,5,5	0.14	0	5,5,5	0.32	0
4	GOL	J	501	-	5,5,5	0.21	0	5,5,5	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	201	-	-	2/4/4/4	-
4	GOL	J	501	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	J	501	GOL	O1-C1-C2-C3
4	J	501	GOL	O1-C1-C2-O2
4	D	201	GOL	C1-C2-C3-O3
4	D	201	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	201	GOL	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	134/134 (100%)	-0.03	1 (0%) 84 83	29, 42, 71, 93	0
1	C	134/134 (100%)	0.47	4 (2%) 52 49	36, 60, 104, 116	0
1	E	132/134 (98%)	0.75	7 (5%) 32 28	39, 68, 125, 144	0
1	G	133/134 (99%)	0.10	3 (2%) 61 58	32, 50, 90, 127	0
1	I	134/134 (100%)	-0.06	2 (1%) 72 70	30, 43, 69, 79	0
1	K	133/134 (99%)	0.91	14 (10%) 11 9	47, 84, 139, 155	0
2	B	113/114 (99%)	0.15	2 (1%) 67 65	25, 48, 80, 93	0
2	D	109/114 (95%)	0.45	7 (6%) 25 22	23, 47, 101, 121	0
2	F	112/114 (98%)	0.34	7 (6%) 26 23	28, 46, 89, 110	0
2	H	108/114 (94%)	0.27	7 (6%) 25 22	25, 50, 93, 105	0
2	J	111/114 (97%)	0.28	7 (6%) 26 23	27, 47, 89, 133	0
2	L	113/114 (99%)	0.16	2 (1%) 67 65	30, 48, 75, 81	0
All	All	1466/1488 (98%)	0.32	63 (4%) 40 36	23, 52, 106, 155	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	62	ALA	5.1
1	C	36	ALA	4.9
2	H	-1	GLY	4.8
1	E	1	MET	4.2
1	G	1	MET	4.1
1	K	-1	PRO	4.0
2	F	68	VAL	3.8
2	J	68	VAL	3.8
1	K	1	MET	3.7
2	F	65	PRO	3.5
2	H	67	LEU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	J	62	ALA	3.4
2	L	71	MET	3.4
2	F	66	SER	3.3
2	D	71	MET	3.1
2	J	69	CYS	3.1
2	J	63	PRO	3.0
2	J	65	PRO	3.0
1	E	2	VAL	3.0
2	H	69	CYS	3.0
1	K	132	GLY	3.0
1	C	132	GLY	2.8
1	E	3	THR	2.8
1	A	1	MET	2.8
2	F	71	MET	2.8
2	D	72	TYR	2.8
1	K	23	GLN	2.7
2	H	72	TYR	2.7
2	D	68	VAL	2.7
2	F	2	ALA	2.7
1	K	120	GLN	2.6
1	K	45	TYR	2.6
2	J	99	CYS	2.6
1	C	38	GLY	2.5
1	K	131	ASN	2.5
1	K	62	ARG	2.5
2	H	68	VAL	2.5
1	C	37	GLY	2.4
2	H	70	PRO	2.4
1	K	30	ARG	2.4
1	G	-1	PRO	2.4
1	K	2	VAL	2.4
1	K	126	ASN	2.4
1	K	119	ALA	2.4
2	D	67	LEU	2.3
2	F	67	LEU	2.3
2	D	73	PHE	2.3
1	E	19	ALA	2.2
1	E	61	GLY	2.2
1	E	24	ILE	2.2
2	H	1	SER	2.2
2	J	1	SER	2.2
1	I	112	CYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	K	130	ALA	2.1
2	D	99	CYS	2.1
1	I	-2	GLY	2.1
2	L	72	TYR	2.1
2	F	61	ALA	2.1
1	K	3	THR	2.1
2	B	71	MET	2.0
1	E	132	GLY	2.0
2	B	1	SER	2.0
1	G	70	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	J	501	6/6	0.75	0.21	69,71,73,73	0
4	GOL	D	201	6/6	0.87	0.13	47,53,63,66	0
3	ZN	K	201	1/1	0.99	0.04	58,58,58,58	0
3	ZN	C	201	1/1	0.99	0.04	43,43,43,43	0
3	ZN	I	201	1/1	0.99	0.03	36,36,36,36	0
3	ZN	E	201	1/1	1.00	0.02	50,50,50,50	0
3	ZN	G	201	1/1	1.00	0.06	41,41,41,41	0
3	ZN	A	201	1/1	1.00	0.02	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.