



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 11:17 PM UTC

PDB ID : 4JNY / pdb\_00004jny  
Title : Crystal structure of PutA86-630 mutant D370A complexed with L-Tetrahydr  
o-2-furoic acid  
Authors : Tanner, J.J.  
Deposited on : 2013-03-16  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	602	<div> <div>13%</div> <div>76%</div> <div>18%</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional protein PutA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	491	3732	2369	655	690	18	0	3	0

There are 19 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	370	ALA	ASP	engineered mutation	UNP P09546
A	670	SER	-	expression tag	UNP P09546
A	671	SER	-	expression tag	UNP P09546
A	672	SER	-	expression tag	UNP P09546
A	673	VAL	-	expression tag	UNP P09546
A	674	ASP	-	expression tag	UNP P09546
A	675	LYS	-	expression tag	UNP P09546
A	676	LEU	-	expression tag	UNP P09546
A	677	ALA	-	expression tag	UNP P09546
A	678	ALA	-	expression tag	UNP P09546
A	679	ALA	-	expression tag	UNP P09546
A	680	LEU	-	expression tag	UNP P09546
A	681	GLU	-	expression tag	UNP P09546
A	682	HIS	-	expression tag	UNP P09546
A	683	HIS	-	expression tag	UNP P09546
A	684	HIS	-	expression tag	UNP P09546
A	685	HIS	-	expression tag	UNP P09546
A	686	HIS	-	expression tag	UNP P09546
A	687	HIS	-	expression tag	UNP P09546

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is TETRAHYDROFURAN-2-CARBOXYLIC ACID (CCD ID: TFB) (formula: C<sub>5</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	5	3		

- Molecule 4 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: C<sub>10</sub>H<sub>22</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			16	10	6		
4	A	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	214	Total	O	0	0
			214	214		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Bifunctional protein PutA



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.33Å 142.32Å 146.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.83 – 1.90 39.83 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (39.83-1.90) 99.7 (39.83-1.90)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.43 (at 1.89Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, $R_{free}$	0.190 , 0.219 0.191 , 0.220	Depositor DCC
$R_{free}$ test set	3082 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.8	Xtriage
Anisotropy	0.623	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 48.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.003 for -h,-l,-k	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4033	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1PE, TFB, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/3809	0.74	2/5172 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	175	ILE	CA-C-N	6.45	125.68	118.97
1	A	175	ILE	C-N-CA	6.45	125.68	118.97

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3732	0	3651	15	0
2	A	53	0	31	1	0
3	A	8	0	7	0	0
4	A	26	0	34	0	0
5	A	214	0	0	0	0
All	All	4033	0	3723	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.



All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:ARG:NH2	1:A:348:GLU:OE1	2.40	0.54
1:A:334:HIS:HB2	1:A:349:LEU:HG	1.90	0.53
1:A:116:LEU:HD23	1:A:120:VAL:HG12	1.95	0.48
1:A:546:HIS:CE1	1:A:547:GLU:HG3	2.50	0.47
1:A:373:GLU:H	1:A:373:GLU:CD	2.23	0.46
1:A:146:GLY:O	1:A:149:GLN:N	2.48	0.45
1:A:106:ALA:O	1:A:110:LEU:HG	2.17	0.45
1:A:151:LEU:C	1:A:153:GLN:H	2.25	0.44
2:A:2001:FAD:H4'	2:A:2001:FAD:H1'1	1.83	0.44
1:A:175:ILE:HA	1:A:176:PRO:HD3	1.86	0.44
1:A:135:LEU:HD23	1:A:570:ILE:HD13	1.99	0.43
1:A:135:LEU:HD12	1:A:135:LEU:HA	1.83	0.43
1:A:583:ASP:HA	1:A:584:PRO:HD3	1.91	0.41
1:A:254:LEU:HD12	1:A:254:LEU:HA	1.94	0.41
1:A:269:LEU:C	1:A:271:ASN:H	2.29	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	486/602 (81%)	470 (97%)	12 (2%)	4 (1%)	<b>16</b> <b>8</b>

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	147	MET
1	A	152	LEU
1	A	156	SER
1	A	146	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	362/489 (74%)	355 (98%)	7 (2%)	50 47

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	148	VAL
1	A	254	LEU
1	A	271	ASN
1	A	349	LEU
1	A	406	TYR
1	A	532	LYS
1	A	553	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	298	GLN
1	A	514	HIS
1	A	568	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	A	2001	-	58,58,58	2.24	14 (24%)	85,89,89	1.80	23 (27%)
3	TFB	A	2002	-	8,8,8	1.55	2 (25%)	7,10,10	1.53	2 (28%)
4	1PE	A	2003	-	15,15,15	0.54	0	14,14,14	1.48	0
4	1PE	A	2004	-	9,9,15	0.52	0	8,8,14	1.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	2001	-	-	5/34/50/50	0/6/6/6
3	TFB	A	2002	-	-	0/4/11/11	0/1/1/1
4	1PE	A	2003	-	-	5/13/13/13	-
4	1PE	A	2004	-	-	5/7/7/13	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	FAD	PA-O3P	-6.97	1.52	1.59
2	A	2001	FAD	O4-C4	6.93	1.36	1.23
2	A	2001	FAD	O2-C2	5.71	1.35	1.24
2	A	2001	FAD	C6A-N6A	4.59	1.45	1.34
2	A	2001	FAD	C4X-N5	4.28	1.40	1.30
2	A	2001	FAD	P-O3P	4.12	1.63	1.59
2	A	2001	FAD	C2-N1	3.12	1.43	1.36
2	A	2001	FAD	O2'-C2'	-3.08	1.36	1.43
3	A	2002	TFB	CB-CA	-2.91	1.46	1.52
2	A	2001	FAD	PA-O5B	-2.68	1.48	1.59
2	A	2001	FAD	C10-N1	2.53	1.38	1.33

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	FAD	O4B-C4B	-2.50	1.39	1.45
2	A	2001	FAD	C5A-C4A	2.24	1.43	1.39
3	A	2002	TFB	OXT-C	2.22	1.28	1.22
2	A	2001	FAD	C4A-N9A	2.18	1.42	1.37
2	A	2001	FAD	PA-O2A	-2.11	1.45	1.55

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	FAD	N3A-C2A-N1A	-6.17	119.24	128.58
2	A	2001	FAD	C5A-C4A-N3A	-4.63	120.34	126.72
2	A	2001	FAD	N3A-C4A-N9A	3.90	133.80	127.17
2	A	2001	FAD	C5A-N7A-C8A	3.77	109.37	103.45
2	A	2001	FAD	C4-C4X-N5	3.76	123.40	118.21
2	A	2001	FAD	C2A-N3A-C4A	3.52	120.44	111.83
2	A	2001	FAD	O2P-P-O3P	-3.34	98.25	107.27
2	A	2001	FAD	N9A-C8A-N7A	-3.17	109.44	113.94
2	A	2001	FAD	C4-N3-C2	-3.04	120.24	125.64
2	A	2001	FAD	C4X-C4-N3	2.99	120.85	113.25
2	A	2001	FAD	C4A-C5A-N7A	-2.78	107.41	110.58
2	A	2001	FAD	O4-C4-C4X	-2.47	120.01	126.53
2	A	2001	FAD	O3P-P-O1P	2.41	117.95	110.70
2	A	2001	FAD	O5'-P-O1P	2.40	118.44	108.94
2	A	2001	FAD	C2A-N1A-C6A	2.38	122.64	118.73
3	A	2002	TFB	CB-CA-C	-2.32	109.26	113.00
2	A	2001	FAD	C4X-C10-N10	2.32	119.81	116.48
3	A	2002	TFB	OB-C-CA	2.30	118.70	112.71
2	A	2001	FAD	C9-C9A-N10	-2.26	118.81	121.85
2	A	2001	FAD	C4'-C3'-C2'	2.23	117.28	113.57
2	A	2001	FAD	C5X-C9A-N10	2.20	119.96	117.97
2	A	2001	FAD	C10-C4X-N5	-2.18	120.35	124.81
2	A	2001	FAD	C4A-N9A-C8A	2.06	107.90	105.74
2	A	2001	FAD	O2-C2-N1	-2.06	118.39	121.80
2	A	2001	FAD	O2P-P-O5'	-2.03	98.35	107.57

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2001	FAD	N10-C1'-C2'-O2'
2	A	2001	FAD	N10-C1'-C2'-C3'
2	A	2001	FAD	C1'-C2'-C3'-C4'

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
4	A	2004	1PE	OH4-C13-C23-OH3
4	A	2003	1PE	OH7-C16-C26-OH6
4	A	2004	1PE	OH6-C15-C25-OH5
4	A	2003	1PE	OH4-C13-C23-OH3
4	A	2003	1PE	OH6-C15-C25-OH5
4	A	2004	1PE	C24-C14-OH5-C25
4	A	2003	1PE	C14-C24-OH4-C13
4	A	2004	1PE	C15-C25-OH5-C14
4	A	2003	1PE	C12-C22-OH3-C23
2	A	2001	FAD	O2'-C2'-C3'-C4'
2	A	2001	FAD	O2'-C2'-C3'-O3'
4	A	2004	1PE	OH5-C14-C24-OH4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	491/602 (81%)	0.81	81 (16%) 4 4	14, 39, 88, 106	3 (0%)

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	236	ILE	7.6
1	A	153	GLN	7.2
1	A	157	LEU	6.7
1	A	183	ALA	6.5
1	A	152	LEU	6.3
1	A	205	PHE	5.7
1	A	179	ALA	5.6
1	A	151	LEU	5.0
1	A	228	LEU	5.0
1	A	209	ALA	4.9
1	A	215	PHE	4.9
1	A	206	VAL	4.9
1	A	175	ILE	4.8
1	A	227	SER	4.8
1	A	213	LEU	4.8
1	A	156	SER	4.7
1	A	184	LEU	4.7
1	A	204	LEU	4.5
1	A	155	PHE	4.4
1	A	165	LEU	4.4
1	A	235	ILE	4.3
1	A	149	GLN	4.3
1	A	182	ASP	4.2
1	A	146	GLY	4.2
1	A	147	MET	4.2
1	A	229	SER	4.2
1	A	87	PRO	4.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	145	ALA	3.9
1	A	158	SER	3.9
1	A	231	SER	3.9
1	A	210	THR	3.8
1	A	211	TRP	3.8
1	A	208	ALA	3.8
1	A	148	VAL	3.7
1	A	141	ALA	3.7
1	A	177	ASP	3.5
1	A	180	THR	3.5
1	A	150	GLY	3.5
1	A	342	TYR	3.5
1	A	226	ALA	3.4
1	A	173	LEU	3.3
1	A	154	GLU	3.2
1	A	138	GLN	3.1
1	A	160	GLN	3.1
1	A	242	PRO	3.1
1	A	137	ASN	3.1
1	A	159	SER	3.1
1	A	450	LEU	3.0
1	A	444	ARG	3.0
1	A	214	LEU	3.0
1	A	207	ASN	2.9
1	A	230	ARG	2.9
1	A	142	SER	2.9
1	A	163	VAL	2.8
1	A	161	GLU	2.8
1	A	181	ARG	2.8
1	A	358	LEU	2.8
1	A	109	MET	2.7
1	A	241	GLU	2.7
1	A	270	ALA	2.6
1	A	505	PRO	2.6
1	A	232	LEU	2.6
1	A	243	LEU	2.6
1	A	164	ALA	2.5
1	A	501	GLN	2.5
1	A	143	GLY	2.5
1	A	169	ALA	2.4
1	A	168	LEU	2.4
1	A	246	LYS	2.3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	343	ASP	2.3
1	A	609	ARG	2.3
1	A	250	MET	2.3
1	A	176	PRO	2.3
1	A	118	GLN	2.2
1	A	244	ILE	2.2
1	A	234	ARG	2.2
1	A	212	GLY	2.2
1	A	271	ASN	2.0
1	A	344	ARG	2.0
1	A	610	ASP	2.0
1	A	451	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	1PE	A	2003	16/16	0.89	0.15	56,67,71,75	0
4	1PE	A	2004	10/16	0.89	0.16	61,64,67,67	0
2	FAD	A	2001	53/53	0.97	0.06	22,26,28,30	0
3	TFB	A	2002	8/8	0.97	0.06	26,29,30,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.