



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:32 AM UTC

PDB ID : 4PFP / pdb\_00004pfp  
Title : Myosin VI motor domain in the Pi release state (with Pi) space group P21  
Authors : Isabet, T.; Benisty, H.; Llinas, P.; Sweeney, H.L.; Houdusse, A.  
Deposited on : 2014-04-30  
Resolution : 2.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	788	
1	C	788	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12526 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Unconventional myosin-VI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	760	Total	C	N	O	S	0	6	0
			5988	3822	1032	1105	29			
1	C	766	Total	C	N	O	S	0	4	0
			6051	3854	1035	1132	30			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is S-1,2-PROPANEDIOL (CCD ID: PGO) (formula:  $C_3H_8O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			5	3	2		

- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula:  $O_4P$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		
5	C	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	170	Total	O	0	0
			170	170		
6	C	241	Total	O	0	0
			241	241		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Unconventional myosin-VI



#### • Molecule 1: Unconventional myosin-VI



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.65Å 94.89Å 97.30Å 90.00° 99.55° 90.00°	Depositor
Resolution (Å)	23.39 – 2.32 23.39 – 2.32	Depositor EDS
% Data completeness (in resolution range)	99.8 (23.39-2.32) 99.7 (23.39-2.32)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.28 (at 2.32Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, $R_{free}$	0.189 , 0.229 0.199 , 0.239	Depositor DCC
$R_{free}$ test set	3902 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.2	Xtriage
Anisotropy	0.526	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 40.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12526	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.74 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.6209e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PGO, MG, PO4, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.94	4/6121 (0.1%)	1.32	23/8270 (0.3%)
1	C	0.86	1/6175 (0.0%)	1.28	12/8340 (0.1%)
All	All	0.90	5/12296 (0.0%)	1.30	35/16610 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	392	THR	C-N	-33.18	0.92	1.33
1	A	393	ARG	C-N	14.35	1.53	1.33
1	A	5	LYS	C-N	-11.31	1.19	1.33
1	A	25	GLY	C-N	6.26	1.41	1.33
1	C	669	ILE	CA-CB	5.13	1.60	1.54

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	5	LYS	O-C-N	-11.65	104.36	123.00
1	A	330	ASP	CA-CB-CG	9.41	122.01	112.60
1	A	393	ARG	CA-C-N	7.94	134.02	123.06
1	A	393	ARG	C-N-CA	7.94	134.02	123.06
1	A	518	ILE	N-CA-C	7.92	117.98	110.53
1	A	25	GLY	CA-C-N	-7.39	112.14	119.90

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	25	GLY	C-N-CA	-7.39	112.14	119.90
1	A	393	ARG	O-C-N	-7.38	115.52	123.42
1	A	450	TYR	CA-C-N	-6.60	112.45	122.81
1	A	450	TYR	C-N-CA	-6.60	112.45	122.81
1	C	173	GLY	CA-C-N	6.53	129.32	120.38
1	C	173	GLY	C-N-CA	6.53	129.32	120.38
1	C	396	LEU	CA-C-N	6.05	132.59	121.70
1	C	396	LEU	C-N-CA	6.05	132.59	121.70
1	A	90	VAL	N-CA-C	-5.84	97.20	109.34
1	C	513	ASP	CA-CB-CG	5.82	118.42	112.60
1	C	485	ASN	CA-C-N	5.82	128.07	120.28
1	C	485	ASN	C-N-CA	5.82	128.07	120.28
1	C	90	VAL	N-CA-C	-5.79	97.30	109.34
1	A	577	PHE	CA-CB-CG	5.70	119.50	113.80
1	A	731	PRO	CA-C-N	5.69	128.18	120.38
1	A	731	PRO	C-N-CA	5.69	128.18	120.38
1	C	229	LEU	N-CA-C	5.67	119.06	111.24
1	C	577	PHE	CA-CB-CG	5.57	119.37	113.80
1	A	703	GLY	CA-C-N	5.53	127.75	122.27
1	A	703	GLY	C-N-CA	5.53	127.75	122.27
1	A	464	GLU	CB-CG-CD	5.39	121.77	112.60
1	C	478	GLU	CA-C-N	5.38	127.50	120.28
1	C	478	GLU	C-N-CA	5.38	127.50	120.28
1	A	579	ILE	N-CA-CB	5.29	117.40	111.21
1	A	475	TYR	CA-C-N	5.16	127.15	120.44
1	A	475	TYR	C-N-CA	5.16	127.15	120.44
1	A	747	ILE	CA-C-N	5.12	127.65	120.28
1	A	747	ILE	C-N-CA	5.12	127.65	120.28
1	A	103	ILE	N-CA-C	-5.10	104.24	108.63

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	5	LYS	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5988	0	5795	43	0
1	C	6051	0	5862	25	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
3	A	27	0	12	0	0
3	C	27	0	12	0	0
4	A	5	0	8	0	0
5	A	10	0	0	0	0
5	C	5	0	0	0	0
6	A	170	0	0	2	0
6	C	241	0	0	3	0
All	All	12526	0	11689	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:GLY:O	1:A:358:THR:HG23	1.78	0.83
1:C:68:LEU:HD12	1:C:702:GLN:HG2	1.64	0.79
1:A:299:GLU:CD	6:A:901:HOH:O	2.26	0.79
1:A:356:GLY:C	1:A:358:THR:HG23	2.09	0.78
1:C:723:PRO:HD2	1:C:726:LEU:HD12	1.67	0.77
1:A:357:SER:HA	1:A:358:THR:C	2.10	0.75
1:C:457:ILE:HD13	1:C:480:LEU:HD12	1.69	0.72
1:A:357:SER:HA	1:A:358:THR:HG23	1.77	0.65
1:A:449:SER:C	1:A:450:TYR:HA	2.23	0.64
1:A:358:THR:OG1	1:A:359:SER:N	2.33	0.60
1:A:357:SER:CA	1:A:358:THR:HG23	2.33	0.58
1:C:592:GLN:HA	6:C:1072:HOH:O	2.04	0.57
1:A:177:ASP:O	1:A:181:ARG:HG3	2.04	0.57
1:A:357:SER:CA	1:A:358:THR:CG2	2.85	0.54
1:A:739:PHE:CE2	1:A:756:VAL:HG21	2.45	0.52
1:C:512:GLN:HA	1:C:515:ILE:HD12	1.91	0.52
1:A:73:LEU:HD22	1:A:698:LEU:HD21	1.92	0.52
1:A:464:GLU:HG2	1:A:465:HIS:CD2	2.45	0.51
1:C:515:ILE:HD11	6:C:905:HOH:O	2.10	0.51
1:A:357:SER:HA	1:A:358:THR:CG2	2.41	0.50
1:A:394:VAL:HG12	1:A:604:SER:HB3	1.92	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:739:PHE:CE2	1:C:756:VAL:HG21	2.47	0.50
1:A:489:LEU:HD21	1:A:697:VAL:HG12	1.94	0.49
1:C:127:VAL:HG11	1:C:160:ASN:OD1	2.12	0.49
1:A:87:TYR:HB3	1:A:94:LEU:HD11	1.95	0.48
1:C:716:ASN:O	1:C:719:LYS:HG2	2.13	0.48
1:A:551:HIS:HB2	1:A:557:LEU:HD22	1.95	0.48
1:A:393:ARG:HB2	1:A:411:LEU:HD21	1.96	0.48
1:A:449:SER:C	1:A:450:TYR:CA	2.86	0.48
1:A:357:SER:CA	1:A:358:THR:C	2.85	0.48
1:A:705:PHE:HB3	1:A:757:PHE:HB3	1.95	0.48
1:C:87:TYR:HB3	1:C:94:LEU:HD11	1.95	0.48
1:C:516:ASP:O	1:C:520:ALA:HB2	2.16	0.46
1:C:739:PHE:HE2	1:C:756:VAL:HG21	1.80	0.46
1:C:11:HIS:HD2	1:C:14:ASP:H	1.64	0.46
1:A:134:ALA:O	1:A:145:GLN:HG3	2.15	0.46
1:A:357:SER:C	1:A:358:THR:HG22	2.41	0.46
1:C:134:ALA:O	1:C:145:GLN:HG3	2.16	0.46
1:C:268:PRO:HB2	1:C:279:THR:HB	1.98	0.45
1:C:68:LEU:CD1	1:C:702:GLN:HG2	2.40	0.45
1:A:356:GLY:C	1:A:358:THR:CG2	2.86	0.45
1:C:166:ARG:O	1:C:170:GLU:HB2	2.18	0.44
1:A:166:ARG:O	1:A:170:GLU:HB2	2.18	0.44
1:A:517:LEU:O	1:A:525:ILE:HG13	2.17	0.44
1:A:268:PRO:HB2	1:A:279:THR:HB	2.00	0.44
1:A:520:ALA:O	1:A:524:GLY:HA3	2.18	0.43
1:A:521:ARG:O	1:A:523:VAL:N	2.48	0.43
1:C:647:PHE:CE2	1:C:651:LEU:HD11	2.53	0.43
1:A:723:PRO:HD2	1:A:726:LEU:HD12	2.00	0.43
1:A:771:LYS:HB3	1:A:776:HIS:CD2	2.54	0.43
1:C:705:PHE:HB3	1:C:757:PHE:HB3	1.99	0.43
1:C:233:SER:HA	1:C:425:LYS:HE2	2.00	0.43
1:A:357:SER:N	1:A:358:THR:HG23	2.32	0.43
1:A:490:LYS:O	1:A:494:GLU:HG3	2.20	0.42
1:A:742:LEU:HD22	1:A:783:ARG:HH11	1.83	0.42
1:C:742:LEU:HD22	1:C:783:ARG:HH11	1.84	0.42
1:A:229:LEU:HD11	1:A:647:PHE:CE2	2.54	0.42
1:A:742:LEU:HD12	1:A:769:ILE:HD11	2.01	0.42
1:A:181:ARG:HG2	1:A:327:ILE:HG22	2.02	0.42
1:C:519:GLU:HB2	1:C:648:LYS:HE3	2.01	0.42
1:C:29:LEU:HG	1:C:45:ILE:HG12	2.00	0.42
1:A:462:TYR:CD2	1:A:586:VAL:HG22	2.54	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:759:ARG:HD3	6:A:909:HOH:O	2.18	0.42
1:A:475:TYR:OH	1:A:648:LYS:HG3	2.21	0.41
1:A:233:SER:HA	1:A:425:LYS:HE2	2.02	0.41
1:A:739:PHE:HE2	1:A:756:VAL:HG21	1.84	0.41
1:C:719:LYS:HA	1:C:722:MET:HG2	2.01	0.41
1:C:479:LYS:HA	6:C:905:HOH:O	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	758/788 (96%)	745 (98%)	13 (2%)	0	100	100
1	C	764/788 (97%)	751 (98%)	13 (2%)	0	100	100
All	All	1522/1576 (97%)	1496 (98%)	26 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	630/696 (90%)	620 (98%)	10 (2%)	55	72
1	C	644/696 (92%)	628 (98%)	16 (2%)	42	59

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1274/1392 (92%)	1248 (98%)	26 (2%)	48 66

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	SER
1	A	330	ASP
1	A	357	SER
1	A	359	SER
1	A	462	TYR
1	A	464	GLU
1	A	523	VAL
1	A	639	SER
1	A	732	ARG
1	A	733	LEU
1	C	46[A]	ASN
1	C	46[B]	ASN
1	C	179	ASP
1	C	180	ASP
1	C	339	ARG
1	C	359	SER
1	C	377	GLU
1	C	395	MET
1	C	406	VAL
1	C	407	ILE
1	C	409	VAL
1	C	478	GLU
1	C	614	ASP
1	C	615	LYS
1	C	733	LEU
1	C	779	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	160	ASN
1	A	293	GLN
1	A	370	GLN
1	A	482	GLN
1	A	507	HIS
1	A	716	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	11	HIS
1	C	186	ASN
1	C	293	GLN
1	C	370	GLN
1	C	481	GLN
1	C	485	ASN
1	C	650	GLN
1	C	712	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ADP	C	801	2	28,29,29	0.38	0	43,45,45	0.47	0
5	PO4	A	804	2	4,4,4	2.09	3 (75%)	6,6,6	0.86	0
5	PO4	A	805	-	4,4,4	2.54	1 (25%)	6,6,6	0.87	0
4	PGO	A	803	-	4,4,4	0.88	0	4,4,4	0.63	0
3	ADP	A	802	2	28,29,29	0.47	0	43,45,45	0.81	2 (4%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	PO4	C	802	2	4,4,4	2.59	2 (50%)	6,6,6	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	C	801	2	-	3/16/32/32	0/3/3/3
4	PGO	A	803	-	-	0/2/2/2	-
3	ADP	A	802	2	-	2/16/32/32	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	802	PO4	P-O1	4.36	1.60	1.50
5	A	805	PO4	P-O1	4.25	1.60	1.50
5	A	804	PO4	P-O1	2.27	1.55	1.50
5	A	804	PO4	P-O2	2.13	1.60	1.54
5	A	804	PO4	P-O4	2.12	1.60	1.54
5	C	802	PO4	P-O4	2.05	1.60	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	ADP	O3A-PA-O1A	-2.46	103.30	110.70
3	A	802	ADP	O3'-C3'-C4'	2.13	117.21	111.08

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	802	ADP	PA-O3A-PB-O3B
3	C	801	ADP	PA-O3A-PB-O1B
3	A	802	ADP	PA-O3A-PB-O2B
3	C	801	ADP	PA-O3A-PB-O2B
3	C	801	ADP	PA-O3A-PB-O3B


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	3

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	449:SER	C	450:TYR	N	2.84
1	A	5:LYS	C	6:PRO	N	1.19
1	A	392:THR	C	393:ARG	N	0.92

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	760/788 (96%)	0.14	22 (2%) 53 56	23, 57, 86, 114	7 (0%)
1	C	766/788 (97%)	-0.08	17 (2%) 62 65	14, 48, 75, 110	4 (0%)
All	All	1526/1576 (96%)	0.03	39 (2%) 57 60	14, 53, 83, 114	11 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	178	ILE	5.7
1	C	480	LEU	4.8
1	A	393	ARG	4.2
1	C	177	ASP	3.8
1	A	177	ASP	3.7
1	C	509	VAL	3.4
1	A	567	ILE	3.3
1	A	549	GLN	3.3
1	C	179	ASP	3.1
1	C	38	GLY	3.0
1	C	459	GLY	2.9
1	A	310	LEU	2.9
1	A	610	CYS	2.9
1	A	789	ILE	2.8
1	A	406	VAL	2.8
1	C	507	HIS	2.6
1	A	522	LEU	2.6
1	A	545	SER	2.6
1	A	462	TYR	2.6
1	A	622	GLU	2.5
1	C	397	THR	2.5
1	A	555	PHE	2.5
1	C	723	PRO	2.4
1	C	508	TYR	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	447	THR	2.3
1	C	582	PHE	2.3
1	C	458	ALA	2.3
1	A	517	LEU	2.2
1	A	5	LYS	2.2
1	C	462	TYR	2.1
1	A	178	ILE	2.1
1	C	726	LEU	2.1
1	C	584	GLY	2.1
1	A	174	THR	2.1
1	A	639	SER	2.1
1	A	730	ASP	2.1
1	A	557	LEU	2.0
1	A	580	ARG	2.0
1	C	789	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PGO	A	803	5/5	0.83	0.15	68,69,71,71	0
5	PO4	A	805	5/5	0.85	0.20	79,80,84,86	0
5	PO4	A	804	5/5	0.88	0.16	18,29,33,36	5
3	ADP	C	801	27/27	0.99	0.04	25,29,36,37	0
2	MG	A	801	1/1	0.99	0.04	29,29,29,29	0
2	MG	C	803	1/1	0.99	0.05	31,31,31,31	0
3	ADP	A	802	27/27	0.99	0.05	24,34,47,49	0
5	PO4	C	802	5/5	0.99	0.07	32,36,38,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.