



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:36 AM UTC

PDB ID : 4RCA / pdb_00004rca
Title : Crystal structure of human PTPdelta and human Slitrk1 complex
Authors : Kim, H.M.; Park, B.S.; Kim, D.; Lee, S.G.
Deposited on : 2014-09-15
Resolution : 2.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	<div> <div>6%</div> <div> <div></div> <div>81%</div> <div>16%</div> <div>.</div> </div> </div>
2	B	251	<div> <div>10%</div> <div> <div></div> <div>68%</div> <div>24%</div> <div>.</div> <div>.</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Receptor-type tyrosine-protein phosphatase delta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	300	2301	1431	410	447	13	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	323	SER	-	expression tag	UNP P23468
A	324	ARG	-	expression tag	UNP P23468
A	325	LEU	-	expression tag	UNP P23468
A	326	VAL	-	expression tag	UNP P23468
A	327	PRO	-	expression tag	UNP P23468
A	328	ARG	-	expression tag	UNP P23468

- Molecule 2 is a protein called SLIT and NTRK-like protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	241	1937	1235	327	366	9	0	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	265	SER	-	expression tag	UNP Q96PX8
B	266	ARG	-	expression tag	UNP Q96PX8
B	267	LEU	-	expression tag	UNP Q96PX8
B	268	VAL	-	expression tag	UNP Q96PX8
B	269	PRO	-	expression tag	UNP Q96PX8
B	270	ARG	-	expression tag	UNP Q96PX8

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Receptor-type tyrosine-protein phosphatase delta



- Molecule 2: SLIT and NTRK-like protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.98Å 60.20Å 83.68Å 90.00° 102.19° 90.00°	Depositor
Resolution (Å)	38.62 – 2.99 38.62 – 2.99	Depositor EDS
% Data completeness (in resolution range)	98.2 (38.62-2.99) 98.1 (38.62-2.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496, REFMAC 5.7.0029	Depositor
R, R_{free}	0.263 , 0.269 (Not available) , 0.243	Depositor DCC
R_{free} test set	758 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	52.6	Xtriage
Anisotropy	0.596	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	4287	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2345	0.76	0/3189
2	B	0.30	0/1977	0.77	2/2684 (0.1%)
All	All	0.31	0/4322	0.76	2/5873 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	252	LEU	CA-C-N	5.43	128.97	120.82
2	B	252	LEU	C-N-CA	5.43	128.97	120.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2301	0	2287	29	0
2	B	1937	0	1911	36	0
3	A	28	0	26	4	0
3	B	14	0	13	1	0
4	B	5	0	0	0	0
5	B	2	0	0	0	0
All	All	4287	0	4237	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:ASN:HD21	3:A:401:NAG:C1	1.70	1.03
2:B:253:ASN:HD21	3:B:302:NAG:C1	1.74	1.00
1:A:254:ASN:HD21	3:A:402:NAG:C1	1.76	0.98
1:A:290:GLU:OE2	3:A:402:NAG:H83	1.73	0.89
2:B:194:TYR:HA	2:B:198:LEU:HB2	1.54	0.89
1:A:242:PRO:HB2	1:A:315:ALA:HB2	1.73	0.68
2:B:34:ILE:HG12	2:B:35:GLU:HA	1.74	0.68
2:B:66:HIS:O	2:B:92:ASN:ND2	2.35	0.60
2:B:83:ASN:HA	2:B:107:LEU:HD12	1.83	0.59
2:B:216:CYS:HB2	2:B:247:LEU:HD23	1.83	0.59
1:A:290:GLU:OE2	3:A:402:NAG:C8	2.48	0.59
1:A:59:LYS:HB2	1:A:64:VAL:HG22	1.86	0.57
1:A:58:ASN:HA	1:A:63:LYS:HA	1.87	0.56
1:A:277:THR:HG21	1:A:287:ASN:HD21	1.71	0.55
1:A:305:MET:HG2	1:A:310:VAL:HG13	1.89	0.55
2:B:145:ILE:O	2:B:170:PRO:HG2	2.07	0.55
2:B:34:ILE:HG12	2:B:35:GLU:HG3	1.89	0.54
1:A:38:GLY:HA2	1:A:86:PRO:HB2	1.89	0.54
2:B:258:GLN:N	2:B:258:GLN:OE1	2.41	0.54
1:A:258:VAL:HG22	1:A:286:ARG:HG3	1.91	0.51
2:B:230:ILE:HG12	2:B:231:PRO:HD2	1.93	0.51
2:B:217:THR:HG22	2:B:218:CYS:H	1.76	0.51
2:B:224:LYS:NZ	2:B:228:GLU:OE2	2.45	0.50
2:B:73:LEU:HD21	2:B:89:MET:HE1	1.93	0.50
2:B:247:LEU:HD22	2:B:259:ASP:OD1	2.11	0.50
2:B:122:ARG:HA	2:B:146:ASP:HB2	1.93	0.50
1:A:265:PRO:HB3	1:A:306:SER:HB3	1.94	0.49
2:B:88:HIS:HA	2:B:112:HIS:HB2	1.94	0.49
2:B:65:LEU:HD12	2:B:87:LEU:HD11	1.94	0.49
2:B:23:VAL:CG2	2:B:24:CYS:HA	2.43	0.49
1:A:284:ILE:HG12	1:A:284:ILE:O	2.13	0.48
2:B:194:TYR:CZ	2:B:199:GLU:HB3	2.47	0.48
1:A:139:ARG:HG2	1:A:140:THR:HG23	1.96	0.48
2:B:23:VAL:HG22	2:B:24:CYS:HA	1.96	0.48
1:A:202:GLN:HB3	1:A:225:VAL:HG23	1.95	0.48
1:A:87:LEU:HD13	1:A:115:VAL:HG22	1.95	0.47
2:B:134:TYR:HA	2:B:158:VAL:HG13	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:209:LEU:HB3	2:B:214:TRP:HZ2	1.80	0.46
2:B:169:LEU:HD12	2:B:192:LEU:HD11	1.96	0.46
1:A:26:PHE:CE2	1:A:109:VAL:HG23	2.51	0.45
2:B:110:ARG:HG3	2:B:134:TYR:CD1	2.51	0.45
1:A:284:ILE:HD13	1:A:284:ILE:H	1.81	0.45
1:A:174:ARG:NH1	1:A:195:GLU:O	2.48	0.44
2:B:189:LEU:H	2:B:213:PRO:HD2	1.82	0.44
2:B:103:LEU:HA	2:B:128:GLY:HA3	1.99	0.44
1:A:55:ILE:HG13	1:A:79:GLY:HA2	1.99	0.44
1:A:138:GLU:HB2	1:A:141:ARG:HD3	2.00	0.43
1:A:93:GLU:HA	1:A:94:ALA:HA	1.68	0.43
2:B:194:TYR:HB2	2:B:222:SER:HB3	1.99	0.43
1:A:87:LEU:HA	1:A:92:ASP:OD2	2.19	0.43
2:B:44:LYS:HA	2:B:67:GLY:O	2.19	0.42
2:B:66:HIS:CD2	2:B:91:ASN:HB2	2.55	0.42
1:A:66:ASN:H	1:A:69:PHE:HE1	1.67	0.42
1:A:26:PHE:CD2	1:A:107:ILE:HD12	2.55	0.42
1:A:158:THR:HG23	1:A:208:VAL:HB	2.02	0.42
2:B:27:LYS:O	2:B:27:LYS:HG3	2.19	0.42
2:B:106:GLN:O	2:B:131:ASP:HB2	2.20	0.41
2:B:247:LEU:HD11	2:B:250:LYS:HD2	2.01	0.41
1:A:42:SER:HA	1:A:82:LEU:O	2.20	0.41
1:A:92:ASP:O	1:A:96:TYR:OH	2.34	0.41
2:B:23:VAL:HG21	2:B:29:CYS:HB2	2.01	0.41
2:B:34:ILE:CG1	2:B:35:GLU:HA	2.47	0.41
1:A:254:ASN:HA	1:A:289:LEU:O	2.20	0.41
2:B:162:ASN:HB3	2:B:183:ASP:OD1	2.20	0.40
2:B:238:ARG:NH1	2:B:251:ASP:OD1	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	298/308 (97%)	289 (97%)	9 (3%)	0	100	100
2	B	239/251 (95%)	217 (91%)	21 (9%)	1 (0%)	30	65
All	All	537/559 (96%)	506 (94%)	30 (6%)	1 (0%)	43	76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	66	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/266 (97%)	253 (98%)	4 (2%)	55	79
2	B	219/228 (96%)	196 (90%)	23 (10%)	6	27
All	All	476/494 (96%)	449 (94%)	27 (6%)	18	52

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	156	GLU
1	A	157	ILE
1	A	281	ASP
1	A	284	ILE
2	B	31	CYS
2	B	34	ILE
2	B	35	GLU
2	B	57	THR
2	B	83	ASN
2	B	96	GLU
2	B	103	LEU
2	B	142	LEU
2	B	158	VAL
2	B	166	ILE
2	B	173	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	188	ARG
2	B	190	LYS
2	B	196	GLU
2	B	204	ILE
2	B	217	THR
2	B	221	LEU
2	B	230	ILE
2	B	232	LYS
2	B	238	ARG
2	B	250	LYS
2	B	258	GLN
2	B	261	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	ASN
1	A	103	ASN
1	A	133	GLN
1	A	193	GLN
1	A	254	ASN
1	A	299	ASN
2	B	66	HIS
2	B	76	ASN
2	B	83	ASN
2	B	91	ASN
2	B	95	HIS
2	B	200	GLN
2	B	253	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	402	1	14,14,15	0.64	0	17,19,21	1.33	1 (5%)
4	SO4	B	301	-	4,4,4	0.23	0	6,6,6	0.10	0
3	NAG	A	401	1	14,14,15	0.34	0	17,19,21	0.44	0
3	NAG	B	302	2	14,14,15	0.88	1 (7%)	17,19,21	1.81	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	402	1	-	4/6/23/26	0/1/1/1
3	NAG	A	401	1	-	4/6/23/26	0/1/1/1
3	NAG	B	302	2	-	4/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	NAG	O5-C1	-2.55	1.39	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	NAG	C1-O5-C5	6.63	121.07	112.19
3	A	402	NAG	C1-O5-C5	-4.10	106.69	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	NAG	C4-C3-C2	2.14	114.15	111.02

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	302	NAG	C4-C5-C6-O6
3	A	401	NAG	O5-C5-C6-O6
3	A	401	NAG	C4-C5-C6-O6
3	A	401	NAG	C8-C7-N2-C2
3	A	401	NAG	O7-C7-N2-C2
3	A	402	NAG	C8-C7-N2-C2
3	A	402	NAG	O7-C7-N2-C2
3	B	302	NAG	C8-C7-N2-C2
3	B	302	NAG	O7-C7-N2-C2
3	B	302	NAG	O5-C5-C6-O6
3	A	402	NAG	C4-C5-C6-O6
3	A	402	NAG	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	NAG	3	0
3	A	401	NAG	1	0
3	B	302	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	300/308 (97%)	0.88	20 (6%) 24 12	37, 74, 107, 138	0
2	B	241/251 (96%)	0.95	26 (10%) 11 6	48, 71, 106, 115	0
All	All	541/559 (96%)	0.91	46 (8%) 16 8	37, 72, 106, 138	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	259	ASP	7.7
2	B	141	LEU	6.6
1	A	35	GLY	4.3
2	B	107	LEU	4.0
1	A	172	ASN	3.9
2	B	247	LEU	3.8
1	A	170	ASN	3.4
1	A	171	ASN	3.4
2	B	91	ASN	3.4
1	A	96	TYR	3.2
1	A	92	ASP	3.1
1	A	207	CYS	3.1
2	B	249	GLY	3.0
2	B	65	LEU	3.0
2	B	34	ILE	2.9
1	A	183	ILE	2.8
1	A	284	ILE	2.8
2	B	69	SER	2.8
2	B	71	THR	2.8
2	B	66	HIS	2.7
1	A	54	LYS	2.7
2	B	23	VAL	2.7
2	B	188	ARG	2.6
2	B	89	MET	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	262	PRO	2.5
1	A	33	GLN	2.4
1	A	66	ASN	2.4
2	B	172	ASN	2.3
1	A	120	GLN	2.3
2	B	28	ILE	2.3
2	B	215	ASP	2.3
2	B	47	PHE	2.2
2	B	78	PHE	2.2
2	B	93	GLY	2.2
1	A	53	PRO	2.1
1	A	104	VAL	2.1
2	B	108	VAL	2.1
2	B	204	ILE	2.1
1	A	64	VAL	2.1
1	A	61	GLY	2.1
1	A	89	THR	2.1
2	B	257	GLU	2.1
2	B	137	ALA	2.0
1	A	173	GLY	2.0
2	B	203	GLY	2.0
1	A	288	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	B	302	14/15	0.44	0.23	82,93,94,95	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	A	402	14/15	0.49	0.17	109,113,114,114	0
3	NAG	A	401	14/15	0.69	0.14	97,100,103,105	0
4	SO4	B	301	5/5	0.83	0.21	111,112,113,113	0

6.5 Other polymers [i](#)

There are no such residues in this entry.