



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:32 AM UTC

PDB ID : 5CEH / pdb\_00005ceh  
Title : Structure of histone lysine demethylase KDM5A in complex with selective inhibitor  
Authors : Kiefer, J.R.; Vinogradova, M.  
Deposited on : 2015-07-06  
Resolution : 3.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	NOT EXECUTED
Xtriage (Phenix)	:	2.0
EDS	:	NOT EXECUTED
Buster-report	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2452 (3.18-3.10)
Ramachandran outliers	187476	2324 (3.18-3.10)
Sidechain outliers	187428	2324 (3.18-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	790	 59% 12% • 27%
2	B	10	 70% 30%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific demethylase 5A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	577	Total	C	N	O	S	0	0	0
			4591	2952	755	846	38			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	11	SER	-	expression tag	UNP P29375
A	798	GLY	-	expression tag	UNP P29375
A	799	ASN	-	expression tag	UNP P29375
A	800	SER	-	expression tag	UNP P29375

- Molecule 2 is a protein called Unknown Peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	10	Total	C	N	O	0	0	0
			50	30	10	10			


- Molecule 3 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ni	0	0
			1	1		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		

- Molecule 5 is 7-oxo-5-phenyl-6-(propan-2-yl)-4,7-dihydropyrazolo[1,5-a]pyrimidine-3-carbonitrile (CCD ID: 50P) (formula: C<sub>16</sub>H<sub>14</sub>N<sub>4</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			21	16	4	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	7	Total	O	0	0
			7	7		

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Lysine-specific demethylase 5A



- Molecule 2: Unknown Peptide



## 4 Data and refinement statistics

EDS was not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	159.31Å 159.31Å 92.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.83 – 3.14	Depositor
% Data completeness (in resolution range)	96.0 (39.83-3.14)	Depositor
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 3.12Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.221 , 0.238	Depositor
Wilson B-factor (Å <sup>2</sup> )	101.8	Xtriage
Anisotropy	0.375	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
Total number of atoms	4672	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, ZN, 50P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	1/4719 (0.0%)	0.86	9/6415 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	459	ILE	CA-CB	5.29	1.61	1.54

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	489	SER	CB-CA-C	-7.30	106.25	115.89
1	A	177	SER	N-CA-C	7.28	118.77	108.00
1	A	177	SER	CB-CA-C	-6.15	107.73	116.34
1	A	430	LYS	N-CA-C	-6.11	101.24	110.28
1	A	64	GLU	N-CA-C	-5.76	103.09	110.53
1	A	428	ARG	NE-CZ-NH2	-5.73	114.04	119.20
1	A	714	LYS	CB-CA-C	-5.41	109.35	115.79
1	A	720	TYR	CA-C-N	5.36	125.44	119.87
1	A	720	TYR	C-N-CA	5.36	125.44	119.87

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	778	ASN	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4591	0	4388	43	1
2	B	50	0	13	2	0
3	A	1	0	0	0	0
4	A	2	0	0	0	0
5	A	21	0	0	1	0
6	A	7	0	0	0	0
All	All	4672	0	4401	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:457:ALA:O	1:A:459:ILE:N	2.15	0.79
1:A:391:GLU:OE2	1:A:554:ARG:NH1	2.23	0.70
1:A:707:CYS:SG	1:A:709:CYS:HB2	2.40	0.61
1:A:426:ASP:O	1:A:428:ARG:N	2.34	0.61
1:A:55:LYS:O	1:A:58:GLN:NE2	2.36	0.59
1:A:714:LYS:CB	1:A:715:CYS:HA	2.34	0.57
1:A:714:LYS:CB	1:A:715:CYS:CA	2.83	0.56
1:A:92:ASP:OD2	1:A:96:LYS:NZ	2.31	0.55
1:A:679:CYS:SG	1:A:702:HIS:CE1	3.01	0.54
1:A:677:SER:N	1:A:697:LEU:O	2.41	0.53
1:A:724:ASP:OD1	1:A:725:LEU:N	2.46	0.49
1:A:676:CYS:HB3	1:A:679:CYS:HB2	1.96	0.48
1:A:542:PRO:HB3	1:A:552:VAL:HG11	1.94	0.48
1:A:369:MET:HG3	1:A:370:ALA:N	2.27	0.47
1:A:679:CYS:HB3	1:A:681:THR:N	2.29	0.47
1:A:599:VAL:O	1:A:603:ARG:N	2.46	0.47
1:A:679:CYS:HB3	1:A:680:ARG:CA	2.44	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:TYR:O	1:A:166:LEU:N	2.47	0.47
1:A:679:CYS:CB	1:A:680:ARG:HA	2.44	0.47
2:B:910:UNK:O	2:B:913:UNK:N	2.48	0.47
1:A:589:ALA:HB2	1:A:631:LEU:HD12	1.97	0.46
1:A:646:GLU:OE2	1:A:686:SER:OG	2.20	0.46
2:B:913:UNK:O	2:B:914:UNK:C	2.65	0.45
1:A:114:ILE:HD13	1:A:114:ILE:H	1.81	0.45
1:A:660:MET:HB3	1:A:714:LYS:CB	2.46	0.45
1:A:485:GLU:HB2	1:A:488:TRP:O	2.17	0.44
1:A:679:CYS:HB3	1:A:680:ARG:HA	2.00	0.43
1:A:390:VAL:HB	1:A:576:GLN:OE1	2.19	0.43
1:A:24:PRO:HB3	1:A:35:PHE:CZ	2.54	0.43
1:A:375:SER:O	1:A:379:ASN:N	2.52	0.42
1:A:656:MET:O	1:A:711:MET:CB	2.67	0.42
1:A:458:HIS:ND1	1:A:458:HIS:C	2.77	0.42
1:A:676:CYS:SG	1:A:679:CYS:HB2	2.60	0.42
1:A:76:ARG:O	1:A:80:LEU:N	2.54	0.41
1:A:374:LYS:HE2	1:A:385:VAL:CG1	2.50	0.41
1:A:24:PRO:HB3	1:A:35:PHE:CE1	2.56	0.41
1:A:721:PRO:O	1:A:722:LEU:HB2	2.21	0.41
1:A:167:TYR:N	1:A:168:PRO:CD	2.83	0.41
1:A:480:PHE:HB3	5:A:904:50P:C5	2.51	0.41
1:A:719:ARG:HG3	1:A:720:TYR:CE1	2.56	0.41
1:A:459:ILE:HG13	1:A:459:ILE:O	2.20	0.41
1:A:425:LYS:NZ	1:A:436:GLU:OE1	2.44	0.40
1:A:463:ILE:H	1:A:463:ILE:HG13	1.68	0.40
1:A:777:GLU:CB	1:A:781:PHE:HB2	2.52	0.40
1:A:40:ARG:N	1:A:41:PRO:CD	2.84	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:428:ARG:CZ	1:A:428:ARG:NH2[6_977]	1.35	0.85

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/790 (41%)	302 (93%)	20 (6%)	3 (1%)	14	41

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	458	HIS
1	A	457	ALA
1	A	711	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	498/710 (70%)	450 (90%)	48 (10%)	8	27

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	GLU
1	A	30	THR
1	A	38	ARG
1	A	39	ILE
1	A	55	LYS
1	A	65	VAL
1	A	100	LEU
1	A	105	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	107	ILE
1	A	109	VAL
1	A	113	LYS
1	A	114	ILE
1	A	117	LEU
1	A	122	LYS
1	A	137	LYS
1	A	138	LYS
1	A	166	LEU
1	A	182	GLN
1	A	363	LEU
1	A	369	MET
1	A	384	MET
1	A	385	VAL
1	A	401	ILE
1	A	448	MET
1	A	450	VAL
1	A	455	VAL
1	A	459	ILE
1	A	463	ILE
1	A	466	MET
1	A	475	MET
1	A	484	ILE
1	A	501	LYS
1	A	512	GLU
1	A	555	THR
1	A	576	GLN
1	A	614	GLU
1	A	616	LEU
1	A	641	LEU
1	A	655	GLN
1	A	669	VAL
1	A	671	ASP
1	A	675	GLN
1	A	679	CYS
1	A	690	CYS
1	A	695	GLU
1	A	698	VAL
1	A	720	TYR
1	A	780	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	GLN
1	A	613	HIS
1	A	693	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

Mogul was not executed - this section is therefore empty.

## 5.5 Carbohydrates [i](#)

Mogul was not executed - this section is therefore empty.

## 5.6 Ligand geometry [i](#)

Mogul was not executed - this section is therefore empty.

## 5.7 Other polymers [i](#)

Mogul was not executed - this section is therefore empty.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.




### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 50P A 904:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [\(i\)](#)

EDS was not executed - this section is therefore empty.