



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:16 PM UTC

PDB ID : 5FSE / pdb_00005fse
Title : 2.07 Å resolution 1,4-Benzoquinone inhibited *Sporosarcina pasteurii* urease
Authors : Mazzei, L.; Cianci, M.; Musiani, F.; Ciurli, S.
Deposited on : 2016-01-04
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	100	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 89% 10% . </div> </div>
2	B	126	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green, yellow, orange, red);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 89% 8% . </div> </div>
3	C	570	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 90% 8% . </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	C	1573	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 6799 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UREASE SUBUNIT GAMMA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	6	0
			824	519	140	158	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	ALA	LEU	conflict	UNP P41022
A	22	LYS	ARG	conflict	UNP P41022

- Molecule 2 is a protein called UREASE SUBUNIT BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	122	Total	C	N	O	S	0	2	0
			969	599	174	195	1			

- Molecule 3 is a protein called UREASE SUBUNIT ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	570	Total	C	N	O	S	0	16	0
			4438	2785	763	864	26			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	19	GLN	ARG	conflict	UNP P41020
C	28	TRP	GLY	conflict	UNP P41020
C	29	ILE	-	insertion	UNP P41020
C	36	THR	TYR	conflict	UNP P41020
C	37	THR	TYR	conflict	UNP P41020
C	38	TYR	LEU	conflict	UNP P41020
C	42	ALA	VAL	conflict	UNP P41020
C	263	LEU	VAL	conflict	UNP P41020

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	403	ALA	LEU	conflict	UNP P41020
C	420	ILE	MET	conflict	UNP P41020

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	2	Total	Ni	0	0
			2	2		

- Molecule 7 is HYDROXIDE ION (CCD ID: OH) (formula: HO).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	1	Total	O	0	0
			1	1		

- Molecule 8 is benzene-1,4-diol (CCD ID: HQE) (formula: C₆H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	C	1	Total	C	O	0	0
			8	6	2		
8	C	1	Total	C	O	0	0
			8	6	2		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	70	Total 70	O 70	0	0
9	B	93	Total 93	O 93	0	0
9	C	301	Total 301	O 301	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: UREASE SUBUNIT GAMMA



- Molecule 2: UREASE SUBUNIT BETA



- Molecule 3: UREASE SUBUNIT ALPHA



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.83Å 131.83Å 188.58Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	114.17 – 2.07 114.17 – 2.07	Depositor EDS
% Data completeness (in resolution range)	98.4 (114.17-2.07) 98.4 (114.17-2.07)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.07Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.145 , 0.192 0.158 , 0.198	Depositor DCC
R_{free} test set	2944 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	27.3	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6799	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NI, SO4, OH, EDO, HQE, CXM, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.19	1/831 (0.1%)	1.06	1/1117 (0.1%)
2	B	1.12	2/981 (0.2%)	1.04	0/1320
3	C	1.11	7/4519 (0.2%)	1.04	4/6121 (0.1%)
All	All	1.12	10/6331 (0.2%)	1.05	5/8558 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	95	VAL	C-O	-6.67	1.17	1.24
3	C	23	ALA	C-O	-5.86	1.18	1.24
1	A	97	ASN	C-O	-5.65	1.21	1.23
3	C	189	GLY	C-O	-5.64	1.20	1.24
3	C	102	ASN	C-O	-5.51	1.20	1.25
3	C	281	GLY	C-O	5.49	1.28	1.23
3	C	23	ALA	CA-C	-5.33	1.48	1.53
3	C	191	PRO	CA-C	-5.17	1.48	1.53
2	B	91	GLY	C-O	-5.12	1.19	1.24
2	B	27	LYS	C-O	-5.09	1.17	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	520	CYS	CB-CA-C	-6.99	97.49	109.65
3	C	234	ARG	CG-CD-NE	-5.78	99.28	112.00
1	A	13	ILE	CB-CA-C	-5.51	104.80	112.02
3	C	121	VAL	N-CA-C	5.33	115.56	108.11
3	C	60	TYR	N-CA-C	5.12	118.29	110.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	366[A]	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	824	0	840	5	0
2	B	969	0	949	4	0
3	C	4438	0	4407	36	0
4	A	4	0	6	0	0
4	B	4	0	6	0	0
4	C	32	0	48	7	0
5	A	5	0	0	1	0
5	B	5	0	0	0	0
5	C	35	0	0	1	0
6	C	2	0	0	0	0
7	C	1	0	0	0	0
8	C	16	0	8	1	0
9	A	70	0	0	0	0
9	B	93	0	0	1	0
9	C	301	0	0	3	1
All	All	6799	0	6264	43	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:64[B]:GLU:HG2	4:C:1573:EDO:H11	1.68	0.75
3:C:64[B]:GLU:HG2	4:C:1573:EDO:C1	2.18	0.74
3:C:507[A]:LYS:HD3	9:C:2250:HOH:O	1.93	0.69
3:C:93:TYR:OH	4:C:1578:EDO:H12	1.92	0.69
3:C:452[B]:LYS:HE2	3:C:491:ASN:OD1	1.94	0.67
1:A:79[B]:ASP:OD2	1:A:96[B]:HIS:ND1	2.28	0.67
3:C:287[A]:ILE:O	3:C:287[A]:ILE:HD13	1.99	0.61
3:C:452[B]:LYS:NZ	9:C:2230:HOH:O	2.33	0.61
1:A:71:GLU:N	5:A:1102:SO4:O3	2.28	0.58
3:C:323:HIS:NE2	5:C:1579:SO4:O4	2.37	0.58
3:C:301:THR:CG2	3:C:363:ASP:HB2	2.36	0.55
3:C:191:PRO:O	3:C:452[B]:LYS:HE3	2.09	0.53
3:C:64[B]:GLU:HG2	4:C:1573:EDO:H12	1.88	0.52
3:C:220:KCX:CX	3:C:222:HIS:HD2	2.24	0.49
8:C:1584:HQE:H5	9:C:2110:HOH:O	2.12	0.49
3:C:303:PRO:HG3	3:C:368:GLY:HA2	1.93	0.49
3:C:64[B]:GLU:CG	4:C:1573:EDO:H11	2.40	0.49
3:C:287[A]:ILE:HD13	3:C:287[A]:ILE:C	2.38	0.48
3:C:513:ARG:HD3	4:C:1578:EDO:H11	1.97	0.47
3:C:301:THR:HG21	3:C:363:ASP:HB2	1.95	0.47
3:C:158:PHE:CE2	3:C:418[A]:GLN:CG	2.98	0.47
3:C:139:HIS:CD2	3:C:366[B]:ALA:HB2	2.51	0.46
3:C:70:LEU:HD11	3:C:86:ASP:HB3	1.98	0.45
3:C:64[A]:GLU:O	3:C:65:ASN:HB2	2.16	0.45
3:C:138:VAL:O	3:C:159:GLY:HA3	2.15	0.45
3:C:513:ARG:HD3	4:C:1578:EDO:C1	2.46	0.45
2:B:111[B]:GLU:HG2	9:B:2086:HOH:O	2.17	0.43
3:C:282:GLY:O	3:C:283:HIS:C	2.61	0.43
3:C:362:THR:O	3:C:368:GLY:HA3	2.18	0.43
3:C:178:TRP:O	3:C:182:LYS:HG2	2.19	0.42
3:C:302:ASN:N	3:C:303:PRO:CD	2.82	0.42
3:C:320:MET:HA	3:C:325:LEU:HD12	2.02	0.42
3:C:393[A]:GLU:CD	3:C:393[A]:GLU:H	2.27	0.42
1:A:96[B]:HIS:O	1:A:97:ASN:C	2.63	0.41
3:C:274:PHE:O	3:C:275:HIS:C	2.63	0.41
3:C:320:MET:HA	3:C:325:LEU:HB2	2.03	0.41
1:A:73:VAL:N	1:A:74:PRO:CD	2.84	0.41
1:A:12[A]:GLN:HE22	3:C:475[A]:MET:HE1	1.87	0.40
2:B:10:GLY:HA2	3:C:21:ARG:O	2.21	0.40
3:C:220:KCX:CX	3:C:222:HIS:CD2	3.04	0.40
3:C:461:ILE:HD11	3:C:469:PRO:HB3	2.04	0.40
2:B:95:GLU:O	3:C:104:ASP:HB3	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:118:LYS:HB2	2:B:125:VAL:HG11	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:C:2051:HOH:O	9:C:2051:HOH:O[11_555]	1.92	0.28

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	104/100 (104%)	104 (100%)	0	0	100	100
2	B	122/126 (97%)	117 (96%)	4 (3%)	1 (1%)	16	8
3	C	583/570 (102%)	562 (96%)	19 (3%)	2 (0%)	36	30
All	All	809/796 (102%)	783 (97%)	23 (3%)	3 (0%)	30	23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	367	MET
2	B	99	ILE
3	C	275	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/83 (107%)	86 (97%)	3 (3%)	32	27
2	B	103/105 (98%)	101 (98%)	2 (2%)	50	49
3	C	473/458 (103%)	459 (97%)	14 (3%)	36	32
All	All	665/646 (103%)	646 (97%)	19 (3%)	45	33

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81[A]	GLN
1	A	81[B]	GLN
1	A	87	PRO
2	B	18	GLU
2	B	110	LYS
3	C	1	MET
3	C	249	HIS
3	C	287[A]	ILE
3	C	287[B]	ILE
3	C	305	ARG
3	C	326	LYS
3	C	393[A]	GLU
3	C	393[B]	GLU
3	C	395	LYS
3	C	452[A]	LYS
3	C	452[B]	LYS
3	C	507[A]	LYS
3	C	507[B]	LYS
3	C	511	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	179	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	KCX	C	220	3,6	10,11,12	2.13	1 (10%)	6,12,14	1.91	1 (16%)
1	CXM	A	1	1	9,10,11	2.21	2 (22%)	9,11,13	1.33	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KCX	C	220	3,6	-	0/9/10/12	-
1	CXM	A	1	1	-	2/9/10/12	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	220	KCX	OQ1-CX	6.17	1.33	1.21
1	A	1	CXM	ON1-CN	5.95	1.32	1.21
1	A	1	CXM	CN-N	2.22	1.39	1.35

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	220	KCX	OQ1-CX-NZ	-4.00	118.85	124.92
1	A	1	CXM	ON1-CN-N	-2.26	121.16	124.86

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1	CXM	CB-CA-N-CN
1	A	1	CXM	C-CA-N-CN

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	220	KCX	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 2 are monoatomic and 1 is modelled with single atom - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	SO4	C	1585	-	4,4,4	0.67	0	6,6,6	0.41	0
4	EDO	C	1571	-	3,3,3	0.34	0	2,2,2	0.52	0
5	SO4	C	1588	-	4,4,4	0.50	0	6,6,6	0.21	0
5	SO4	C	1579	-	4,4,4	0.86	0	6,6,6	0.56	0
5	SO4	C	1582	-	4,4,4	0.62	0	6,6,6	0.87	0
4	EDO	C	1573	-	3,3,3	0.30	0	2,2,2	0.55	0
5	SO4	A	1102	-	4,4,4	0.69	0	6,6,6	0.57	0
5	SO4	B	1128	-	4,4,4	0.50	0	6,6,6	0.39	0
4	EDO	C	1578	-	3,3,3	0.88	0	2,2,2	0.14	0
4	EDO	C	1577	-	3,3,3	0.46	0	2,2,2	0.65	0
4	EDO	B	1127	-	3,3,3	0.88	0	2,2,2	0.24	0
5	SO4	C	1586	-	4,4,4	0.49	0	6,6,6	0.47	0
4	EDO	C	1575	-	3,3,3	0.29	0	2,2,2	0.47	0
8	HQE	C	1584	3	8,8,8	2.50	1 (12%)	10,10,10	1.35	1 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	C	1587	-	4,4,4	0.60	0	6,6,6	0.69	0
4	EDO	C	1572	-	3,3,3	0.35	0	2,2,2	0.62	0
4	EDO	A	1101	-	3,3,3	0.33	0	2,2,2	0.43	0
8	HQE	C	1583	3	8,8,8	2.70	3 (37%)	10,10,10	1.26	1 (10%)
5	SO4	C	1581	-	4,4,4	0.56	0	6,6,6	0.59	0
4	EDO	C	1576	-	3,3,3	0.40	0	2,2,2	0.74	0
4	EDO	C	1574	-	3,3,3	0.37	0	2,2,2	1.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	1571	-	-	0/1/1/1	-
4	EDO	C	1572	-	-	1/1/1/1	-
4	EDO	B	1127	-	-	1/1/1/1	-
4	EDO	A	1101	-	-	0/1/1/1	-
4	EDO	C	1573	-	-	0/1/1/1	-
4	EDO	C	1575	-	-	0/1/1/1	-
8	HQE	C	1583	3	-	-	0/1/1/1
4	EDO	C	1578	-	-	0/1/1/1	-
8	HQE	C	1584	3	-	-	0/1/1/1
4	EDO	C	1576	-	-	1/1/1/1	-
4	EDO	C	1574	-	-	1/1/1/1	-
4	EDO	C	1577	-	-	1/1/1/1	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	C	1583	HQE	C5-C6	6.69	1.51	1.39
8	C	1584	HQE	C2-C3	6.33	1.50	1.39
8	C	1583	HQE	C2-C3	2.36	1.43	1.39
8	C	1583	HQE	C4-C3	2.20	1.43	1.39

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	C	1584	HQE	C2-C1-C6	2.37	122.38	119.88
8	C	1583	HQE	C4-C5-C6	-2.20	117.55	119.88

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	1572	EDO	O1-C1-C2-O2
4	C	1574	EDO	O1-C1-C2-O2
4	B	1127	EDO	O1-C1-C2-O2
4	C	1577	EDO	O1-C1-C2-O2
4	C	1576	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	1579	SO4	1	0
4	C	1573	EDO	4	0
5	A	1102	SO4	1	0
4	C	1578	EDO	3	0
8	C	1584	HQE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	99/100 (99%)	-0.58	1 (1%) 79 82	14, 26, 36, 61	6 (6%)
2	B	122/126 (96%)	-0.42	0 100 100	18, 29, 45, 81	2 (1%)
3	C	569/570 (99%)	-0.52	5 (0%) 81 83	10, 26, 49, 78	17 (2%)
All	All	790/796 (99%)	-0.52	6 (0%) 82 84	10, 26, 47, 81	25 (3%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	100	SER	2.4
3	C	366[A]	ALA	2.4
3	C	320	MET	2.4
3	C	327	GLN	2.3
3	C	326	LYS	2.2
3	C	550	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	KCX	C	220	12/13	0.97	0.06	21,23,27,28	0
1	CXM	A	1	11/12	0.98	0.06	26,31,36,41	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	C	1582	5/5	0.63	0.16	73,77,78,81	0
5	SO4	C	1581	5/5	0.73	0.11	66,94,101,106	0
5	SO4	A	1102	5/5	0.74	0.16	85,91,94,107	0
5	SO4	C	1579	5/5	0.79	0.14	63,68,79,95	0
5	SO4	B	1128	5/5	0.81	0.10	71,79,82,85	0
5	SO4	C	1586	5/5	0.82	0.11	85,101,112,118	0
5	SO4	C	1588	5/5	0.84	0.09	83,84,93,108	0
4	EDO	C	1578	4/4	0.85	0.16	43,43,43,49	0
4	EDO	C	1577	4/4	0.85	0.16	63,63,65,65	0
8	HQE	C	1584	8/8	0.85	0.14	54,67,71,75	0
4	EDO	C	1573	4/4	0.87	0.14	46,48,48,49	0
4	EDO	C	1574	4/4	0.88	0.13	41,42,42,48	0
5	SO4	C	1587	5/5	0.88	0.13	71,86,93,94	0
4	EDO	C	1572	4/4	0.90	0.15	54,56,58,63	0
5	SO4	C	1585	5/5	0.90	0.08	54,55,68,75	0
4	EDO	B	1127	4/4	0.91	0.12	43,48,49,57	0
4	EDO	C	1576	4/4	0.93	0.11	42,48,50,52	0
4	EDO	C	1571	4/4	0.94	0.09	38,43,44,49	0
8	HQE	C	1583	8/8	0.95	0.07	38,40,43,44	0
4	EDO	C	1575	4/4	0.97	0.07	41,41,43,43	0
4	EDO	A	1101	4/4	0.98	0.05	30,33,33,34	0
7	OH	C	1580	1/1	0.99	0.05	31,31,31,31	0
6	NI	C	601	1/1	1.00	0.02	29,29,29,29	0
6	NI	C	600	1/1	1.00	0.02	35,35,35,35	0

6.5 Other polymers ⓘ

There are no such residues in this entry.