



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:25 AM UTC

PDB ID : 5L7I / pdb_0000517i
Title : Structure of human Smoothened in complex with Vismodegib
Authors : Byrne, E.X.B.; Sircar, R.; Miller, P.S.; Hedger, G.; Luchetti, G.; Nachtergaele, S.; Tully, M.D.; Mydock-McGrane, L.; Covey, D.F.; Rambo, R.F.; Sansom, M.S.P.; Newstead, S.; Rohatgi, R.; Siebold, C.
Deposited on : 2016-06-03
Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	638	<div> <div>2%</div> <div>78%</div> <div>10%</div> <div>12%</div> </div>
1	B	638	<div> <div>7%</div> <div>80%</div> <div>9%</div> <div>10%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Smoothened homolog,Soluble cytochrome b562,Smoothened homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	564	Total	C	N	O	S	0	0	0
			4464	2870	764	798	32			
1	B	572	Total	C	N	O	S	0	0	0
			4524	2909	773	810	32			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	329	PHE	VAL	conflict	UNP Q99835
A	1011	ALA	-	linker	UNP Q99835
A	1012	ARG	-	linker	UNP Q99835
A	1013	ARG	-	linker	UNP Q99835
A	1014	GLN	-	linker	UNP Q99835
A	1015	LEU	-	linker	UNP Q99835
A	1022	TRP	MET	conflict	UNP P0ABE7
A	1117	ILE	HIS	conflict	UNP P0ABE7
A	1121	LEU	-	linker	UNP P0ABE7
A	1122	GLU	-	linker	UNP P0ABE7
A	1123	ARG	-	linker	UNP P0ABE7
A	1124	ALA	-	linker	UNP P0ABE7
A	1125	ARG	-	linker	UNP P0ABE7
A	1126	SER	-	linker	UNP P0ABE7
A	1127	THR	-	linker	UNP P0ABE7
A	1128	LEU	-	linker	UNP P0ABE7
A	556	GLY	-	expression tag	UNP Q99835
A	557	THR	-	expression tag	UNP Q99835
A	558	GLU	-	expression tag	UNP Q99835
A	559	THR	-	expression tag	UNP Q99835
A	560	SER	-	expression tag	UNP Q99835
A	561	GLN	-	expression tag	UNP Q99835
A	562	VAL	-	expression tag	UNP Q99835
A	563	ALA	-	expression tag	UNP Q99835

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	564	PRO	-	expression tag	UNP Q99835
A	565	ALA	-	expression tag	UNP Q99835
B	329	PHE	VAL	conflict	UNP Q99835
B	1011	ALA	-	linker	UNP Q99835
B	1012	ARG	-	linker	UNP Q99835
B	1013	ARG	-	linker	UNP Q99835
B	1014	GLN	-	linker	UNP Q99835
B	1015	LEU	-	linker	UNP Q99835
B	1022	TRP	MET	conflict	UNP P0ABE7
B	1117	ILE	HIS	conflict	UNP P0ABE7
B	1121	LEU	-	linker	UNP P0ABE7
B	1122	GLU	-	linker	UNP P0ABE7
B	1123	ARG	-	linker	UNP P0ABE7
B	1124	ALA	-	linker	UNP P0ABE7
B	1125	ARG	-	linker	UNP P0ABE7
B	1126	SER	-	linker	UNP P0ABE7
B	1127	THR	-	linker	UNP P0ABE7
B	1128	LEU	-	linker	UNP P0ABE7
B	556	GLY	-	expression tag	UNP Q99835
B	557	THR	-	expression tag	UNP Q99835
B	558	GLU	-	expression tag	UNP Q99835
B	559	THR	-	expression tag	UNP Q99835
B	560	SER	-	expression tag	UNP Q99835
B	561	GLN	-	expression tag	UNP Q99835
B	562	VAL	-	expression tag	UNP Q99835
B	563	ALA	-	expression tag	UNP Q99835
B	564	PRO	-	expression tag	UNP Q99835
B	565	ALA	-	expression tag	UNP Q99835

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is 2-chloranyl- {N}-(4-chloranyl-3-pyridin-2-yl-phenyl)-4-methylsulfonyl-benzamide (CCD ID: VIS) (formula: $C_{19}H_{14}Cl_2N_2O_3S$).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	S	0	0
			27	19	2	2	3	1		
3	B	1	Total	C	Cl	N	O	S	0	0
			27	19	2	2	3	1		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		

- Molecule 5 is [(Z)-octadec-9-enyl] (2R)-2,3-bis(oxidanyl)propanoate (CCD ID: MPG) (formula: C₂₁H₄₀O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			25	21	4		
5	B	1	Total	C	O	0	0
			25	21	4		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Smoothened homolog,Soluble cytochrome b562,Smoothened homolog



- Molecule 1: Smoothened homolog,Soluble cytochrome b562,Smoothened homolog



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	52.82Å 71.29Å 107.02Å 91.53° 98.20° 105.92°	Depositor
Resolution (Å)	59.32 – 3.30 59.32 – 3.30	Depositor EDS
% Data completeness (in resolution range)	94.5 (59.32-3.30) 92.4 (59.32-3.30)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 3.33Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.245 , 0.257 0.273 , 0.299	Depositor DCC
R_{free} test set	1017 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	94.1	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 80.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	9107	wwPDB-VP
Average B, all atoms (Å ²)	123.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MPG, NAG, NA, VIS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/4574	1.34	6/6212 (0.1%)
1	B	0.74	0/4636	1.34	14/6299 (0.2%)
All	All	0.74	0/9210	1.34	20/12511 (0.2%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	450	LEU	CA-C-N	6.09	128.44	120.28
1	B	450	LEU	C-N-CA	6.09	128.44	120.28
1	A	60	PRO	CA-C-N	5.77	129.46	120.75
1	A	60	PRO	C-N-CA	5.77	129.46	120.75
1	B	446	ASN	CA-C-N	5.71	128.72	120.38
1	B	446	ASN	C-N-CA	5.71	128.72	120.38
1	A	474	PHE	CA-C-N	5.38	127.44	120.44
1	A	474	PHE	C-N-CA	5.38	127.44	120.44
1	B	1036	ASP	CA-CB-CG	5.30	117.90	112.60
1	B	61	LEU	CA-C-N	5.28	127.30	120.44
1	B	61	LEU	C-N-CA	5.28	127.30	120.44
1	B	172	ASP	CA-CB-CG	5.22	117.82	112.60
1	B	550	CYS	CA-C-N	5.16	127.15	120.44
1	B	550	CYS	C-N-CA	5.16	127.15	120.44
1	A	61	LEU	CA-C-N	5.14	127.16	120.28
1	A	61	LEU	C-N-CA	5.14	127.16	120.28
1	B	171	PRO	CA-C-N	5.10	127.38	120.65
1	B	171	PRO	C-N-CA	5.10	127.38	120.65
1	B	155	ALA	CA-C-N	5.04	127.36	120.46
1	B	155	ALA	C-N-CA	5.04	127.36	120.46

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4464	0	4391	25	0
1	B	4524	0	4454	23	0
2	A	14	0	13	0	0
3	A	27	0	0	0	0
3	B	27	0	0	0	0
4	B	1	0	0	0	0
5	B	50	0	80	1	0
All	All	9107	0	8938	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1049:ARG:HH11	1:B:1091:LEU:HD21	1.52	0.73
1:B:296:ARG:HG3	1:B:302:ARG:HG3	1.78	0.65
1:B:382:ASP:HB2	1:B:394:TYR:HB2	1.81	0.63
1:A:382:ASP:HB2	1:A:394:TYR:HB2	1.81	0.63
1:B:363:LEU:HD23	1:B:367:LEU:HD11	1.89	0.55
1:B:312:LEU:HD21	5:B:1203:MPG:H112	1.90	0.54
1:B:252:PHE:HD1	1:B:259:SER:HB3	1.72	0.54
1:A:252:PHE:HD1	1:A:259:SER:HB3	1.72	0.53
1:A:1015:LEU:HD21	1:A:1128:LEU:HD13	1.91	0.53
1:A:1131:ILE:HA	1:A:449:MET:HB2	1.92	0.52
1:A:325:LEU:HD23	1:A:326:MET:HE2	1.91	0.52
1:B:325:LEU:HD23	1:B:326:MET:HE2	1.91	0.52
1:A:115:ALA:HB2	1:A:210:VAL:HG13	1.92	0.51
1:B:473:ASP:HA	1:B:480:TRP:HZ3	1.75	0.50
1:A:128:ALA:HB2	1:A:182:VAL:HG21	1.94	0.49
1:B:128:ALA:HB2	1:B:182:VAL:HG21	1.93	0.49
1:B:1106:ALA:HA	1:B:1109:LEU:HD12	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:ARG:HH22	1:A:140:GLU:HG2	1.79	0.48
1:B:312:LEU:HD23	1:B:316:ILE:HD11	1.96	0.47
1:B:241:THR:O	1:B:245:THR:HG22	2.15	0.47
1:A:241:THR:O	1:A:245:THR:HG22	2.15	0.47
1:A:68:ALA:HB2	1:A:91:LEU:HD21	1.98	0.46
1:A:1021:ASN:HA	1:A:1024:THR:OG1	2.17	0.45
1:A:197:LEU:HB3	1:A:213:CYS:HB3	1.99	0.45
1:A:1083:LEU:HG	1:A:1109:LEU:HD23	1.98	0.45
1:A:1018:LEU:HD11	1:A:1076:PHE:HZ	1.81	0.45
1:A:541:THR:HA	1:A:544:ILE:HD12	1.98	0.45
1:B:473:ASP:HA	1:B:480:TRP:CZ3	2.52	0.44
1:A:248:THR:HA	1:A:536:VAL:HG21	2.00	0.43
1:A:256:TRP:NE1	1:B:312:LEU:HD13	2.32	0.43
1:A:329:PHE:HB3	1:A:411:VAL:HG11	2.00	0.43
1:B:302:ARG:HD3	1:B:305:GLU:HG3	2.00	0.43
1:B:340:HIS:CE1	1:B:421:ARG:HB3	2.53	0.43
1:B:248:THR:HA	1:B:536:VAL:HG21	1.99	0.43
1:A:172:ASP:O	1:A:173:ARG:HG2	2.18	0.43
1:B:541:THR:HA	1:B:544:ILE:HD12	2.00	0.43
1:A:473:ASP:HA	1:A:480:TRP:HZ3	1.83	0.42
1:B:352:PRO:C	1:B:354:SER:H	2.27	0.42
1:A:293:ILE:O	1:A:302:ARG:HD2	2.20	0.42
1:A:450:LEU:O	1:A:454:ILE:HG12	2.20	0.42
1:A:461:GLY:O	1:A:465:ILE:HG13	2.19	0.42
1:B:329:PHE:HB3	1:B:411:VAL:HG11	2.01	0.42
1:B:1025:LEU:HD23	1:B:1048:MET:HE3	2.02	0.41
1:B:461:GLY:O	1:B:465:ILE:HG12	2.20	0.41
1:B:1016:ALA:O	1:B:1020:ASP:HB2	2.20	0.41
1:A:1083:LEU:O	1:A:1087:ILE:HG13	2.20	0.41
1:A:275:PHE:O	1:A:279:ILE:HG13	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	556/638 (87%)	540 (97%)	14 (2%)	2 (0%)	30	60
1	B	566/638 (89%)	545 (96%)	19 (3%)	2 (0%)	30	60
All	All	1122/1276 (88%)	1085 (97%)	33 (3%)	4 (0%)	30	60

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	313	SER
1	B	446	ASN
1	A	313	SER
1	A	115	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	474/529 (90%)	457 (96%)	17 (4%)	31	58
1	B	480/529 (91%)	467 (97%)	13 (3%)	39	63
All	All	954/1058 (90%)	924 (97%)	30 (3%)	35	60

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	SER
1	A	267	LEU
1	A	303	LEU
1	A	308	SER
1	A	311	THR
1	A	317	ILE
1	A	362	LEU
1	A	389	ILE
1	A	1015	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1025	LEU
1	A	1049	ARG
1	A	1081	ASP
1	A	1083	LEU
1	A	1100	LYS
1	A	1107	GLU
1	A	1109	LEU
1	A	477	GLN
1	B	91	LEU
1	B	194	GLU
1	B	257	ARG
1	B	311	THR
1	B	317	ILE
1	B	362	LEU
1	B	389	ILE
1	B	1025	LEU
1	B	1037	ASN
1	B	1083	LEU
1	B	1100	LYS
1	B	1107	GLU
1	B	477	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	202	ASN
1	A	260	ASN
1	A	271	ASN
1	A	1078	HIS
1	A	1118	GLN
1	A	477	GLN
1	B	63	HIS
1	B	202	ASN
1	B	260	ASN
1	B	271	ASN
1	B	340	HIS
1	B	380	GLN
1	B	1037	ASN
1	B	1086	GLN
1	B	1114	ASN
1	B	1118	GLN
1	B	476	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	477	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	MPG	B	1204	-	24,24,24	1.33	1 (4%)	24,25,25	1.21	2 (8%)
2	NAG	A	1201	1	14,14,15	0.32	0	17,19,21	0.45	0
5	MPG	B	1203	-	24,24,24	1.32	1 (4%)	24,25,25	1.21	3 (12%)
3	VIS	B	1202	-	29,29,29	0.20	0	42,42,42	0.31	0
3	VIS	A	1202	-	29,29,29	0.19	0	42,42,42	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MPG	B	1204	-	-	14/25/25/25	-
2	NAG	A	1201	1	-	0/6/23/26	0/1/1/1
5	MPG	B	1203	-	-	14/25/25/25	-
3	VIS	B	1202	-	-	6/18/18/18	0/3/3/3
3	VIS	A	1202	-	-	6/18/18/18	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1204	MPG	O1-CX3	6.23	1.46	1.33
5	B	1203	MPG	O1-CX3	6.08	1.46	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1203	MPG	O1-CX3-CXD	3.88	120.77	111.78
5	B	1204	MPG	O1-CX3-CXD	3.73	120.40	111.78
5	B	1204	MPG	O1-CX3-O4	-3.13	118.41	124.14
5	B	1203	MPG	O1-CX3-O4	-3.11	118.44	124.14
5	B	1203	MPG	C1-O1-CX3	2.34	121.68	116.53

There are no chirality outliers.

All (40) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	1203	MPG	CXD-CX3-O1-C1
5	B	1203	MPG	O3-C21-CXD-O2
5	B	1204	MPG	CXD-CX3-O1-C1
5	B	1204	MPG	O3-C21-CXD-O2
5	B	1204	MPG	O3-C21-CXD-CX3
5	B	1203	MPG	O4-CX3-O1-C1
5	B	1204	MPG	O4-CX3-O1-C1
3	B	1202	VIS	C9-C8-N-C7
3	A	1202	VIS	C9-C8-N-C7
3	B	1202	VIS	C13-C8-N-C7
3	A	1202	VIS	C13-C8-N-C7
5	B	1203	MPG	C4-C5-C6-C7
5	B	1204	MPG	C4-C5-C6-C7
5	B	1204	MPG	C2-C3-C4-C5
5	B	1203	MPG	C12-C13-C14-C15
5	B	1204	MPG	C1-C2-C3-C4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	B	1204	MPG	C14-C15-C16-C17
5	B	1203	MPG	C14-C15-C16-C17
5	B	1204	MPG	C12-C13-C14-C15
5	B	1203	MPG	C6-C7-C8-C9
5	B	1203	MPG	C10-C11-C12-C13
3	A	1202	VIS	C13-C12-C14-N1
3	B	1202	VIS	C13-C12-C14-C15
5	B	1203	MPG	O3-C21-CXD-CX3
5	B	1204	MPG	C6-C7-C8-C9
3	B	1202	VIS	C13-C12-C14-N1
3	A	1202	VIS	C13-C12-C14-C15
5	B	1204	MPG	O1-C1-C2-C3
5	B	1203	MPG	C1-C2-C3-C4
5	B	1203	MPG	C5-C6-C7-C8
3	B	1202	VIS	C11-C12-C14-N1
5	B	1203	MPG	C15-C16-C17-C18
3	B	1202	VIS	C11-C12-C14-C15
5	B	1203	MPG	C9-C10-C11-C12
5	B	1204	MPG	C3-C4-C5-C6
3	A	1202	VIS	C11-C12-C14-N1
5	B	1204	MPG	C13-C14-C15-C16
3	A	1202	VIS	C11-C12-C14-C15
5	B	1203	MPG	C3-C4-C5-C6
5	B	1204	MPG	C10-C11-C12-C13



There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1203	MPG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient


equivalents in the CSD to analyse the geometry.



Ligand VIS B 1202



Ligand VIS A 1202



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	564/638 (88%)	0.07	12 (2%) 63 44	68, 127, 170, 206	0
1	B	572/638 (89%)	0.24	44 (7%) 19 14	62, 116, 173, 254	0
All	All	1136/1276 (89%)	0.16	56 (4%) 35 23	62, 120, 171, 254	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1011	ALA	10.3
1	B	343	PHE	8.8
1	A	1129	SER	7.3
1	A	1130	LYS	6.8
1	B	220	PRO	6.7
1	B	426	LEU	6.7
1	B	257	ARG	5.4
1	B	219	ASN	5.3
1	B	425	THR	5.2
1	B	215	ILE	4.9
1	B	428	SER	4.8
1	B	218	GLN	4.7
1	B	484	PHE	4.6
1	B	488	VAL	4.4
1	B	221	LEU	4.2
1	B	339	TRP	4.1
1	B	446	ASN	3.9
1	B	227	HIS	3.9
1	B	346	LEU	3.8
1	B	207	TYR	3.8
1	A	446	ASN	3.7
1	B	1012	ARG	3.5
1	B	1131	ILE	3.5
1	B	342	SER	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	210	VAL	3.1
1	B	450	LEU	3.1
1	B	1014	GLN	3.1
1	B	196	PRO	3.1
1	A	69	PRO	3.0
1	B	112	LEU	2.9
1	B	197	LEU	2.9
1	B	447	GLU	2.9
1	B	295	CYS	2.6
1	A	1011	ALA	2.6
1	B	1017	ASP	2.6
1	A	283	ALA	2.5
1	B	1015	LEU	2.5
1	B	1035	ALA	2.4
1	B	1038	ALA	2.4
1	A	1126	SER	2.4
1	B	345	ALA	2.4
1	A	137	ASP	2.3
1	B	516	LEU	2.3
1	B	216	GLN	2.3
1	B	495	THR	2.3
1	B	360	PHE	2.2
1	B	1055	ALA	2.2
1	B	213	CYS	2.2
1	B	303	LEU	2.2
1	A	448	THR	2.2
1	B	427	PHE	2.1
1	B	195	VAL	2.1
1	B	420	ILE	2.1
1	A	1105	ALA	2.1
1	A	71	GLU	2.1
1	A	136	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	A	1201	14/15	0.66	0.12	163,164,165,165	0
5	MPG	B	1204	25/25	0.79	0.12	81,93,121,121	0
5	MPG	B	1203	25/25	0.81	0.14	74,85,105,105	0
3	VIS	B	1202	27/27	0.84	0.15	135,142,155,156	0
3	VIS	A	1202	27/27	0.86	0.12	109,113,120,120	0
4	NA	B	1201	1/1	0.97	0.06	49,49,49,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around MPG B 1203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around VIS B 1202:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.