



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:35 AM UTC

PDB ID : 5N7T / pdb\_00005n7t  
Title : Pseudomonas fluorescens kynurenine 3-monooxygenase (KMO) in complex with 3-(5,6-dichloro-2-oxo-2,3-dihydro-1,3-benzoxazol-3-yl)propanoic acid  
Authors : Rowland, P.  
Deposited on : 2017-02-21  
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	<div> <div>5%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div>.</div> </div> </div>
1	B	461	<div> <div>5%</div> <div> <div></div> <div>94%</div> <div>.</div> <div>.</div> </div> </div>

2 Entry composition ⓘ

There are 6 unique types of molecules in this entry. The entry contains 8116 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Kynurenine 3-monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	448	Total	C	N	O	S	0	3	0
			3482	2182	643	639	18			
1	B	451	Total	C	N	O	S	0	4	0
			3511	2201	650	642	18			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	252	SER	CYS	engineered mutation	UNP Q84HF5
A	461	SER	CYS	engineered mutation	UNP Q84HF5
B	252	SER	CYS	engineered mutation	UNP Q84HF5
B	461	SER	CYS	engineered mutation	UNP Q84HF5

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is 3-(5,6-DICHLORO-2-OXOBENZO[D]OXAZOL-3(2H)-YL)PROPANOIC ACID (CCD ID: JHY) (formula: C<sub>10</sub>H<sub>7</sub>Cl<sub>2</sub>NO<sub>4</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			17	10	2	1	4		
4	B	1	Total	C	Cl	N	O	0	0
			17	10	2	1	4		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	502	Total	O	0	0
			502	502		
6	B	431	Total	O	0	0
			431	431		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kynurenine 3-monooxygenase



- Molecule 1: Kynurenine 3-monooxygenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.73Å 53.11Å 136.49Å 90.00° 103.81° 90.00°	Depositor
Resolution (Å)	132.55 – 1.81 132.55 – 1.81	Depositor EDS
% Data completeness (in resolution range)	99.2 (132.55-1.81) 95.8 (132.55-1.81)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 1.81Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, $R_{free}$	0.163 , 0.188 (Not available) , 0.192	Depositor DCC
$R_{free}$ test set	3509 reflections (3.64%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.0	Xtriage
Anisotropy	0.259	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 53.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8116	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.68 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.0358e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: JHY, FAD, CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.85	0/3566	1.11	4/4843 (0.1%)
1	B	0.83	0/3599	1.08	2/4888 (0.0%)
All	All	0.84	0/7165	1.09	6/9731 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	330	ASP	CA-CB-CG	7.04	119.64	112.60
1	B	330	ASP	CA-CB-CG	6.60	119.20	112.60
1	B	272	ASP	CA-CB-CG	5.77	118.37	112.60
1	A	52	SER	CA-C-N	5.50	127.94	120.63
1	A	52	SER	C-N-CA	5.50	127.94	120.63
1	A	165	ALA	N-CA-C	-5.11	103.99	111.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3482	0	3446	14	0
1	B	3511	0	3487	8	0
2	A	53	0	31	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	53	0	31	0	0
3	A	1	0	0	1	0
3	B	1	0	0	1	0
4	A	17	0	6	0	0
4	B	17	0	6	0	0
5	A	24	0	32	0	0
5	B	24	0	32	0	0
6	A	502	0	0	2	0
6	B	431	0	0	1	0
All	All	8116	0	7071	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:84:ARG:HD2	1:B:215:ILE:HD12	1.62	0.81
1:A:77:GLU:HB3	1:A:229:LEU:HD11	1.76	0.68
1:A:184:GLU:HG2	1:A:297:LEU:HD22	1.77	0.65
1:B:185:HIS:HD2	6:B:637:HOH:O	1.82	0.63
1:A:318:PRO:HA	3:A:502:CL:CL	2.40	0.58
1:B:318:PRO:HA	3:B:502:CL:CL	2.41	0.58
1:A:77:GLU:HB2	1:A:113:ARG:NH2	2.21	0.55
1:A:229:LEU:HG	6:A:913:HOH:O	2.07	0.55
1:B:84:ARG:HD2	1:B:215:ILE:CD1	2.34	0.54
1:A:219:GLY:HA2	1:A:375:SER:HB3	1.92	0.51
1:A:379:SER:HB2	1:A:382:TYR:HB3	1.96	0.47
1:B:86:VAL:HG12	1:B:88:VAL:HG23	1.96	0.47
1:A:77:GLU:CB	1:A:229:LEU:HD11	2.42	0.46
1:B:213:LEU:HD21	1:B:224:ILE:HD11	1.98	0.46
1:A:51:ARG:HB3	1:A:53:ILE:HG12	1.99	0.45
1:A:84:ARG:HG3	1:A:215:ILE:HD12	1.99	0.44
1:B:55:LEU:HD21	1:B:111:ARG:NH2	2.34	0.42
2:A:501:FAD:H9	2:A:501:FAD:H1'1	1.88	0.42
1:A:277:LEU:HD13	1:A:280:LEU:HD13	2.02	0.42
1:A:299:THR:HA	6:A:858:HOH:O	2.20	0.42
1:A:17:LEU:HD13	1:A:324[B]:MET:HG3	2.01	0.41
1:B:63:HIS:CG	1:B:411:ARG:HD3	2.56	0.41
1:A:438:LEU:HA	1:A:441:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	447/461 (97%)	436 (98%)	9 (2%)	2 (0%)	30	19
1	B	453/461 (98%)	446 (98%)	7 (2%)	0	100	100
All	All	900/922 (98%)	882 (98%)	16 (2%)	2 (0%)	43	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	433	ALA
1	A	380	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/372 (98%)	359 (99%)	5 (1%)	59	49
1	B	367/372 (99%)	361 (98%)	6 (2%)	55	44
All	All	731/744 (98%)	720 (98%)	11 (2%)	57	46

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	55	LEU
1	A	109	ILE
1	A	427	LEU
1	A	438	LEU
1	B	45	GLU
1	B	101	ASP
1	B	104	GLU
1	B	272	ASP
1	B	427	LEU
1	B	438	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	32	GLN
1	A	63	HIS
1	A	390	GLN
1	A	434	ASN
1	B	34	ASN
1	B	197	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	A	504	-	5,5,5	0.19	0	5,5,5	0.79	0
2	FAD	A	501	-	58,58,58	0.32	0	85,89,89	0.47	0
5	GOL	B	504	-	5,5,5	0.06	0	5,5,5	0.34	0
4	JHY	A	503	-	18,18,18	0.63	0	26,26,26	0.91	0
4	JHY	B	503	-	18,18,18	0.61	0	26,26,26	0.94	1 (3%)
5	GOL	B	506	-	5,5,5	0.08	0	5,5,5	0.33	0
5	GOL	B	507	-	5,5,5	0.11	0	5,5,5	0.42	0
5	GOL	A	507	-	5,5,5	0.07	0	5,5,5	0.61	0
5	GOL	A	506	-	5,5,5	0.07	0	5,5,5	0.22	0
2	FAD	B	501	-	58,58,58	0.40	0	85,89,89	0.48	0
5	GOL	A	505	-	5,5,5	0.14	0	5,5,5	0.26	0
5	GOL	B	505	-	5,5,5	0.12	0	5,5,5	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	504	-	-	2/4/4/4	-
2	FAD	A	501	-	-	1/34/50/50	0/6/6/6
5	GOL	B	504	-	-	0/4/4/4	-
4	JHY	A	503	-	-	0/5/5/5	0/2/2/2
4	JHY	B	503	-	-	2/5/5/5	0/2/2/2
5	GOL	B	506	-	-	0/4/4/4	-
5	GOL	B	507	-	-	0/4/4/4	-
5	GOL	A	507	-	-	0/4/4/4	-
5	GOL	A	506	-	-	0/4/4/4	-
2	FAD	B	501	-	-	1/34/50/50	0/6/6/6
5	GOL	A	505	-	-	0/4/4/4	-
5	GOL	B	505	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	JHY	O2-C8-C7	2.03	129.21	124.95

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	504	GOL	C1-C2-C3-O3
5	A	504	GOL	O2-C2-C3-O3
4	B	503	JHY	O-C-C1-C2
2	A	501	FAD	C1'-C2'-C3'-O3'
2	B	501	FAD	C1'-C2'-C3'-O3'
4	B	503	JHY	O1-C-C1-C2

There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## Ligand JHY B 503





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	448/461 (97%)	-0.06	21 (4%) 36 43	10, 26, 68, 89	3 (0%)
1	B	451/461 (97%)	0.09	24 (5%) 32 37	11, 30, 66, 92	4 (0%)
All	All	899/922 (97%)	0.01	45 (5%) 34 40	10, 28, 67, 92	7 (0%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	245	ALA	4.3
1	B	382	TYR	4.3
1	B	383	LEU	4.2
1	B	151	SER	3.8
1	A	90	GLY	3.5
1	A	457	LEU	3.4
1	A	438	LEU	3.3
1	A	434	ASN	3.2
1	B	457	LEU	3.1
1	B	150	VAL	3.0
1	A	380	PRO	3.0
1	B	101	ASP	2.9
1	A	432	VAL	2.9
1	B	380	PRO	2.8
1	B	177	ALA	2.8
1	B	393	ALA	2.8
1	A	381	THR	2.8
1	A	383	LEU	2.8
1	B	246	ALA	2.7
1	B	381	THR	2.7
1	A	375	SER	2.7
1	B	185	HIS	2.7
1	B	102	ASP	2.6
1	A	382	TYR	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	394	GLN	2.5
1	A	443	LEU	2.5
1	B	379	SER	2.5
1	B	155	LEU	2.5
1	B	384	LEU	2.5
1	A	245	ALA	2.5
1	A	379	SER	2.5
1	B	448	HIS	2.4
1	A	451	THR	2.4
1	B	378	ALA	2.4
1	B	100	ARG	2.4
1	A	453	CYS	2.4
1	A	433	ALA	2.3
1	A	441	ILE	2.3
1	A	430	PHE	2.3
1	B	152	GLY	2.3
1	A	100	ARG	2.2
1	B	394	GLN	2.1
1	B	248	PRO	2.0
1	A	246	ALA	2.0
1	B	436	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	B	507	6/6	0.79	0.18	59,62,63,63	0
5	GOL	A	507	6/6	0.83	0.16	40,45,50,52	0

*Continued on next page...*




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	B	506	6/6	0.83	0.16	59,61,61,62	0
5	GOL	A	506	6/6	0.83	0.18	74,76,77,77	0
5	GOL	A	504	6/6	0.87	0.14	25,42,43,48	0
5	GOL	A	505	6/6	0.88	0.13	33,45,45,48	0
5	GOL	B	505	6/6	0.90	0.11	30,39,42,43	0
5	GOL	B	504	6/6	0.92	0.09	25,36,38,39	0
4	JHY	A	503	17/17	0.95	0.08	20,25,32,41	0
4	JHY	B	503	17/17	0.96	0.06	24,25,31,35	0
2	FAD	B	501	53/53	0.99	0.04	16,19,24,27	0
3	CL	B	502	1/1	0.99	0.07	26,26,26,26	0
2	FAD	A	501	53/53	0.99	0.04	14,17,20,21	0
3	CL	A	502	1/1	1.00	0.06	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




#### Electron density around JHY B 503:

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FAD B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FAD A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.