



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 10:21 AM UTC

PDB ID : 5WG3 / pdb\_00005wg3  
Title : Human GRK2 in complex with Gbetagamma subunits and CCG258748  
Authors : Bouley, R.; Tesmer, J.J.G.  
Deposited on : 2017-07-13  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	689	
2	B	340	
3	G	71	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8184 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-adrenergic receptor kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	618	Total	C	N	O	S	0	0	0
			5059	3224	881	920	34			

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	339	Total	C	N	O	S	0	0	0
			2607	1607	468	511	21			


- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	57	Total	C	N	O	S	0	0	0
			438	274	77	84	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	68	SER	CYS	engineered mutation	UNP P63212

- Molecule 4 is 2-fluoro-5-[(3S,4R)-3-[[[(1H-indazol-5-yl)oxy]methyl]piperidin-4-yl]-N-[(1H-pyrazol-3-yl)methyl]benzamide (CCD ID: AFM) (formula: C<sub>24</sub>H<sub>25</sub>FN<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0	0
			33	24	1	6	2		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	26	Total	O	0	0
			26	26		
6	B	14	Total	O	0	0
			14	14		
6	G	6	Total	O	0	0
			6	6		



- Molecule 1: Beta-adrenergic receptor kinase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.27Å 241.57Å 214.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.07 – 2.90 49.07 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (49.07-2.90) 99.2 (49.07-2.90)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.54 (at 2.91Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.198 , 0.248 0.200 , 0.250	Depositor DCC
$R_{free}$ test set	1765 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	69.1	Xtriage
Anisotropy	0.748	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 65.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8184	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AFM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.14	0/5169	0.32	0/6944
2	B	0.16	0/2654	0.36	0/3597
3	G	0.13	0/444	0.28	0/599
All	All	0.15	0/8267	0.33	0/11140

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5059	0	5039	42	0
2	B	2607	0	2510	23	0
3	G	438	0	447	2	0
4	A	33	0	0	0	0
5	A	1	0	0	0	0
6	A	26	0	0	1	0
6	B	14	0	0	0	0
6	G	6	0	0	0	0
All	All	8184	0	7996	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:ARG:NH2	1:A:166:ILE:O	2.29	0.66
1:A:240:ARG:HH21	1:A:511:LEU:HB2	1.62	0.64
2:B:51:LEU:HB2	2:B:336:LEU:HB2	1.81	0.63
2:B:45:MET:HE3	2:B:308:LEU:HD21	1.81	0.63
2:B:54:HIS:O	2:B:334:SER:HB2	1.98	0.62
1:A:315:TYR:OH	1:A:334:SER:O	2.17	0.61
2:B:294:CYS:HB3	2:B:308:LEU:HB2	1.83	0.59
1:A:319:LYS:HE3	1:A:321:ALA:HB3	1.83	0.59
1:A:173:ARG:NH2	1:A:534:ASP:OD1	2.37	0.57
1:A:223:ASP:HB3	1:A:226:ARG:HB2	1.87	0.56
2:B:160:SER:HB3	2:B:190:LEU:HD23	1.87	0.55
1:A:587:ARG:HD2	1:A:600:LEU:HD22	1.87	0.55
1:A:609:VAL:HG22	1:A:622:LEU:HD22	1.89	0.55
1:A:442:ARG:HB2	1:A:446:GLU:HG3	1.90	0.53
1:A:294:MET:HE3	1:A:381:LEU:HB3	1.89	0.53
1:A:664:MET:HE3	2:B:101:MET:HG3	1.91	0.53
2:B:320:VAL:HG22	2:B:327:VAL:HG22	1.92	0.52
2:B:271:CYS:HB2	2:B:290:ASP:HB3	1.91	0.52
1:A:40:ARG:NH1	1:A:172:THR:OG1	2.44	0.51
2:B:79:LEU:HG	2:B:95:LEU:HD21	1.94	0.50
2:B:192:LEU:HD23	2:B:199:PHE:HB3	1.93	0.50
1:A:605:GLU:HB3	1:A:624:ILE:HG23	1.92	0.50
1:A:358:ALA:HB3	1:A:361:VAL:HG23	1.94	0.49
1:A:314:VAL:HG12	1:A:370:SER:HA	1.94	0.49
1:A:606:ILE:O	1:A:655:GLN:NE2	2.45	0.49
2:B:27:ASP:OD1	2:B:28:ALA:N	2.45	0.49
1:A:565:MET:HE3	1:A:634:CYS:SG	2.52	0.49
1:A:603:MET:HB3	1:A:651:TYR:HA	1.94	0.49
1:A:382:PHE:HB3	1:A:390:PRO:HG3	1.94	0.48
1:A:426:GLU:O	1:A:430:GLN:HG2	2.14	0.48
1:A:294:MET:HE2	1:A:298:ALA:HB2	1.95	0.48
2:B:321:THR:OG1	2:B:323:ASP:OD1	2.32	0.47
1:A:601:LEU:HD11	1:A:624:ILE:HD12	1.95	0.47
1:A:360:GLU:CD	1:A:436:ARG:HH22	2.22	0.47
1:A:433:VAL:HG13	1:A:436:ARG:HH21	1.79	0.46
2:B:146:LEU:HD11	2:B:159:THR:HB	1.97	0.46
1:A:388:HIS:HD2	1:A:392:ARG:HB2	1.81	0.46
1:A:565:MET:HE2	1:A:632:LEU:HB3	1.98	0.46

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:350:SER:HB3	1:A:364:LYS:HZ2	1.80	0.45
1:A:71:PHE:HB2	1:A:170:LYS:HG3	1.98	0.45
2:B:30:LEU:HD23	2:B:262:MET:HE2	1.98	0.45
1:A:105:SER:HB3	1:A:136:LEU:HD21	1.97	0.45
2:B:274:THR:OG1	2:B:315:VAL:O	2.21	0.45
2:B:340:ASN:ND2	3:G:59:ASN:OD1	2.47	0.45
1:A:113:ILE:HA	1:A:122:HIS:CE1	2.52	0.44
2:B:126:LEU:HA	2:B:133:VAL:HG12	1.98	0.44
2:B:48:ARG:HG3	2:B:340:ASN:HB3	1.98	0.44
1:A:296:PHE:CD2	1:A:469:PRO:HD3	2.52	0.44
2:B:166:CYS:HB2	2:B:180:PHE:HB2	1.98	0.44
1:A:568:MET:HE3	1:A:576:TRP:CE2	2.53	0.43
2:B:164:THR:HG22	2:B:185:GLY:C	2.43	0.43
2:B:51:LEU:HB3	2:B:82:TRP:CE3	2.53	0.43
1:A:172:THR:O	1:A:176:GLN:HG3	2.19	0.43
1:A:628:LYS:HA	1:A:628:LYS:HD2	1.84	0.43
3:G:21:MET:HE3	3:G:21:MET:HB3	1.94	0.42
1:A:62:LYS:HD3	1:A:62:LYS:HA	1.89	0.42
1:A:526:PHE:O	1:A:530:ASN:ND2	2.53	0.42
1:A:326:ASP:OD1	1:A:330:HIS:N	2.53	0.41
2:B:211:TRP:CZ3	2:B:218:CYS:HB2	2.55	0.41
2:B:199:PHE:HD1	2:B:213:VAL:HG22	1.86	0.41
1:A:516:ARG:NH1	6:A:802:HOH:O	2.54	0.40
1:A:320:PRO:HD3	1:A:380:MET:HG3	2.04	0.40
1:A:56:GLU:N	1:A:56:GLU:OE1	2.53	0.40
1:A:139:LYS:HE3	1:A:139:LYS:HB2	1.91	0.40
1:A:581:PHE:CD2	1:A:590:TRP:HB3	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	606/689 (88%)	576 (95%)	29 (5%)	1 (0%)	43	72
2	B	337/340 (99%)	318 (94%)	17 (5%)	2 (1%)	21	51
3	G	55/71 (78%)	53 (96%)	2 (4%)	0	100	100
All	All	998/1100 (91%)	947 (95%)	48 (5%)	3 (0%)	36	65

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	54	HIS
1	A	316	ARG
2	B	141	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	553/609 (91%)	548 (99%)	5 (1%)	70	90
2	B	282/283 (100%)	280 (99%)	2 (1%)	76	92
3	G	46/58 (79%)	44 (96%)	2 (4%)	26	60
All	All	881/950 (93%)	872 (99%)	9 (1%)	68	89

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	VAL
1	A	98	GLU
1	A	411	VAL
1	A	433	VAL
1	A	641	VAL
2	B	54	HIS
2	B	292	PHE
3	G	16	VAL
3	G	58	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	519	GLN
1	A	613	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	AFM	A	701	-	37,37,37	1.74	8 (21%)	43,51,51	2.00	14 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AFM	A	701	-	-	8/18/29/29	0/5/5/5

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	701	AFM	C25-N26	5.56	1.46	1.33
4	A	701	AFM	C05-C06	4.84	1.59	1.51
4	A	701	AFM	C22-C06	-3.10	1.49	1.53
4	A	701	AFM	C08-C07	2.38	1.56	1.51
4	A	701	AFM	N14-N15	-2.25	1.35	1.36
4	A	701	AFM	C27-C28	2.12	1.53	1.49
4	A	701	AFM	C17-C13	-2.06	1.37	1.41
4	A	701	AFM	O33-C25	-2.01	1.18	1.23

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	701	AFM	C27-C28-N32	6.30	126.33	120.01
4	A	701	AFM	C16-N15-N14	4.28	108.25	106.02
4	A	701	AFM	C17-C16-N15	-4.11	108.68	111.68
4	A	701	AFM	C13-N14-N15	-3.07	110.02	111.52
4	A	701	AFM	C03-C02-C24	-2.94	119.91	123.07
4	A	701	AFM	C19-C07-C06	2.81	114.18	109.88
4	A	701	AFM	C23-C24-C02	2.80	120.31	116.83
4	A	701	AFM	C12-C13-C17	-2.66	119.72	122.13
4	A	701	AFM	C22-C06-C07	2.56	112.98	109.75
4	A	701	AFM	C05-C06-C07	-2.51	108.84	113.67
4	A	701	AFM	C22-C06-C05	-2.50	107.94	112.54
4	A	701	AFM	C28-N32-N31	2.24	110.47	105.38
4	A	701	AFM	C30-N31-N32	-2.20	107.27	111.60
4	A	701	AFM	C18-C17-C13	2.05	120.39	119.13

There are no chirality outliers.


All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	701	AFM	N26-C27-C28-C29
4	A	701	AFM	N26-C27-C28-N32
4	A	701	AFM	C06-C07-C08-O09
4	A	701	AFM	C19-C07-C08-O09
4	A	701	AFM	C02-C24-C25-N26
4	A	701	AFM	C11-C10-O09-C08
4	A	701	AFM	C18-C10-O09-C08
4	A	701	AFM	C02-C24-C25-O33

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	618/689 (89%)	0.01	9 (1%) 72 64	50, 99, 152, 179	0
2	B	339/340 (99%)	-0.36	0 100 100	56, 76, 111, 161	0
3	G	57/71 (80%)	-0.20	0 100 100	67, 91, 137, 148	0
All	All	1014/1100 (92%)	-0.13	9 (0%) 81 75	50, 90, 146, 179	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	411	VAL	5.0
1	A	482	ALA	2.9
1	A	479	ALA	2.6
1	A	396	THR	2.6
1	A	359	PRO	2.5
1	A	477	VAL	2.4
1	A	75	HIS	2.1
1	A	362	LEU	2.0
1	A	407	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	AFM	A	701	33/33	0.95	0.10	51,85,104,112	0
5	MG	A	702	1/1	0.95	0.06	139,139,139,139	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.