



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:58 PM UTC

PDB ID : 5ZQU / pdb\_00005zqu  
Title : Crystal structure of tetrameric RXRalpha-LBD complexed with partial agonist CBt-PMN  
Authors : Miyashita, Y.; Numoto, N.; Arulmozhiraja, S.; Nakano, S.; Matsuo, N.; Shimizu, K.; Kakuta, H.; Ito, S.; Ikura, T.; Ito, N.; Tokiwa, H.  
Deposited on : 2018-04-20  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	<div> <div>3%</div> <div>78%</div> <div>19%</div> </div>
1	B	243	<div> <div>4%</div> <div>76%</div> <div>5%</div> <div>19%</div> </div>
1	C	243	<div> <div>4%</div> <div>75%</div> <div>5%</div> <div>19%</div> </div>
1	D	243	<div> <div>5%</div> <div>75%</div> <div>6%</div> <div>19%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Retinoic acid receptor RXR-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	198	Total	C	N	O	S	0	0	0
			1566	1007	272	278	9			
1	B	198	Total	C	N	O	S	0	0	0
			1562	1005	270	278	9			
1	C	197	Total	C	N	O	S	0	0	0
			1555	1000	269	277	9			
1	D	196	Total	C	N	O	S	0	0	0
			1549	998	269	273	9			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	220	GLY	-	expression tag	UNP P19793
A	221	SER	-	expression tag	UNP P19793
A	222	SER	-	expression tag	UNP P19793
A	223	HIS	-	expression tag	UNP P19793
B	220	GLY	-	expression tag	UNP P19793
B	221	SER	-	expression tag	UNP P19793
B	222	SER	-	expression tag	UNP P19793
B	223	HIS	-	expression tag	UNP P19793
C	220	GLY	-	expression tag	UNP P19793
C	221	SER	-	expression tag	UNP P19793
C	222	SER	-	expression tag	UNP P19793
C	223	HIS	-	expression tag	UNP P19793
D	220	GLY	-	expression tag	UNP P19793
D	221	SER	-	expression tag	UNP P19793
D	222	SER	-	expression tag	UNP P19793
D	223	HIS	-	expression tag	UNP P19793

- Molecule 2 is 1-(3,5,5,8,8-pentamethyl-6,7-dihydronaphthalen-2-yl)benzotriazole-5-carboxylic acid (CCD ID: 9HF) (formula: C<sub>22</sub>H<sub>25</sub>N<sub>3</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			27	22	3	2		
2	A	1	Total	C	N	O	0	0
			27	22	3	2		
2	B	1	Total	C	N	O	0	0
			27	22	3	2		
2	C	1	Total	C	N	O	0	0
			27	22	3	2		
2	D	1	Total	C	N	O	0	0
			27	22	3	2		
2	D	1	Total	C	N	O	0	0
			27	22	3	2		

- Molecule 3 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Br	0	0
			4	4		
3	B	3	Total	Br	0	0
			3	3		
3	C	2	Total	Br	0	0
			2	2		
3	D	2	Total	Br	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	42	Total 42	O 42	0	0
4	B	36	Total 36	O 36	0	0
4	C	19	Total 19	O 19	0	0
4	D	22	Total 22	O 22	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Retinoic acid receptor RXR-alpha




- Molecule 1: Retinoic acid receptor RXR-alpha




- Molecule 1: Retinoic acid receptor RXR-alpha



- Molecule 1: Retinoic acid receptor RXR-alpha





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.57Å 100.49Å 111.01Å 90.00° 99.27° 90.00°	Depositor
Resolution (Å)	36.52 – 2.60 36.52 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (36.52-2.60) 91.9 (36.52-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.07 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.222 , 0.270 0.231 , 0.276	Depositor DCC
$R_{free}$ test set	1587 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.4	Xtriage
Anisotropy	0.281	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 31.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.037 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6524	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 9HF, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.29	0/1600	0.74	0/2164
1	B	0.29	0/1596	0.74	0/2159
1	C	0.29	0/1588	0.75	2/2148 (0.1%)
1	D	0.29	0/1583	0.75	0/2140
All	All	0.29	0/6367	0.75	2/8611 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	263	ASP	CA-C-N	5.44	124.90	119.24
1	C	263	ASP	C-N-CA	5.44	124.90	119.24

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1566	0	1599	4	0
1	B	1562	0	1598	7	0
1	C	1555	0	1590	8	0
1	D	1549	0	1588	8	0
2	A	54	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	27	0	0	0	0
2	C	27	0	0	0	0
2	D	54	0	0	0	0
3	A	4	0	0	0	0
3	B	3	0	0	1	0
3	C	2	0	0	0	0
3	D	2	0	0	1	0
4	A	42	0	0	0	0
4	B	36	0	0	0	0
4	C	19	0	0	1	0
4	D	22	0	0	0	0
All	All	6524	0	6375	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:452:MET:HG3	1:D:294:LEU:HD21	1.86	0.56
1:C:268:ILE:HG23	1:C:442:ILE:HD11	1.87	0.55
1:C:310:ILE:HA	1:C:313:PHE:CE2	2.42	0.54
1:B:310:ILE:HA	1:B:313:PHE:CE2	2.43	0.54
1:D:434:GLU:HG3	1:D:435:HIS:CD2	2.45	0.52
1:D:310:ILE:HA	1:D:313:PHE:CE2	2.45	0.51
1:A:310:ILE:HA	1:A:313:PHE:CE2	2.46	0.50
1:A:320:VAL:HG21	1:A:325:LEU:HB2	1.93	0.50
1:C:348:ARG:HH21	1:C:431:LYS:HG3	1.77	0.49
1:A:394:GLU:HA	1:A:397:TYR:CE2	2.48	0.48
1:D:320:VAL:HG21	1:D:325:LEU:HB2	1.96	0.48
1:D:348:ARG:HD3	3:D:504:BR:BR	2.69	0.48
1:B:371:ARG:HD3	3:B:502:BR:BR	2.69	0.47
1:C:315:HIS:HE1	1:C:364:LYS:HD3	1.80	0.47
1:C:323:GLY:HA3	1:C:331:HIS:HE1	1.81	0.46
1:B:394:GLU:HA	1:B:397:TYR:CE2	2.52	0.45
1:C:394:GLU:HA	1:C:397:TYR:CE2	2.52	0.44
1:D:394:GLU:HA	1:D:397:TYR:CE2	2.52	0.43
1:A:385:ASN:ND2	1:A:388:GLU:HB2	2.33	0.43
1:B:262:ASN:HB3	1:B:263:ASP:H	1.65	0.42
1:B:457:ALA:HB1	1:B:458:PRO:HD2	2.01	0.42
1:B:434:GLU:HG2	1:B:435:HIS:HD2	1.86	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:321:LYS:HE3	1:C:321:LYS:HB2	1.97	0.41
1:D:441:LEU:HD23	1:D:441:LEU:HA	1.84	0.41
1:D:385:ASN:ND2	1:D:388:GLU:HB2	2.36	0.41
1:C:431:LYS:NZ	4:C:602:HOH:O	2.48	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/243 (81%)	193 (98%)	3 (2%)	0	100	100
1	B	196/243 (81%)	191 (97%)	5 (3%)	0	100	100
1	C	195/243 (80%)	192 (98%)	3 (2%)	0	100	100
1	D	194/243 (80%)	191 (98%)	3 (2%)	0	100	100
All	All	781/972 (80%)	767 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/209 (81%)	170 (100%)	0	100	100
1	B	170/209 (81%)	169 (99%)	1 (1%)	78	91

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	169/209 (81%)	165 (98%)	4 (2%)	43	70
1	D	168/209 (80%)	167 (99%)	1 (1%)	78	91
All	All	677/836 (81%)	671 (99%)	6 (1%)	70	87

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	442	ILE
1	C	263	ASP
1	C	321	LYS
1	C	442	ILE
1	C	445	THR
1	D	438	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	335	ASN
1	A	385	ASN
1	A	406	HIS
1	A	460	GLN
1	B	333	HIS
1	B	385	ASN
1	C	275	GLN
1	C	288	HIS
1	C	331	HIS
1	C	335	ASN
1	C	385	ASN
1	D	331	HIS
1	D	385	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 11 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	9HF	A	501	-	30,30,30	4.44	12 (40%)	47,48,48	1.68	10 (21%)
2	9HF	C	501	-	30,30,30	4.20	12 (40%)	47,48,48	2.25	12 (25%)
2	9HF	D	501	-	30,30,30	4.47	12 (40%)	47,48,48	1.69	12 (25%)
2	9HF	D	502	-	30,30,30	4.41	12 (40%)	47,48,48	1.85	11 (23%)
2	9HF	B	501	-	30,30,30	4.28	12 (40%)	47,48,48	2.30	13 (27%)
2	9HF	A	502	-	30,30,30	4.34	12 (40%)	47,48,48	1.95	10 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9HF	A	501	-	-	0/8/27/27	0/4/4/4
2	9HF	C	501	-	-	1/8/27/27	0/4/4/4
2	9HF	D	501	-	-	0/8/27/27	0/4/4/4
2	9HF	D	502	-	-	4/8/27/27	0/4/4/4
2	9HF	B	501	-	-	0/8/27/27	0/4/4/4
2	9HF	A	502	-	-	4/8/27/27	0/4/4/4

All (72) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	501	9HF	NAL-NAM	-15.00	1.22	1.38
2	A	501	9HF	NAL-NAM	-14.78	1.23	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	502	9HF	NAL-NAM	-14.74	1.23	1.38
2	A	502	9HF	NAL-NAM	-14.10	1.23	1.38
2	B	501	9HF	NAL-NAM	-12.78	1.25	1.38
2	C	501	9HF	NAL-NAM	-11.83	1.26	1.38
2	D	501	9HF	CAD-CAE	-10.92	1.39	1.53
2	B	501	9HF	CAP-CAI	-10.51	1.39	1.53
2	A	501	9HF	CAD-CAE	-10.42	1.39	1.53
2	C	501	9HF	CAP-CAI	-10.14	1.40	1.53
2	A	502	9HF	CAP-CAI	-10.02	1.40	1.53
2	D	502	9HF	CAD-CAE	-9.78	1.40	1.53
2	D	502	9HF	CAP-CAI	-9.75	1.40	1.53
2	C	501	9HF	CAD-CAE	-9.74	1.40	1.53
2	A	502	9HF	CAD-CAE	-9.52	1.40	1.53
2	A	501	9HF	CAP-CAI	-9.47	1.41	1.53
2	B	501	9HF	CAD-CAE	-9.39	1.41	1.53
2	D	501	9HF	CAP-CAI	-9.31	1.41	1.53
2	C	501	9HF	NAN-NAM	-6.38	1.21	1.30
2	B	501	9HF	NAN-NAM	-6.37	1.21	1.30
2	B	501	9HF	CAB-CAG	-6.20	1.39	1.51
2	C	501	9HF	CAB-CAG	-6.17	1.39	1.51
2	A	501	9HF	CAB-CAG	-6.15	1.39	1.51
2	D	502	9HF	CAB-CAG	-5.99	1.39	1.51
2	A	502	9HF	CAB-CAG	-5.98	1.39	1.51
2	D	501	9HF	CAB-CAG	-5.93	1.39	1.51
2	D	501	9HF	CAX-CAY	-4.78	1.39	1.49
2	A	501	9HF	CAX-CAY	-4.75	1.39	1.49
2	A	502	9HF	CAX-CAY	-4.69	1.39	1.49
2	D	502	9HF	CAX-CAY	-4.67	1.39	1.49
2	A	501	9HF	CAR-CAS	-4.66	1.32	1.40
2	C	501	9HF	CAX-CAY	-4.65	1.39	1.49
2	D	501	9HF	CAR-CAS	-4.64	1.32	1.40
2	A	502	9HF	CAR-CAS	-4.60	1.32	1.40
2	D	502	9HF	CAR-CAS	-4.59	1.32	1.40
2	B	501	9HF	CAX-CAY	-4.58	1.39	1.49
2	D	502	9HF	CAK-NAL	-4.41	1.33	1.43
2	D	501	9HF	CAK-NAL	-4.37	1.33	1.43
2	A	502	9HF	CAK-NAL	-4.34	1.34	1.43
2	B	501	9HF	CAR-CAS	-4.33	1.33	1.40
2	C	501	9HF	CAR-CAS	-4.26	1.33	1.40
2	A	501	9HF	CAK-NAL	-4.25	1.34	1.43
2	D	502	9HF	NAN-NAM	-4.20	1.24	1.30
2	C	501	9HF	CAV-CAS	-4.15	1.33	1.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	502	9HF	NAN-NAM	-4.13	1.24	1.30
2	A	502	9HF	CAV-CAS	-4.13	1.33	1.40
2	D	502	9HF	CAV-CAS	-4.07	1.33	1.40
2	A	501	9HF	CAV-CAS	-4.05	1.33	1.40
2	A	501	9HF	NAN-NAM	-4.00	1.24	1.30
2	B	501	9HF	CAV-CAS	-3.99	1.33	1.40
2	D	501	9HF	CAV-CAS	-3.97	1.33	1.40
2	D	501	9HF	CAR-NAL	-3.86	1.33	1.38
2	C	501	9HF	CAK-NAL	-3.83	1.35	1.43
2	B	501	9HF	CAK-NAL	-3.77	1.35	1.43
2	D	502	9HF	CAU-CAR	-3.76	1.33	1.39
2	D	502	9HF	CAR-NAL	-3.74	1.33	1.38
2	D	501	9HF	CAU-CAR	-3.73	1.33	1.39
2	A	502	9HF	CAU-CAR	-3.70	1.33	1.39
2	A	501	9HF	CAU-CAR	-3.69	1.33	1.39
2	A	502	9HF	CAR-NAL	-3.67	1.33	1.38
2	C	501	9HF	CAU-CAR	-3.61	1.34	1.39
2	B	501	9HF	CAU-CAR	-3.57	1.34	1.39
2	A	501	9HF	CAR-NAL	-3.56	1.33	1.38
2	B	501	9HF	CAS-NAN	-3.37	1.33	1.38
2	C	501	9HF	CAS-NAN	-3.36	1.33	1.38
2	D	502	9HF	CAS-NAN	-3.23	1.33	1.38
2	A	502	9HF	CAS-NAN	-3.21	1.33	1.38
2	A	501	9HF	CAS-NAN	-3.20	1.33	1.38
2	D	501	9HF	CAS-NAN	-3.12	1.33	1.38
2	D	501	9HF	NAN-NAM	-3.11	1.26	1.30
2	C	501	9HF	CAR-NAL	-3.04	1.34	1.38
2	B	501	9HF	CAR-NAL	-2.81	1.34	1.38

All (68) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	9HF	CAG-CAK-NAL	8.05	125.09	119.57
2	C	501	9HF	CAG-CAK-NAL	7.74	124.88	119.57
2	A	502	9HF	CAG-CAK-NAL	7.56	124.76	119.57
2	B	501	9HF	NAL-NAM-NAN	6.78	112.56	108.77
2	C	501	9HF	NAL-NAM-NAN	6.75	112.54	108.77
2	D	502	9HF	CAG-CAK-NAL	6.45	123.99	119.57
2	C	501	9HF	CAR-NAL-NAM	-6.41	107.03	109.79
2	B	501	9HF	CAR-NAL-NAM	-6.06	107.18	109.79
2	A	501	9HF	NAL-NAM-NAN	4.84	111.47	108.77
2	D	502	9HF	NAL-NAM-NAN	4.66	111.37	108.77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	501	9HF	CAK-NAL-NAM	4.63	127.73	120.11
2	A	502	9HF	NAL-NAM-NAN	4.61	111.34	108.77
2	D	501	9HF	CAR-CAS-NAN	-4.55	104.91	108.70
2	D	502	9HF	CAR-CAS-NAN	-4.38	105.06	108.70
2	A	501	9HF	CAR-CAS-NAN	-4.31	105.12	108.70
2	A	502	9HF	CAR-CAS-NAN	-4.09	105.30	108.70
2	D	502	9HF	CAK-NAL-NAM	3.90	126.52	120.11
2	D	501	9HF	CAS-CAR-NAL	3.88	106.44	103.97
2	B	501	9HF	CAR-CAS-NAN	-3.86	105.49	108.70
2	A	502	9HF	CAK-NAL-NAM	3.84	126.42	120.11
2	D	501	9HF	NAL-NAM-NAN	3.84	110.91	108.77
2	B	501	9HF	CAJ-CAK-CAG	-3.75	117.55	121.79
2	A	501	9HF	CAK-NAL-NAM	3.75	126.27	120.11
2	C	501	9HF	CAJ-CAK-CAG	-3.62	117.71	121.79
2	C	501	9HF	CAR-CAS-NAN	-3.53	105.76	108.70
2	A	501	9HF	CAS-CAR-NAL	3.34	106.10	103.97
2	A	502	9HF	CAS-CAR-NAL	3.20	106.01	103.97
2	A	502	9HF	CAR-NAL-NAM	-3.19	108.42	109.79
2	D	502	9HF	CAS-CAR-NAL	3.18	106.00	103.97
2	A	501	9HF	CAR-NAL-NAM	-3.17	108.43	109.79
2	D	502	9HF	CAR-NAL-NAM	-2.87	108.56	109.79
2	D	501	9HF	CAU-CAW-CAX	-2.75	117.86	120.80
2	A	502	9HF	CAU-CAW-CAX	-2.66	117.96	120.80
2	D	501	9HF	CAC-CAD-CAE	-2.65	105.55	110.12
2	A	502	9HF	CAJ-CAK-CAG	-2.61	118.84	121.79
2	A	501	9HF	CAU-CAW-CAX	-2.58	118.04	120.80
2	A	502	9HF	CAJ-CAK-NAL	-2.58	116.38	118.91
2	A	501	9HF	CAJ-CAK-CAG	-2.51	118.95	121.79
2	B	501	9HF	CAF-CAG-CAK	2.51	120.04	117.17
2	B	501	9HF	CAH-CAO-CAP	-2.43	104.93	113.84
2	C	501	9HF	CAF-CAG-CAK	2.40	119.91	117.17
2	C	501	9HF	CAU-CAW-CAX	-2.37	118.27	120.80
2	C	501	9HF	CAS-CAR-NAL	2.36	105.48	103.97
2	D	502	9HF	CAJ-CAK-CAG	-2.36	119.12	121.79
2	A	501	9HF	CAG-CAK-NAL	2.36	121.19	119.57
2	A	501	9HF	CAF-CAG-CAK	2.35	119.86	117.17
2	D	502	9HF	CAU-CAW-CAX	-2.31	118.33	120.80
2	B	501	9HF	CAS-CAR-NAL	2.28	105.42	103.97
2	B	501	9HF	CAU-CAW-CAX	-2.26	118.38	120.80
2	B	501	9HF	CAV-CAS-CAR	2.25	122.23	120.16
2	C	501	9HF	CAH-CAO-CAP	-2.24	105.61	113.84
2	C	501	9HF	CAO-CAH-CAD	-2.21	105.73	113.84

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	501	9HF	CAR-NAL-NAM	-2.20	108.84	109.79
2	D	501	9HF	CAG-CAK-NAL	2.17	121.06	119.57
2	B	501	9HF	CAT-CAP-CAI	-2.17	106.38	110.12
2	B	501	9HF	CAB-CAG-CAF	-2.15	115.79	119.57
2	A	501	9HF	CAO-CAH-CAD	-2.14	105.97	113.84
2	A	502	9HF	CAH-CAO-CAP	-2.14	105.98	113.84
2	C	501	9HF	CAV-CAS-CAR	2.10	122.10	120.16
2	C	501	9HF	CAB-CAG-CAF	-2.10	115.87	119.57
2	D	501	9HF	OAZ-CAY-CAX	2.09	120.20	114.84
2	D	502	9HF	OAZ-CAY-CAX	2.09	120.19	114.84
2	D	502	9HF	CAJ-CAK-NAL	-2.08	116.87	118.91
2	D	501	9HF	CAJ-CAK-CAG	-2.08	119.44	121.79
2	D	502	9HF	CAH-CAO-CAP	-2.07	106.25	113.84
2	B	501	9HF	OAZ-CAY-CAX	2.06	120.14	114.84
2	D	501	9HF	CAH-CAO-CAP	-2.04	106.36	113.84
2	D	501	9HF	CAO-CAH-CAD	-2.01	106.48	113.84

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	502	9HF	CAW-CAX-CAY-OAZ
2	D	502	9HF	CAV-CAX-CAY-OAZ
2	A	502	9HF	CAW-CAX-CAY-OAZ
2	D	502	9HF	CAW-CAX-CAY-OBA
2	A	502	9HF	CAV-CAX-CAY-OAZ
2	D	502	9HF	CAV-CAX-CAY-OBA
2	A	502	9HF	CAW-CAX-CAY-OBA
2	A	502	9HF	CAV-CAX-CAY-OBA
2	C	501	9HF	CAJ-CAK-NAL-CAR

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the


average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




## Ligand 9HF D 501



## Ligand 9HF D 502





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	198/243 (81%)	0.32	7 (3%) 47 41	36, 54, 87, 121	0
1	B	198/243 (81%)	0.36	10 (5%) 33 28	33, 57, 91, 123	0
1	C	197/243 (81%)	0.56	10 (5%) 33 28	46, 64, 99, 114	0
1	D	196/243 (80%)	0.57	12 (6%) 27 22	42, 64, 93, 149	0
All	All	789/972 (81%)	0.45	39 (4%) 35 29	33, 60, 95, 149	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	325	LEU	4.4
1	D	264	PRO	4.3
1	D	265	VAL	3.9
1	B	261	PRO	3.7
1	D	266	THR	3.5
1	C	322	ASP	3.5
1	D	438	PHE	3.4
1	A	265	VAL	3.4
1	A	266	THR	3.2
1	B	458	PRO	3.2
1	D	294	LEU	3.1
1	A	439	PHE	3.1
1	A	438	PHE	3.0
1	B	331	HIS	3.0
1	C	346	PHE	3.0
1	B	324	ILE	2.9
1	B	325	LEU	2.9
1	A	264	PRO	2.8
1	C	326	LEU	2.6
1	C	333	HIS	2.6
1	C	262	ASN	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	320	VAL	2.5
1	D	436	LEU	2.4
1	B	330	LEU	2.4
1	C	331	HIS	2.4
1	A	263	ASP	2.4
1	C	328	THR	2.4
1	B	321	LYS	2.4
1	A	436	LEU	2.3
1	C	330	LEU	2.3
1	D	439	PHE	2.2
1	B	326	LEU	2.2
1	B	345	ILE	2.2
1	D	437	PHE	2.2
1	D	333	HIS	2.2
1	D	267	ASN	2.1
1	D	459	HIS	2.1
1	D	285	ARG	2.1
1	B	335	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	9HF	A	501	27/27	0.81	0.15	59,68,77,85	0
2	9HF	D	501	27/27	0.86	0.14	69,73,78,82	0
2	9HF	C	501	27/27	0.87	0.12	63,70,81,97	0
2	9HF	B	501	27/27	0.87	0.12	64,68,75,82	0
2	9HF	A	502	27/27	0.88	0.17	63,66,69,70	27

*Continued on next page...*




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	9HF	D	502	27/27	0.91	0.16	65,74,83,86	27
3	BR	B	504	1/1	0.95	0.08	77,77,77,77	1
3	BR	C	503	1/1	0.95	0.07	73,73,73,73	1
3	BR	B	503	1/1	0.96	0.06	69,69,69,69	1
3	BR	A	506	1/1	0.97	0.09	74,74,74,74	1
3	BR	A	504	1/1	0.98	0.04	46,46,46,46	1
3	BR	A	505	1/1	0.98	0.05	69,69,69,69	1
3	BR	C	502	1/1	0.98	0.04	55,55,55,55	1
3	BR	A	503	1/1	0.98	0.05	45,45,45,45	1
3	BR	D	503	1/1	0.98	0.07	60,60,60,60	1
3	BR	D	504	1/1	0.98	0.04	76,76,76,76	1
3	BR	B	502	1/1	0.99	0.04	62,62,62,62	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 9HF A 501:**




2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around 9HF D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 9HF C 501:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around 9HF B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 9HF A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 9HF D 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.