



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:42 PM UTC

PDB ID : 6N6L / pdb\_00006n6l  
Title : Crystal Structure of ATPase delta 1-79 Spa47 R189A R191A mutant  
Authors : Morales, Y.; Johnson, S.J.; Demler, H.J.; Dickenson, N.E.  
Deposited on : 2018-11-26  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	<div> <div>6%</div> <div> <div></div> <div>87%</div> <div>12%</div> <div>.</div> </div> </div>
1	B	352	<div> <div>10%</div> <div> <div></div> <div>84%</div> <div>15%</div> <div>.</div> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5667 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP synthase SpaL/MxiB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	348	Total	C	N	O	S	0	1	0
			2716	1727	458	522	9			
1	B	348	Total	C	N	O	S	0	1	0
			2715	1724	461	521	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	79	ASN	-	expression tag	UNP P0A1C1
A	189	ALA	ARG	engineered mutation	UNP P0A1C1
A	191	ALA	ARG	engineered mutation	UNP P0A1C1
B	79	ASN	-	expression tag	UNP P0A1C1
B	189	ALA	ARG	engineered mutation	UNP P0A1C1
B	191	ALA	ARG	engineered mutation	UNP P0A1C1

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	5	0
			5	4	1		
2	B	1	Total	O	S	5	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	130	Total	O	0	0
			130	130		
3	B	96	Total	O	0	0
			96	96		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ATP synthase SpaL/MxiB



#### • Molecule 1: ATP synthase SpaL/MxiB



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.53Å 152.92Å 54.73Å 90.00° 110.90° 90.00°	Depositor
Resolution (Å)	35.84 – 2.15 35.84 – 2.15	Depositor EDS
% Data completeness (in resolution range)	89.8 (35.84-2.15) 83.3 (35.84-2.15)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.49 (at 2.14Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, $R_{free}$	0.186 , 0.227 0.187 , 0.226	Depositor DCC
$R_{free}$ test set	1763 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.5	Xtriage
Anisotropy	0.450	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 44.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5667	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.19	0/2767	0.43	0/3742
1	B	0.15	0/2765	0.39	0/3738
All	All	0.17	0/5532	0.41	0/7480

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	186	ILE	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2716	0	2720	29	0
1	B	2715	0	2724	38	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	130	0	0	8	0
3	B	96	0	0	6	0
All	All	5667	0	5444	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:314:PRO:O	1:B:318:GLU:OE2	1.59	1.19
1:B:151:GLU:OE1	3:B:601:HOH:O	1.86	0.91
1:B:314:PRO:C	1:B:318:GLU:OE2	2.14	0.90
1:A:223:ARG:NH2	3:A:602:HOH:O	2.11	0.83
1:B:83:LEU:HD23	1:B:84:HIS:ND1	1.94	0.82
1:A:83:LEU:N	3:A:604:HOH:O	2.16	0.78
1:B:398:TYR:O	3:B:603:HOH:O	2.09	0.70
1:B:111:ASN:O	3:B:602:HOH:O	2.09	0.70
1:B:153:GLN:HG3	1:B:324:ASP:HB2	1.74	0.70
1:B:315:LEU:HA	1:B:318:GLU:CD	2.22	0.63
1:B:369:GLU:OE2	3:B:604:HOH:O	2.15	0.62
1:B:312:ALA:O	3:B:606:HOH:O	2.16	0.61
1:B:358:GLU:OE1	1:B:361[B]:ARG:NH1	2.34	0.60
1:B:318:GLU:OE1	3:B:605:HOH:O	2.15	0.60
1:A:223:ARG:HH22	1:A:256:ARG:HH22	1.52	0.57
1:A:387:LYS:HE3	1:A:390:GLU:OE1	2.05	0.57
1:A:427:GLU:OE2	1:A:430:ARG:HD2	2.05	0.57
1:A:326:HIS:ND1	3:A:603:HOH:O	2.16	0.56
1:A:313:ASP:HB2	1:A:314:PRO:HD2	1.86	0.56
1:B:425:ILE:O	1:B:428:THR:HG22	2.06	0.56
1:A:153:GLN:HG3	1:A:324:ASP:HB2	1.88	0.55
1:A:335:GLN:HG3	1:A:336:LYS:N	2.20	0.55
1:B:400:LYS:NZ	1:B:428:THR:O	2.40	0.55
1:B:154:ARG:HD2	1:B:286:LEU:O	2.07	0.55
1:A:129:ARG:NH2	1:A:287:GLU:O	2.29	0.55
1:A:223:ARG:HH22	1:A:256:ARG:NH2	2.05	0.54
1:B:238:ARG:HD2	1:B:292:LEU:HG	1.92	0.52
1:A:332:ASN:OD1	1:A:336:LYS:NZ	2.43	0.51

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:188:GLU:HG3	1:B:192:GLU:OE1	2.09	0.51
1:A:403:VAL:HG13	3:A:627:HOH:O	2.09	0.51
1:B:187:GLY:N	1:B:214:ALA:O	2.45	0.49
1:A:335:GLN:HE21	1:A:336:LYS:CG	2.26	0.49
1:B:116:TYR:O	1:B:117:ARG:NH1	2.44	0.49
1:B:367:PHE:CD1	1:B:428:THR:HG21	2.48	0.48
1:A:88:GLY:HA2	1:A:236:PHE:CE2	2.49	0.47
1:B:83:LEU:CD2	1:B:84:HIS:ND1	2.72	0.47
1:B:358:GLU:OE2	1:B:361[A]:ARG:NH2	2.46	0.47
1:B:345:LEU:O	1:B:346:LYS:HD3	2.16	0.46
1:B:194:THR:HA	1:B:197:VAL:HG12	1.96	0.46
1:B:345:LEU:C	1:B:346:LYS:HD3	2.40	0.46
1:B:249:ASP:HA	1:B:250:SER:HA	1.60	0.46
1:A:249:ASP:HA	1:A:250:SER:HA	1.68	0.45
1:A:396:LYS:O	1:A:396:LYS:HD3	2.17	0.45
1:A:181:TYR:O	1:A:209:CYS:HA	2.17	0.45
1:A:367:PHE:HE1	1:A:428:THR:HG21	1.80	0.45
1:B:235:GLU:O	1:B:239:THR:HG23	2.17	0.45
1:B:282:LEU:HB3	1:B:283:PRO:HD3	1.98	0.44
1:A:130:ALA:HB3	1:A:291:LYS:HG3	1.99	0.44
1:B:185:LEU:HG	1:B:211:LEU:HD11	2.00	0.44
1:A:282:LEU:HB3	1:A:283:PRO:HD3	1.99	0.44
1:A:380:ILE:HG23	1:A:385:GLU:HB2	1.99	0.44
1:B:115:LEU:HD13	1:B:236:PHE:CZ	2.53	0.44
1:A:312:ALA:HB1	1:A:316:ALA:HB3	2.00	0.43
1:A:335:GLN:HE21	1:A:336:LYS:HG3	1.82	0.43
1:B:163:CYS:SG	1:B:331:ARG:HB2	2.58	0.43
1:B:274:TYR:CG	1:B:275:PRO:HA	2.54	0.43
1:A:154:ARG:HD3	3:A:666:HOH:O	2.18	0.43
1:A:256:ARG:CZ	3:A:602:HOH:O	2.67	0.42
1:B:93:GLY:HA2	1:B:106:PHE:CE1	2.54	0.42
1:A:401:ILE:O	1:A:405:GLU:HG3	2.20	0.42
1:B:282:LEU:O	1:B:286:LEU:HG	2.20	0.42
1:A:288:ARG:NH2	3:A:608:HOH:O	2.33	0.41
1:B:165:LYS:HE2	1:B:165:LYS:HB3	1.89	0.41
1:B:193:VAL:O	1:B:197:VAL:HG12	2.20	0.41
1:B:194:THR:HA	1:B:197:VAL:CG1	2.51	0.41
1:B:180:ILE:HG21	1:B:237:PHE:CD2	2.57	0.40
1:A:368:ARG:HD2	3:A:691:HOH:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/352 (99%)	339 (98%)	7 (2%)	1 (0%)	36	34
1	B	347/352 (99%)	336 (97%)	11 (3%)	0	100	100
All	All	694/704 (99%)	675 (97%)	18 (3%)	1 (0%)	48	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	308	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/292 (99%)	290 (100%)	0	100	100
1	B	290/292 (99%)	290 (100%)	0	100	100
All	All	580/584 (99%)	580 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	GLN
1	A	121	ASN
1	A	202	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	335	GLN
1	A	338	GLN
1	B	202	ASN
1	B	242	HIS
1	B	335	GLN
1	B	354	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	501	-	4,4,4	0.23	0	6,6,6	0.08	0
2	SO4	B	501	-	4,4,4	0.24	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	348/352 (98%)	0.36	21 (6%)	27	31	24, 38, 76, 101	1 (0%)
1	B	348/352 (98%)	0.65	35 (10%)	12	14	26, 47, 83, 115	1 (0%)
All	All	696/704 (98%)	0.51	56 (8%)	18	20	24, 43, 80, 115	2 (0%)

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	83	LEU	5.1
1	A	311	PHE	4.7
1	B	99	LEU	4.5
1	A	122	ALA	4.3
1	B	312	ALA	4.0
1	A	270	ALA	3.6
1	B	83	LEU	3.6
1	B	311	PHE	3.5
1	A	310	ASP	3.5
1	B	98	PRO	3.4
1	A	187	GLY	3.2
1	B	276	VAL	3.1
1	A	126[A]	TYR	3.0
1	A	123	PRO	3.0
1	B	271	ARG	3.0
1	B	309	ASP	2.9
1	A	269	PRO	2.9
1	A	312	ALA	2.9
1	B	314	PRO	2.8
1	B	313	ASP	2.8
1	B	429	ILE	2.8
1	A	314	PRO	2.8
1	B	270	ALA	2.7
1	B	274	TYR	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	310	ASP	2.7
1	B	363	MET	2.6
1	B	269	PRO	2.6
1	B	338	GLN	2.6
1	B	315	LEU	2.6
1	A	335	GLN	2.5
1	B	268	SER	2.5
1	B	280	ASP	2.4
1	B	199	TYR	2.4
1	A	430	ARG	2.4
1	A	276	VAL	2.3
1	A	188	GLU	2.3
1	B	197	VAL	2.3
1	B	126	TYR	2.3
1	B	275	PRO	2.3
1	A	189	ALA	2.3
1	B	279	PHE	2.2
1	B	114	ILE	2.2
1	B	108	VAL	2.2
1	A	169	MET	2.2
1	A	313	ASP	2.2
1	A	427	GLU	2.2
1	B	101	GLU	2.2
1	B	187	GLY	2.2
1	B	189	ALA	2.1
1	B	169	MET	2.1
1	A	306	LEU	2.1
1	B	308	ASP	2.1
1	B	278	VAL	2.1
1	A	125	LEU	2.1
1	B	345	LEU	2.1
1	B	188	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	501	5/5	-	-	50,50,50,50	5
2	SO4	B	501	5/5	-	-	52,52,52,52	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.