



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:20 PM UTC

PDB ID : 6ZAA / pdb\_00006zaa  
Title : PI3K Delta in complex with methoxy(methylsulfamoyl)pyridinylN(methylpiperidinyl)dihydrobenzoxazinecarboxamide  
Authors : Convery, M.A.; Hardy, C.J.; Spencer, J.A.; Rowland, P.  
Deposited on : 2020-06-05  
Resolution : 2.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	940	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7061 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit delta isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	859	Total	C	N	O	S	0	0	0
			6913	4418	1184	1257	54			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	105	GLY	-	expression tag	UNP O35904
A	508	GLN	-	insertion	UNP O35904

- Molecule 2 is 4-[6-methoxy-5-(methylsulfamoyl)pyridin-3-yl]- {N}-(1-methylpiperidin-4-yl)-2,3-dihydro-1,4-benzoxazine-6-carboxamide (CCD ID: QD2) (formula: C<sub>22</sub>H<sub>29</sub>N<sub>5</sub>O<sub>5</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			33	22	5	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	115	Total 115	O 115	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	140.05Å 64.56Å 115.46Å 90.00° 101.99° 90.00°	Depositor
Resolution (Å)	68.50 – 2.52 68.50 – 2.52	Depositor EDS
% Data completeness (in resolution range)	77.3 (68.50-2.52) 70.4 (68.50-2.52)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.63 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, $R_{free}$	0.184 , 0.259 (Not available) , 0.281	Depositor DCC
$R_{free}$ test set	1274 reflections (3.72%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	74.7	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 78.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7061	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	96.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QD2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.88	0/7064	1.40	37/9536 (0.4%)

There are no bond length outliers.

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	915	PHE	CA-CB-CG	7.13	120.93	113.80
1	A	745	CYS	CA-C-N	6.51	129.00	120.60
1	A	745	CYS	C-N-CA	6.51	129.00	120.60
1	A	1028	LYS	CA-C-N	6.07	128.42	120.28
1	A	1028	LYS	C-N-CA	6.07	128.42	120.28
1	A	132	ARG	CA-C-N	6.02	128.84	120.65
1	A	132	ARG	C-N-CA	6.02	128.84	120.65
1	A	1000	ASP	CA-CB-CG	6.02	118.62	112.60
1	A	937	ASP	CA-CB-CG	5.87	118.47	112.60
1	A	918	ASN	CA-CB-CG	5.44	118.04	112.60
1	A	629	LEU	CA-C-N	5.43	127.50	120.44
1	A	629	LEU	C-N-CA	5.43	127.50	120.44
1	A	422	ASN	CA-CB-CG	5.38	117.98	112.60
1	A	958	GLY	CA-C-N	5.37	127.47	120.28
1	A	958	GLY	C-N-CA	5.37	127.47	120.28
1	A	874	GLU	CA-C-N	5.36	127.90	120.29
1	A	874	GLU	C-N-CA	5.36	127.90	120.29
1	A	631	LYS	CA-C-N	5.23	127.24	120.44
1	A	631	LYS	C-N-CA	5.23	127.24	120.44
1	A	873	GLU	CB-CG-CD	5.23	121.49	112.60
1	A	933	ILE	CB-CA-C	5.23	117.64	111.32
1	A	1022	ALA	CA-C-N	5.20	127.67	120.29
1	A	1022	ALA	C-N-CA	5.20	127.67	120.29
1	A	982	ARG	CA-C-N	5.20	127.97	120.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	982	ARG	C-N-CA	5.20	127.97	120.38
1	A	113	ILE	CA-C-N	5.17	127.21	120.28
1	A	113	ILE	C-N-CA	5.17	127.21	120.28
1	A	646	PHE	CA-CB-CG	-5.17	108.63	113.80
1	A	898	ASN	CA-CB-CG	5.13	117.73	112.60
1	A	334	ASN	CA-CB-CG	5.12	117.72	112.60
1	A	687	GLU	CB-CG-CD	5.11	121.29	112.60
1	A	593	GLY	CA-C-N	5.11	127.13	120.28
1	A	593	GLY	C-N-CA	5.11	127.13	120.28
1	A	115	SER	CA-C-N	5.09	127.11	120.28
1	A	115	SER	C-N-CA	5.09	127.11	120.28
1	A	243	VAL	N-CA-C	-5.09	102.17	108.89
1	A	203	PHE	CA-CB-CG	5.08	118.88	113.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6913	0	6889	56	0
2	A	33	0	0	1	0
3	A	115	0	0	3	0
All	All	7061	0	6889	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:752:MET:HB2	1:A:758:PRO:HD2	1.65	0.79
1:A:383:ASP:HB3	1:A:556:ASN:O	1.99	0.62
1:A:419:ALA:HB1	1:A:441:MET:HB3	1.82	0.60
1:A:830:HIS:O	1:A:903:GLU:HG3	2.01	0.60

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:875:PHE:O	1:A:879:CYS:HB2	2.02	0.59
1:A:328:ILE:HD11	1:A:474:TYR:HB2	1.86	0.58
1:A:208:SER:OG	1:A:210:LYS:HG2	2.05	0.57
1:A:975:LEU:HD23	1:A:998:LEU:HD23	1.87	0.57
1:A:971:GLY:HA3	1:A:1004:LEU:HD21	1.86	0.56
1:A:279:SER:O	1:A:283:LEU:HD12	2.06	0.56
1:A:157:LEU:HD22	1:A:161:GLU:HB3	1.86	0.56
1:A:597:ILE:HD11	1:A:629:LEU:HD13	1.87	0.55
1:A:834:ILE:HD12	1:A:855:LEU:HD11	1.89	0.53
1:A:191:LEU:O	1:A:272:PRO:HD2	2.09	0.52
1:A:189:ALA:HB1	1:A:206:GLN:HE22	1.75	0.51
1:A:194:VAL:HG21	1:A:216:LEU:CD2	2.39	0.51
1:A:535:ARG:NH2	1:A:563:GLN:HE21	2.09	0.50
1:A:349:HIS:C	1:A:588:PRO:HG3	2.38	0.49
1:A:890:GLY:HA3	1:A:917:GLY:H	1.78	0.49
1:A:452:LEU:HA	3:A:4210:HOH:O	2.13	0.48
1:A:194:VAL:HG21	1:A:216:LEU:HD21	1.95	0.48
1:A:765:SER:HB3	1:A:768:ALA:HB3	1.95	0.47
1:A:247:HIS:CE1	1:A:740:LEU:HD21	2.49	0.47
1:A:245:GLY:HA3	1:A:768:ALA:HB2	1.96	0.47
1:A:216:LEU:HD12	1:A:256:LEU:HD11	1.97	0.47
1:A:617:GLN:HB3	1:A:983:ALA:HB3	1.97	0.46
1:A:441:MET:HB2	1:A:469:ALA:O	2.16	0.46
1:A:784:LEU:HD12	1:A:823:GLY:HA3	1.96	0.46
1:A:859:LEU:HD21	1:A:901:ILE:HD11	1.98	0.46
1:A:534:MET:O	1:A:538:VAL:HG23	2.15	0.46
1:A:964:TYR:CE2	1:A:968:ARG:HD2	2.50	0.46
1:A:860:LYS:HG2	1:A:868:LEU:HD22	1.97	0.46
1:A:153:HIS:HB2	3:A:4161:HOH:O	2.16	0.46
1:A:263:CYS:HA	1:A:266:LEU:HD12	1.98	0.45
1:A:341:LEU:HG	1:A:365:VAL:HG22	1.98	0.45
1:A:995:ILE:HG22	1:A:999:LYS:HE2	1.98	0.45
1:A:609:PHE:HE1	1:A:646:PHE:CD2	2.35	0.44
1:A:617:GLN:HG3	1:A:984:ALA:HB2	2.00	0.44
1:A:321:GLN:O	1:A:381:VAL:HG23	2.18	0.43
1:A:390:LEU:HB2	1:A:425:LEU:HD21	1.99	0.43
1:A:833:THR:HG23	1:A:900:MET:HE3	1.99	0.43
1:A:616:VAL:O	1:A:619:LEU:HB2	2.18	0.43
1:A:289:GLN:HG2	1:A:677:HIS:CE1	2.54	0.42
1:A:940:HIS:CE1	1:A:945:GLY:HA2	2.55	0.42
1:A:832:ASP:C	1:A:900:MET:HE2	2.45	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:777:ILE:O	1:A:824:LEU:HA	2.20	0.41
1:A:386:ARG:HE	1:A:430:ASP:CG	2.28	0.41
1:A:883:CYS:HB3	1:A:932:PHE:HD1	1.85	0.41
1:A:842:SER:C	1:A:844:MET:H	2.29	0.41
1:A:420:TRP:CE2	1:A:442:TRP:HB2	2.55	0.41
1:A:420:TRP:NE1	1:A:442:TRP:HB2	2.36	0.41
1:A:606:ASP:HB2	3:A:4112:HOH:O	2.21	0.41
1:A:859:LEU:CD2	1:A:905:GLY:HA2	2.51	0.41
2:A:4000:QD2:C2	2:A:4000:QD2:C13	2.97	0.40
1:A:645:HIS:CG	1:A:737:PRO:HG3	2.56	0.40
1:A:801:TRP:HB3	1:A:806:LEU:HB3	2.03	0.40
1:A:943:GLN:HG2	1:A:949:ASN:HB3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	845/940 (90%)	806 (95%)	38 (4%)	1 (0%)	48 67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	847	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	757/827 (92%)	713 (94%)	44 (6%)	18	36

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	VAL
1	A	115	SER
1	A	119	LEU
1	A	166	SER
1	A	174	SER
1	A	187	ASN
1	A	200	GLU
1	A	206	GLN
1	A	233	VAL
1	A	264	SER
1	A	267	HIS
1	A	283	LEU
1	A	318	SER
1	A	319	LEU
1	A	324	SER
1	A	334	ASN
1	A	357	THR
1	A	427	ASP
1	A	489	GLU
1	A	496	ARG
1	A	503	ILE
1	A	506	GLU
1	A	509	LEU
1	A	522	GLU
1	A	552	VAL
1	A	586	SER
1	A	602	LYS
1	A	676	THR
1	A	705	LYS
1	A	744	VAL
1	A	748	GLN
1	A	766	GLU
1	A	782	ASP
1	A	787	ASP
1	A	855	LEU
1	A	897	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	915	PHE
1	A	918	ASN
1	A	933	ILE
1	A	946	LYS
1	A	947	THR
1	A	975	LEU
1	A	1006	LYS
1	A	1012	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	193	ASN
1	A	206	GLN
1	A	235	GLN
1	A	247	HIS
1	A	273	HIS
1	A	334	ASN
1	A	344	GLN
1	A	351	ASN
1	A	422	ASN
1	A	462	ASN
1	A	464	ASN
1	A	510	GLN
1	A	526	HIS
1	A	563	GLN
1	A	696	ASN
1	A	721	GLN
1	A	732	GLN
1	A	943	GLN
1	A	970	HIS
1	A	976	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	QD2	A	4000	-	35,36,36	0.32	0	47,52,52	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QD2	A	4000	-	-	3/23/43/43	0/4/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	4000	QD2	C5-C1-S-O1
2	A	4000	QD2	C5-C1-S-N
2	A	4000	QD2	C-N-S-O1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	4000	QD2	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	859/940 (91%)	0.20	27 (3%)	51 48	49, 91, 154, 184	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	521	GLY	3.6
1	A	932	PHE	3.6
1	A	931	PRO	3.6
1	A	478	VAL	3.4
1	A	177	GLY	3.3
1	A	418	ILE	2.8
1	A	934	LEU	2.7
1	A	269	GLY	2.7
1	A	771	ALA	2.7
1	A	504	THR	2.5
1	A	445	VAL	2.4
1	A	933	ILE	2.4
1	A	398	VAL	2.4
1	A	452	LEU	2.3
1	A	621	TYR	2.3
1	A	989	LEU	2.2
1	A	231	PRO	2.2
1	A	323	PHE	2.2
1	A	915	PHE	2.2
1	A	1015	PHE	2.2
1	A	492	LEU	2.1
1	A	580	LEU	2.1
1	A	146	PHE	2.1
1	A	974	PHE	2.1
1	A	364	ASN	2.1
1	A	107	ASP	2.0
1	A	268	SER	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	QD2	A	4000	33/33	0.96	0.09	45,58,66,67	33

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.