



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 05:30 AM UTC

PDB ID : 7E36 / pdb_00007e36
Title : A [6+4]-cycloaddition adduct is the biosynthetic intermediate in streptoseomycin biosynthesis
Authors : Zhang, B.; Ge, H.M.
Deposited on : 2021-02-08
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>9%</div> <div>.</div> </div> </div>
2	B	421	<div> <div>8%</div> <div> <div></div> <div>81%</div> <div>11%</div> <div>6%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CL	A	406	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alkanesulfonate monooxygenase SsuD/methylene tetrahydro methanopterin reductase-like flavin-dependent oxidoreductase (Luciferase family).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2384	1477	456	450	1			

- Molecule 2 is a protein called FMN-dependent oxidoreductase (Nitrilotriacetate monooxygenase family).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	395	Total	C	N	O	S	0	0	0
			3075	1942	559	567	7			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	7	Total	Cl		0	0
			7	7			
4	B	1	Total	Cl		0	0
			1	1			

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	Mg		0	0
			1	1			

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 7 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	231	Total	O	0	0
			231	231		
8	B	260	Total	O	0	0
			260	260		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alkanesulfonate monooxygenase SsuD/methylene tetrahydromethanopterin reductase-like flavin-dependent oxidoreductase (Luciferase family)



- Molecule 2: FMN-dependent oxidoreductase (Nitrilotriacetate monooxygenase family)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.28Å 104.99Å 75.95Å 90.00° 95.95° 90.00°	Depositor
Resolution (Å)	51.00 – 2.00 51.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (51.00-2.00) 99.7 (51.00-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.165 , 0.205 0.166 , 0.206	Depositor DCC
R_{free} test set	2000 reflections (3.69%)	wwPDB-VP
Wilson B-factor (Å ²)	25.6	Xtriage
Anisotropy	0.465	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5989	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, CL, MG, MES, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/2425	0.42	0/3299
2	B	0.21	0/3152	0.48	1/4292 (0.0%)
All	All	0.20	0/5577	0.45	1/7591 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	341	THR	CB-CA-C	5.31	118.35	111.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2384	0	2361	15	0
2	B	3075	0	3000	35	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	7	0	0	4	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
6	B	12	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	8	0	12	1	0
8	A	231	0	0	2	1
8	B	260	0	0	3	1
All	All	5989	0	5385	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:134:SER:HB3	2:B:154:ARG:HD3	1.47	0.94
2:B:405:THR:HG22	2:B:408:ASP:H	1.33	0.92
4:A:407:CL:CL	8:A:721:HOH:O	2.34	0.81
1:A:148:SER:HB2	4:A:406:CL:CL	2.22	0.76
2:B:327:ALA:O	2:B:329:ARG:N	2.19	0.76
1:A:185:LEU:N	4:A:406:CL:CL	2.60	0.71
2:B:328:ARG:O	2:B:330:GLU:N	2.23	0.71
1:A:282:VAL:HG23	1:A:284:ASP:H	1.59	0.68
4:A:403:CL:CL	8:A:710:HOH:O	2.52	0.63
2:B:61:VAL:HG11	2:B:139:GLU:OE2	1.98	0.62
1:A:100:SER:O	1:A:103:ARG:HD3	2.01	0.60
2:B:329:ARG:O	2:B:331:ASN:N	2.34	0.59
1:A:55:ASP:OD1	2:B:173:ARG:NH1	2.32	0.58
2:B:273:LEU:HD21	2:B:342:ALA:HB2	1.87	0.56
1:A:297:ARG:HD2	1:A:302:PRO:HD3	1.90	0.54
2:B:105:THR:HG21	2:B:134:SER:HB2	1.88	0.54
2:B:151:HIS:HA	2:B:154:ARG:HE	1.74	0.52
2:B:151:HIS:HA	2:B:154:ARG:NE	2.25	0.52
2:B:349:THR:CG2	2:B:352:GLN:H	2.25	0.50
1:A:217:VAL:HG21	1:A:231:VAL:HG11	1.94	0.49
2:B:86:LEU:HD22	2:B:98:VAL:HB	1.95	0.49
2:B:108:ASP:OD2	8:B:601:HOH:O	2.20	0.49
2:B:405:THR:HG21	8:B:824:HOH:O	2.12	0.48
1:A:279:ASP:OD1	1:A:280:ARG:N	2.46	0.48
2:B:214:ALA:HB1	2:B:221:ILE:HD12	1.95	0.47
1:A:43:LEU:HD12	1:A:63:ALA:HB2	1.97	0.46
2:B:405:THR:HG22	2:B:408:ASP:N	2.16	0.46
1:A:180:ASP:HB2	2:B:144:GLY:HA2	1.98	0.45
2:B:273:LEU:HD21	2:B:342:ALA:CB	2.46	0.45
1:A:280:ARG:O	1:A:282:VAL:HG22	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:349:THR:OG1	2:B:350:PRO:HD2	2.16	0.45
1:A:53:ALA:N	1:A:54:PRO:HD3	2.32	0.45
2:B:307:VAL:HG22	2:B:334:LEU:HD23	1.99	0.45
1:A:278:ALA:O	1:A:281:SER:OG	2.35	0.44
2:B:203:GLN:HG2	2:B:214:ALA:HA	1.99	0.44
2:B:329:ARG:C	2:B:331:ASN:H	2.24	0.43
1:A:77:PRO:HD2	1:A:105:GLY:O	2.18	0.43
2:B:65:GLY:H	2:B:68:LEU:HG	1.84	0.43
2:B:41:ARG:HH22	7:B:503:TRS:H21	1.83	0.43
2:B:235:TYR:CZ	2:B:239:LYS:HE3	2.53	0.43
2:B:373:PHE:HA	2:B:374:PRO:HD3	1.92	0.42
2:B:293:LEU:HD22	2:B:326:LEU:HD12	2.01	0.42
2:B:8:LEU:HD12	2:B:368:GLY:HA2	2.02	0.42
2:B:327:ALA:N	8:B:620:HOH:O	2.52	0.42
2:B:338:SER:HA	2:B:341:THR:HG22	2.03	0.41
2:B:64:TRP:CE3	2:B:68:LEU:HD21	2.56	0.41
1:A:164:PHE:HZ	2:B:68:LEU:HD22	1.85	0.40
2:B:329:ARG:HE	2:B:329:ARG:HB3	1.68	0.40
2:B:402:LEU:O	2:B:404:GLY:N	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:730:HOH:O	8:B:766:HOH:O[1_554]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	317/324 (98%)	307 (97%)	8 (2%)	2 (1%)	21 17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	391/421 (93%)	370 (95%)	17 (4%)	4 (1%)	12	8
All	All	708/745 (95%)	677 (96%)	25 (4%)	6 (1%)	16	11

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	146	PRO
2	B	328	ARG
2	B	329	ARG
2	B	330	GLU
1	A	16	ALA
1	A	282	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/234 (99%)	226 (98%)	5 (2%)	45	50
2	B	317/338 (94%)	305 (96%)	12 (4%)	29	29
All	All	548/572 (96%)	531 (97%)	17 (3%)	35	37

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ARG
1	A	35	LEU
1	A	169	LEU
1	A	283	ASP
1	A	312	LEU
2	B	5	THR
2	B	42	ARG
2	B	196	GLN
2	B	203	GLN
2	B	287	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	328	ARG
2	B	329	ARG
2	B	336	GLN
2	B	339	LEU
2	B	400	SER
2	B	402	LEU
2	B	405	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	GLN
1	A	225	HIS
1	A	251	GLN
2	B	148	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 9 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	502	-	4,4,4	0.23	0	6,6,6	0.11	0
6	MES	B	501	-	12,12,12	2.15	1 (8%)	15,16,16	2.35	6 (40%)
7	TRS	B	503	-	7,7,7	0.35	0	9,9,9	0.36	0
3	SO4	A	401	-	4,4,4	0.22	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MES	B	501	-	-	1/6/14/14	0/1/1/1
7	TRS	B	503	-	-	0/9/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	501	MES	C8-S	-7.16	1.67	1.77

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	501	MES	C5-N4-C3	5.15	119.93	108.84
6	B	501	MES	C7-N4-C5	3.30	120.04	111.24
6	B	501	MES	C2-C3-N4	-3.25	105.18	110.12
6	B	501	MES	C7-N4-C3	3.14	119.61	111.24
6	B	501	MES	C6-C5-N4	-2.75	105.94	110.12
6	B	501	MES	O1S-S-C8	2.20	110.06	106.73

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	501	MES	C8-C7-N4-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	503	TRS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/324 (98%)	-0.20	16 (5%) 34 33	16, 27, 61, 105	0
2	B	395/421 (93%)	0.02	32 (8%) 18 17	16, 27, 71, 142	0
All	All	714/745 (95%)	-0.08	48 (6%) 24 22	16, 27, 69, 142	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	135	ALA	8.1
2	B	342	ALA	6.0
2	B	146	PRO	5.6
2	B	138	TRP	5.3
1	A	53	ALA	4.6
1	A	282	VAL	4.2
2	B	136	ALA	4.1
2	B	137	PRO	4.0
2	B	310	LEU	3.9
2	B	341	THR	3.9
2	B	60	VAL	3.8
1	A	16	ALA	3.7
1	A	279	ASP	3.5
2	B	5	THR	3.5
2	B	145	PHE	3.3
1	A	281	SER	3.2
2	B	147	GLU	3.2
2	B	295	ALA	3.1
2	B	226	THR	3.1
2	B	134	SER	3.0
2	B	414	ASP	3.0
1	A	52	GLY	2.9
2	B	149	MET	2.8
2	B	148	HIS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	151	HIS	2.7
2	B	404	GLY	2.7
2	B	61	VAL	2.7
1	A	17	GLU	2.7
2	B	327	ALA	2.6
1	A	51	PRO	2.6
1	A	2	SER	2.6
2	B	224	ARG	2.6
2	B	66	THR	2.5
1	A	54	PRO	2.4
2	B	326	LEU	2.4
2	B	403	ASP	2.4
2	B	294	GLY	2.3
1	A	170	HIS	2.3
1	A	278	ALA	2.3
1	A	319	GLY	2.3
1	A	280	ARG	2.2
1	A	50	ARG	2.2
1	A	283	ASP	2.2
2	B	225	HIS	2.1
2	B	133	THR	2.1
2	B	293	LEU	2.0
2	B	296	VAL	2.0
2	B	152	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	TRS	B	503	8/8	0.75	0.18	50,55,60,70	0
4	CL	A	406	1/1	0.78	0.33	74,74,74,74	0
4	CL	A	407	1/1	0.93	0.09	60,60,60,60	0
4	CL	A	405	1/1	0.93	0.11	55,55,55,55	0
6	MES	B	501	12/12	0.94	0.10	33,40,45,55	0
4	CL	B	504	1/1	0.95	0.10	59,59,59,59	0
4	CL	A	404	1/1	0.95	0.14	58,58,58,58	0
3	SO4	A	401	5/5	0.95	0.07	51,57,60,73	0
4	CL	A	403	1/1	0.96	0.31	56,56,56,56	0
3	SO4	B	502	5/5	0.96	0.13	46,49,52,57	0
4	CL	A	408	1/1	0.97	0.22	30,30,30,30	0
4	CL	A	402	1/1	0.97	0.05	42,42,42,42	0
5	MG	A	409	1/1	0.98	0.03	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.