



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 02:42 PM UTC

PDB ID : 7Q8K / pdb\_00007q8k  
Title : Peptide LLKVAL in complex with human cathepsin V C25S mutant  
Authors : Loboda, J.; Sosnowski, P.; Tusar, L.; Vidmar, R.; Vizovisek, M.; Horvat, J.;  
Kosec, G.; Impens, F.; Demol, H.; Turk, B.; Gevaert, K.; Turk, D.  
Deposited on : 2021-11-11  
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.48.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1043 (1.74-1.74)
Clashscore	180529	1119 (1.74-1.74)
Ramachandran outliers	177936	1112 (1.74-1.74)
Sidechain outliers	177891	1112 (1.74-1.74)
RSRZ outliers	164620	1043 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AA	221	<div> <div>3%</div> <div> <div></div> <div>84%</div> <div>16%</div> </div> </div>
1	BA	221	<div> <div>3%</div> <div> <div></div> <div>80%</div> <div>19%</div> <div>.</div> </div> </div>
2	PA	6	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>
2	PB	6	<div> <div></div> <div> <div>50%</div> <div>50%</div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5433 atoms, of which 1550 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cathepsin L2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AA	221	Total	C	H	N	O	S	403	1	0
			2089	1070	392	291	326	10			
1	BA	221	Total	C	H	N	O	S	394	1	0
			2093	1070	394	292	327	10			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AA	25	SER	CYS	engineered mutation	UNP O60911
AA	108	GLN	ASN	engineered mutation	UNP O60911
AA	179	GLN	ASN	engineered mutation	UNP O60911
BA	246	SER	CYS	engineered mutation	UNP O60911
BA	329	GLN	ASN	engineered mutation	UNP O60911
BA	400	GLN	ASN	engineered mutation	UNP O60911

- Molecule 2 is a protein called LLKVAL Peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	PA	3	Total	C	H	N	O	7	0	0
			31	18	6	4	3			
2	PB	3	Total	C	H	N	O	6	0	0
			31	18	6	4	3			

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AA	1	Total	C	O	0	0
			8	6	2		
3	AA	1	Total	C	O	0	0
			8	6	2		
3	AA	1	Total	C	O	0	0
			8	6	2		
3	BA	1	Total	C	O	0	0
			8	6	2		
3	BA	1	Total	C	O	0	0
			8	6	2		
3	BA	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	AA	1	Total	C	O	0	0
			6	3	3		
4	AA	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	BA	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	AA	197	Total	H	O	394	0
			591	394	197		
6	BA	173	Total	H	O	346	0
			519	346	173		
6	PA	3	Total	H	O	6	0
			9	6	3		
6	PB	3	Total	H	O	6	0
			9	6	3		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cathepsin L2



- Molecule 1: Cathepsin L2



- Molecule 2: LLKVAL Peptide



- Molecule 2: LLKVAL Peptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.11Å 96.11Å 125.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.06 – 1.74 48.06 – 1.74	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.06-1.74) 99.5 (48.06-1.74)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.01 (at 1.74Å)	Xtriage
Refinement program	MAIN	Depositor
R, $R_{free}$	0.174 , 0.196 0.173 , 0.179	Depositor DCC
$R_{free}$ test set	2100 reflections (3.49%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.0	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 43.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5433	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	AA	1.25	11/1739 (0.6%)	1.30	18/2350 (0.8%)
1	BA	1.20	11/1741 (0.6%)	1.31	18/2352 (0.8%)
2	PA	0.80	0/24	1.25	0/30
2	PB	0.68	0/24	1.71	0/30
All	All	1.22	22/3528 (0.6%)	1.31	36/4762 (0.8%)

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AA	20	LYS	C-O	-8.38	1.15	1.24
1	BA	241	LYS	C-O	-7.63	1.16	1.24
1	AA	101	ARG	C-O	-7.52	1.18	1.25
1	BA	279	ARG	C-O	-7.23	1.17	1.24
1	AA	154	GLU	C-O	-7.02	1.18	1.25
1	AA	68	GLY	C-O	-6.26	1.16	1.23
1	BA	318	ILE	C-O	-6.26	1.16	1.24
1	AA	56	CYS	CA-C	-6.17	1.44	1.52
1	BA	378	CYS	C-O	-6.07	1.16	1.23
1	BA	430	HIS	C-O	-6.04	1.16	1.23
1	BA	288	GLY	C-O	-5.92	1.17	1.23
1	AA	67	GLY	C-O	-5.89	1.17	1.23
1	AA	22	CYS	C-O	-5.74	1.17	1.23
1	BA	274	LEU	C-O	-5.57	1.17	1.24
1	AA	24	SER	C-O	-5.45	1.16	1.24
1	AA	65	CYS	C-O	-5.40	1.17	1.24
1	AA	58	ARG	C-O	-5.37	1.19	1.24
1	BA	322	ARG	C-O	-5.32	1.18	1.25
1	AA	18	ASN	C-O	-5.26	1.18	1.24
1	BA	243	CYS	C-O	-5.13	1.18	1.23
1	BA	289	GLY	C-O	-5.04	1.18	1.23
1	BA	246	SER	C-O	-5.01	1.17	1.24



All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AA	56	CYS	N-CA-C	10.25	125.99	113.38
1	BA	278	SER	N-CA-C	9.49	123.56	112.93
1	AA	49	SER	N-CA-C	9.30	124.23	108.23
1	BA	270	SER	N-CA-C	8.75	123.28	108.23
1	BA	234	VAL	N-CA-C	8.41	119.88	108.11
1	AA	57	SER	N-CA-C	8.25	122.17	112.93
1	BA	277	CYS	N-CA-C	8.22	123.49	113.38
1	AA	129	THR	N-CA-C	7.97	123.31	113.50
1	AA	13	VAL	N-CA-C	7.78	118.99	108.11
1	BA	350	THR	N-CA-C	7.55	123.64	113.97
1	BA	407	VAL	N-CA-C	7.43	118.52	108.11
1	AA	93	ALA	N-CA-C	7.34	121.65	111.74
1	BA	273	ASN	N-CA-C	-6.58	104.03	111.07
1	BA	311	PRO	N-CA-C	6.58	121.67	111.21
1	AA	109	ASP	CA-CB-CG	6.57	119.17	112.60
1	BA	435	THR	N-CA-C	6.51	118.45	111.36
1	BA	330	ASP	CA-CB-CG	6.47	119.07	112.60
1	AA	42	THR	N-CA-C	6.21	122.76	114.12
1	AA	67	GLY	N-CA-C	6.07	118.21	112.08
1	BA	314	ALA	N-CA-C	6.07	119.94	111.74
1	AA	90	PRO	N-CA-C	6.05	120.83	111.21
1	BA	276	ASP	N-CA-C	6.04	119.85	112.23
1	BA	288	GLY	N-CA-C	5.99	118.13	112.08
1	AA	117	PRO	N-CA-C	5.91	120.37	111.14
1	AA	55	ASP	N-CA-C	5.91	118.67	111.82
1	AA	118	GLY	CA-C-N	-5.90	114.65	122.85
1	AA	118	GLY	C-N-CA	-5.90	114.65	122.85
1	BA	338	PRO	N-CA-C	5.61	119.89	111.14
1	BA	407	VAL	N-CA-CB	-5.49	103.83	111.41
1	AA	52	ASN	N-CA-C	-5.44	105.35	111.28
1	AA	130	VAL	N-CA-C	5.43	116.83	111.67
1	AA	13	VAL	N-CA-CB	-5.29	104.11	111.41
1	BA	276	ASP	CA-CB-CG	5.28	117.88	112.60
1	BA	242	GLN	N-CA-C	5.17	118.71	110.70
1	AA	214	THR	N-CA-C	5.06	116.88	111.36
1	BA	245	SER	N-CA-C	5.06	118.76	112.59

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AA	1697	392	1630	8	0
1	BA	1699	394	1626	15	0
2	PA	25	6	34	0	0
2	PB	25	6	34	0	0
3	AA	24	0	42	6	0
3	BA	24	0	42	7	0
4	AA	12	0	16	2	0
5	BA	1	0	0	0	0
6	AA	197	394	0	3	1
6	BA	173	346	0	4	0
6	PA	3	6	0	0	0
6	PB	3	6	0	0	0
All	All	3883	1550	3424	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BA:398:ASN:CB	1:BA:401:ASN:OD1	1.84	1.24
1:BA:398:ASN:HB3	1:BA:401:ASN:OD1	0.98	1.15
3:AA:304:MPD:H53	3:AA:304:MPD:C1	2.05	0.86
1:BA:372:ILE:HD11	1:BA:395:GLU:HG3	1.66	0.77
4:AA:305:GOL:H11	6:AA:442:HOH:O	1.84	0.76
3:BA:502:MPD:H53	3:BA:504:MPD:C5	2.18	0.74
3:BA:503:MPD:HM1	3:BA:503:MPD:H52	1.70	0.73
3:AA:304:MPD:H53	3:AA:304:MPD:H13	1.71	0.72
3:AA:304:MPD:H53	3:AA:304:MPD:H11	1.71	0.70
3:BA:502:MPD:O2	6:BA:601:HOH:O	2.08	0.70
1:AA:207:ASN:HB3	6:AA:570:HOH:O	1.92	0.68
1:AA:9:LYS:HE2	6:BA:616:HOH:O	1.96	0.66
1:BA:384:ASP:OD2	3:BA:503:MPD:H51	1.98	0.63
3:AA:304:MPD:C1	3:AA:304:MPD:C5	2.76	0.61
4:AA:303:GOL:H32	6:AA:413:HOH:O	1.99	0.61

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:AA:301:MPD:HM1	3:AA:301:MPD:H52	1.82	0.60
1:BA:296:GLN:O	1:BA:300:GLU:HG3	2.03	0.58
3:BA:502:MPD:O2	3:BA:502:MPD:H52	2.05	0.55
1:BA:228:TRP:CE2	1:BA:352:GLY:HA2	2.43	0.54
1:AA:7:TRP:CE2	1:AA:131:GLY:HA2	2.43	0.54
1:AA:116:ALA:CB	1:AA:119:LYS:HD2	2.38	0.53
1:BA:242:GLN:OE1	3:BA:502:MPD:H51	2.13	0.49
1:BA:307:GLU:HB2	6:BA:606:HOH:O	2.13	0.48
1:BA:401:ASN:O	1:BA:402:SER:HB2	2.13	0.48
1:AA:116:ALA:HB1	1:AA:119:LYS:HD2	1.96	0.46
1:BA:295:PHE:CZ	1:BA:440:PRO:HD3	2.51	0.46
1:AA:147:TYR:CZ	1:AA:200:VAL:HG23	2.51	0.45
3:AA:304:MPD:H11	3:AA:304:MPD:C5	2.42	0.45
1:BA:345:MET:HE3	1:BA:392:TYR:CE1	2.52	0.44
1:BA:242:GLN:N	1:BA:242:GLN:CD	2.75	0.44
3:BA:502:MPD:H53	3:BA:504:MPD:H52	1.95	0.44
1:AA:116:ALA:HB3	1:AA:119:LYS:HD2	2.01	0.41
1:BA:342:LYS:HD3	1:BA:342:LYS:HA	1.97	0.41
1:BA:394:PHE:CG	1:BA:399:SER:HA	2.56	0.41
1:AA:146:PHE:O	1:AA:148:LYS:HE2	2.20	0.41
1:BA:342:LYS:HD3	6:BA:639:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:AA:570:HOH:O	6:AA:570:HOH:O[8_555]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AA	220/221 (100%)	215 (98%)	5 (2%)	0	100	100
1	BA	220/221 (100%)	213 (97%)	7 (3%)	0	100	100
2	PA	1/6 (17%)	1 (100%)	0	0	100	100
2	PB	1/6 (17%)	1 (100%)	0	0	100	100
All	All	442/454 (97%)	430 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AA	180/179 (101%)	179 (99%)	1 (1%)	84	78
1	BA	180/179 (101%)	176 (98%)	4 (2%)	47	23
2	PA	3/5 (60%)	3 (100%)	0	100	100
2	PB	3/5 (60%)	3 (100%)	0	100	100
All	All	366/368 (100%)	361 (99%)	5 (1%)	62	45

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AA	157	CYS
1	BA	241	LYS
1	BA	378	CYS
1	BA	381	LYS
1	BA	399	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	AA	179	GLN
1	BA	240	GLN
1	BA	441	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	MPD	AA	302	-	7,7,7	0.46	0	9,10,10	0.26	0
3	MPD	BA	504	-	7,7,7	0.60	0	9,10,10	0.45	0
4	GOL	AA	303	-	5,5,5	0.10	0	5,5,5	0.30	0
3	MPD	AA	301	-	7,7,7	0.53	0	9,10,10	0.41	0
4	GOL	AA	305	-	5,5,5	0.15	0	5,5,5	0.28	0
3	MPD	BA	503	-	7,7,7	0.62	0	9,10,10	0.58	0
3	MPD	AA	304	-	7,7,7	0.40	0	9,10,10	0.28	0
3	MPD	BA	502	-	7,7,7	0.40	0	9,10,10	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	AA	302	-	-	2/5/5/5	-
3	MPD	BA	504	-	-	3/5/5/5	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	AA	303	-	-	0/4/4/4	-
3	MPD	AA	301	-	-	1/5/5/5	-
4	GOL	AA	305	-	-	0/4/4/4	-
3	MPD	BA	503	-	-	2/5/5/5	-
3	MPD	AA	304	-	-	3/5/5/5	-
3	MPD	BA	502	-	-	5/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	BA	502	MPD	C2-C3-C4-O4
3	BA	502	MPD	C2-C3-C4-C5
3	BA	503	MPD	C2-C3-C4-O4
3	AA	301	MPD	CM-C2-C3-C4
3	AA	304	MPD	C1-C2-C3-C4
3	AA	304	MPD	CM-C2-C3-C4
3	BA	502	MPD	C1-C2-C3-C4
3	BA	504	MPD	C1-C2-C3-C4
3	BA	504	MPD	CM-C2-C3-C4
3	AA	302	MPD	C2-C3-C4-C5
3	AA	304	MPD	C2-C3-C4-C5
3	BA	503	MPD	C2-C3-C4-C5
3	BA	504	MPD	C2-C3-C4-C5
3	BA	502	MPD	O2-C2-C3-C4
3	AA	302	MPD	CM-C2-C3-C4
3	BA	502	MPD	CM-C2-C3-C4

There are no ring outliers.

7 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	BA	504	MPD	2	0
4	AA	303	GOL	1	0
3	AA	301	MPD	1	0
4	AA	305	GOL	1	0
3	BA	503	MPD	2	0
3	AA	304	MPD	5	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	BA	502	MPD	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	AA	221/221 (100%)	-0.18	7 (3%)	50	58	11, 17, 29, 51	6 (2%)
1	BA	221/221 (100%)	-0.01	6 (2%)	56	63	12, 20, 36, 71	1 (0%)
2	PA	3/6 (50%)	0.87	0	100	100	21, 21, 21, 29	1 (33%)
2	PB	3/6 (50%)	1.02	0	100	100	23, 23, 27, 32	0
All	All	448/454 (98%)	-0.08	13 (2%)	54	61	11, 19, 34, 71	8 (1%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AA	116	ALA	6.2
1	AA	117	PRO	3.8
1	AA	180	ASN	3.5
1	AA	176	ALA	3.1
1	BA	400[A]	GLN	2.9
1	BA	414	GLU	2.5
1	BA	398	ASN	2.5
1	BA	401	ASN	2.5
1	AA	118	GLY	2.4
1	BA	222	LEU	2.4
1	AA	207	ASN	2.2
1	AA	179	GLN	2.2
1	BA	397	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MPD	AA	301	8/8	0.82	0.37	81,84,97,97	0
3	MPD	AA	304	8/8	0.83	0.29	98,105,110,111	0
3	MPD	BA	502	8/8	0.83	0.19	52,54,57,58	0
4	GOL	AA	305	6/6	0.83	0.25	81,84,88,89	0
3	MPD	AA	302	8/8	0.85	0.29	83,86,97,103	0
3	MPD	BA	504	8/8	0.88	0.24	74,78,83,86	0
4	GOL	AA	303	6/6	0.89	0.37	99,102,104,112	0
3	MPD	BA	503	8/8	0.89	0.13	35,38,41,43	0
5	CL	BA	501	1/1	0.98	0.06	26,26,26,26	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.