



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:49 PM UTC

PDB ID : 7VDR / pdb\_00007vdr  
Title : The structure of cyclin-dependent kinase 5 (CDK5) in complex with p25 and Compound 13  
Authors : Malojcic, G.; Clugston, S.L.; Daniels, M.; Harmange, J.C.; Ledeborer, M.  
Deposited on : 2021-09-07  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	<div> <div>2%</div> <div>92%</div> <div>8%</div> </div>
1	B	292	<div> <div>9%</div> <div>85%</div> <div>9%</div> <div>6%</div> </div>
2	C	209	<div> <div>%</div> <div>65%</div> <div>6%</div> <div>29%</div> </div>
2	D	209	<div> <div>33%</div> <div>57%</div> <div>8%</div> <div>35%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7028 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent-like kinase 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	0	0	0
			2330	1495	400	424	11			
1	B	275	Total	C	N	O	S	0	0	0
			2217	1428	384	395	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q00535
B	1	SER	-	expression tag	UNP Q00535


- Molecule 2 is a protein called Cyclin-dependent kinase 5 activator 1, p25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	148	Total	C	N	O	S	0	0	0
			1196	768	197	220	11			
2	D	136	Total	C	N	O	S	0	0	0
			1104	711	179	204	10			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	99	MET	-	initiating methionine	UNP Q15078
D	99	MET	-	initiating methionine	UNP Q15078

- Molecule 3 is (1R)-1-[7-[(2-fluoranyl-4-pyrazol-1-yl-phenyl)amino]-1,6-naphthyridin-2-yl]-1-(1-methylpiperidin-4-yl)ethanol (CCD ID: 63I) (formula: C<sub>25</sub>H<sub>27</sub>FN<sub>6</sub>O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			33	25	1	6	1		
3	B	1	Total	C	F	N	O	0	0
			33	25	1	6	1		

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula:  $C_6H_{14}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	41	Total O 41 41	0	0
6	B	15	Total O 15 15	0	0
6	C	13	Total O 13 13	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclin-dependent-like kinase 5




- Molecule 1: Cyclin-dependent-like kinase 5



- Molecule 2: Cyclin-dependent kinase 5 activator 1, p25



- Molecule 2: Cyclin-dependent kinase 5 activator 1, p25





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	117.83Å 117.83Å 154.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.65 – 2.55 29.65 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.65-2.55) 99.9 (29.65-2.55)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.20 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.219 , 0.256 0.220 , 0.251	Depositor DCC
$R_{free}$ test set	1980 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	61.2	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7028	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, 63I, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.97	0/2385	1.37	0/3229
1	B	0.99	0/2267	1.37	0/3065
2	C	0.98	0/1224	1.49	0/1660
2	D	1.03	0/1129	1.50	0/1530
All	All	0.99	0/7005	1.41	0/9484

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2330	0	2344	12	0
1	B	2217	0	2257	17	0
2	C	1196	0	1185	9	0
2	D	1104	0	1088	10	0
3	A	33	0	0	1	0
3	B	33	0	0	0	0
4	A	10	0	14	0	0
5	A	32	0	48	0	0
5	C	4	0	6	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	41	0	0	1	0
6	B	15	0	0	0	0
6	C	13	0	0	0	0
All	All	7028	0	6942	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:CYS:HA	1:A:135:ASN:HD21	1.45	0.81
1:B:247:SER:O	1:B:248:LEU:HB2	1.91	0.70
1:B:19:PHE:HE1	1:B:34:ARG:HD3	1.63	0.63
1:B:157:CYS:HB2	1:B:177:LYS:HB3	1.85	0.58
2:D:204:LEU:HD12	2:D:229:CYS:SG	2.45	0.57
1:B:37:LEU:HD23	2:D:258:TRP:CD1	2.42	0.54
1:B:44:VAL:N	1:B:45:PRO:HD2	2.23	0.53
1:B:49:LEU:HA	1:B:52:ILE:HD12	1.92	0.52
1:A:83:CYS:HA	1:A:135:ASN:ND2	2.20	0.52
1:A:43:GLY:O	2:C:242:SER:HB2	2.11	0.51
1:B:63:ILE:HD13	1:B:142:LEU:HB3	1.96	0.48
2:C:246:LYS:HB3	2:C:247:PRO:HD3	1.96	0.47
1:A:50:ARG:NH1	2:C:235:SER:O	2.47	0.47
1:A:152:GLY:HA3	2:C:276:ASN:O	2.15	0.47
1:B:168:ARG:HD3	1:B:172:VAL:HG12	1.97	0.46
1:B:159:SER:HB2	2:D:239:ASN:O	2.16	0.46
2:D:246:LYS:HB3	2:D:247:PRO:HD3	1.98	0.46
1:B:196:ALA:HB1	1:B:255:LEU:HG	1.97	0.46
1:B:209:ASP:OD1	1:B:213:LYS:HE2	2.15	0.45
2:C:177:TRP:CD2	2:C:223:GLN:HG3	2.52	0.44
1:B:114:LEU:HD12	1:B:114:LEU:HA	1.89	0.44
1:B:153:ILE:O	1:B:155:VAL:HG23	2.18	0.43
1:B:49:LEU:HD11	2:D:235:SER:CB	2.48	0.42
2:D:190:TRP:CD1	2:D:244:PRO:HG3	2.55	0.42
1:A:84:ASP:H	1:A:135:ASN:HD22	1.68	0.42
2:D:207:LEU:HB2	2:D:229:CYS:SG	2.59	0.42
2:D:260:ARG:O	2:D:264:VAL:HG23	2.19	0.42
1:B:49:LEU:HB2	2:D:265:ILE:HD11	2.01	0.42
1:B:19:PHE:CE1	1:B:34:ARG:HD3	2.49	0.41
1:A:76:LEU:HD11	2:C:262:LEU:HD11	2.01	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:ILE:HG23	3:A:301:63I:C23	2.51	0.41
1:A:196:ALA:HB1	1:A:255:LEU:HG	2.02	0.41
1:A:213:LYS:NZ	6:A:403:HOH:O	2.54	0.41
2:C:260:ARG:O	2:C:264:VAL:HG23	2.21	0.41
1:A:153:ILE:HG21	2:C:199:ALA:HB1	2.02	0.41
1:A:104:LYS:NZ	1:A:285:TYR:O	2.41	0.40
2:C:173:ASP:HB2	2:C:174:PRO:HD3	2.02	0.40
1:B:57:GLU:CD	2:D:270:SER:HA	2.46	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/292 (99%)	276 (96%)	11 (4%)	1 (0%)	36	45
1	B	269/292 (92%)	253 (94%)	14 (5%)	2 (1%)	18	25
2	C	146/209 (70%)	146 (100%)	0	0	100	100
2	D	130/209 (62%)	126 (97%)	4 (3%)	0	100	100
All	All	833/1002 (83%)	801 (96%)	29 (4%)	3 (0%)	30	39

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	163	VAL
1	B	163	VAL
1	B	248	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/260 (99%)	251 (97%)	7 (3%)	39	58
1	B	246/260 (95%)	245 (100%)	1 (0%)	84	90
2	C	138/187 (74%)	138 (100%)	0	100	100
2	D	128/187 (68%)	125 (98%)	3 (2%)	44	62
All	All	770/894 (86%)	759 (99%)	11 (1%)	59	75

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	9	LYS
1	A	14	THR
1	A	17	THR
1	A	61	LYS
1	A	127	LEU
1	A	232	LYS
1	B	147	LEU
2	D	170	SER
2	D	186	LEU
2	D	259	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	GLN
1	A	135	ASN
2	C	191	GLN
2	D	188	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	EDO	C	401	-	3,3,3	0.08	0	2,2,2	0.16	0
4	PGE	A	302	-	9,9,9	0.23	0	8,8,8	0.09	0
5	EDO	A	305	-	3,3,3	0.04	0	2,2,2	0.09	0
5	EDO	A	309	-	3,3,3	0.14	0	2,2,2	0.29	0
5	EDO	A	308	-	3,3,3	0.07	0	2,2,2	0.10	0
5	EDO	A	310	-	3,3,3	0.08	0	2,2,2	0.12	0
5	EDO	A	303	-	3,3,3	0.10	0	2,2,2	0.19	0
3	63I	A	301	-	35,37,37	0.74	1 (2%)	51,54,54	2.11	10 (19%)
5	EDO	A	307	-	3,3,3	0.08	0	2,2,2	0.07	0
5	EDO	A	306	-	3,3,3	0.09	0	2,2,2	0.03	0
3	63I	B	301	-	35,37,37	0.76	1 (2%)	51,54,54	2.04	11 (21%)
5	EDO	A	304	-	3,3,3	0.06	0	2,2,2	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	C	401	-	-	1/1/1/1	-
4	PGE	A	302	-	-	3/7/7/7	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	305	-	-	0/1/1/1	-
5	EDO	A	309	-	-	0/1/1/1	-
5	EDO	A	308	-	-	1/1/1/1	-
5	EDO	A	310	-	-	0/1/1/1	-
5	EDO	A	303	-	-	1/1/1/1	-
3	63I	A	301	-	-	3/19/30/30	0/5/5/5
5	EDO	A	307	-	-	1/1/1/1	-
5	EDO	A	306	-	-	1/1/1/1	-
3	63I	B	301	-	-	4/19/30/30	0/5/5/5
5	EDO	A	304	-	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	301	63I	N26-N30	2.84	1.40	1.36
3	A	301	63I	N26-N30	2.78	1.40	1.36

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	63I	C11-N33-C32	6.56	122.64	117.37
3	A	301	63I	C27-N26-N30	-6.46	106.43	111.83
3	B	301	63I	C27-N26-N30	-5.98	106.84	111.83
3	B	301	63I	C11-N33-C32	5.95	122.15	117.37
3	A	301	63I	C15-N16-C17	5.42	122.91	117.81
3	B	301	63I	C15-N16-C17	5.24	122.74	117.81
3	A	301	63I	C29-N30-N26	5.20	109.01	103.94
3	B	301	63I	C29-N30-N26	5.16	108.97	103.94
3	A	301	63I	C31-C32-N33	3.12	123.34	118.78
3	B	301	63I	C31-C32-N33	2.93	123.06	118.78
3	A	301	63I	C9-C8-C5	-2.85	108.70	112.45
3	B	301	63I	C22-N26-N30	2.65	125.60	121.08
3	B	301	63I	C3-N2-C7	2.61	113.69	109.54
3	A	301	63I	C22-N26-N30	2.59	125.50	121.08
3	B	301	63I	C23-C24-C19	-2.52	120.85	123.43
3	A	301	63I	C14-C32-N33	-2.44	118.90	122.25
3	A	301	63I	C28-C27-N26	2.38	108.72	106.76
3	B	301	63I	C14-C15-N16	-2.20	120.70	124.25
3	A	301	63I	C31-C17-N16	-2.20	120.04	122.93
3	B	301	63I	C12-C11-C8	-2.18	118.81	122.11
3	B	301	63I	C14-C32-N33	-2.14	119.30	122.25

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	63I	C12-C11-C8-C5
3	B	301	63I	C12-C11-C8-C5
4	A	302	PGE	O1-C1-C2-O2
5	A	303	EDO	O1-C1-C2-O2
5	A	306	EDO	O1-C1-C2-O2
5	A	304	EDO	O1-C1-C2-O2
3	B	301	63I	C12-C11-C8-C9
4	A	302	PGE	O3-C5-C6-O4
3	A	301	63I	C12-C11-C8-C9
3	B	301	63I	N33-C11-C8-O10
5	A	307	EDO	O1-C1-C2-O2
4	A	302	PGE	C1-C2-O2-C3
3	B	301	63I	C12-C11-C8-O10
5	A	308	EDO	O1-C1-C2-O2
5	C	401	EDO	O1-C1-C2-O2
3	A	301	63I	C12-C11-C8-O10


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	63I	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


## Ligand 63I A 301




Bond lengths



Bond angles




Torsions



Rings





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	290/292 (99%)	0.03	6 (2%) 63 64	39, 54, 96, 139	0
1	B	275/292 (94%)	0.75	25 (9%) 15 14	52, 83, 136, 175	0
2	C	148/209 (70%)	0.02	2 (1%) 73 74	40, 53, 78, 87	0
2	D	136/209 (65%)	2.19	68 (50%) 0 0	117, 155, 173, 182	0
All	All	849/1002 (84%)	0.61	101 (11%) 9 8	39, 70, 164, 182	0

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	222	LEU	7.2
2	D	226	LEU	7.1
2	D	175	VAL	6.8
2	D	155	LEU	5.8
2	D	178	LEU	5.3
1	A	15	TYR	5.2
2	D	225	VAL	5.0
1	A	40	ASP	4.9
1	A	14	THR	4.6
2	D	196	ILE	4.4
2	D	169	LEU	4.4
2	D	207	LEU	4.3
2	D	151	LEU	4.2
2	D	163	CYS	4.2
2	D	176	LEU	4.0
2	D	227	LEU	3.7
2	D	203	PHE	3.7
1	B	56	LYS	3.7
1	B	18	VAL	3.6
2	D	229	CYS	3.6
2	D	177	TRP	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	285	VAL	3.5
2	D	281	TYR	3.5
2	D	152	LEU	3.5
2	D	228	THR	3.4
2	C	146	ALA	3.4
2	C	241	ILE	3.3
2	D	181	VAL	3.3
2	D	293	SER	3.2
1	A	291	PRO	3.2
1	B	70	LEU	3.2
1	B	9	LYS	3.2
2	D	224	ALA	3.2
2	D	168	HIS	3.1
2	D	198	PRO	3.0
2	D	262	LEU	3.0
2	D	282	PHE	3.0
1	B	3	LYS	3.0
2	D	241	ILE	2.9
2	D	244	PRO	2.9
2	D	216	VAL	2.9
2	D	166	LEU	2.9
1	B	64	VAL	2.8
2	D	164	TYR	2.8
2	D	250	VAL	2.8
2	D	148	THR	2.8
2	D	267	LEU	2.8
1	B	29	ILE	2.7
2	D	233	SER	2.7
2	D	258	TRP	2.7
2	D	248	PHE	2.7
2	D	174	PRO	2.7
1	B	244	ALA	2.7
2	D	171	PRO	2.7
2	D	230	LEU	2.7
1	B	245	THR	2.7
2	D	275	ILE	2.6
1	B	66	LEU	2.6
1	B	262	LEU	2.6
2	D	273	LEU	2.6
1	B	24	ARG	2.5
2	D	190	TRP	2.5
1	B	49	LEU	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	26	THR	2.5
2	D	220	HIS	2.5
2	D	201	VAL	2.5
2	D	185	LEU	2.4
1	B	288	ASP	2.4
2	D	205	TYR	2.4
1	B	69	VAL	2.4
2	D	195	PHE	2.4
2	D	247	PRO	2.3
1	B	58	LEU	2.3
1	B	55	LEU	2.3
2	D	172	THR	2.3
2	D	283	THR	2.3
2	D	257	PHE	2.3
1	B	78	LEU	2.3
2	D	286	PHE	2.3
2	D	256	ALA	2.2
2	D	223	GLN	2.2
2	D	245	LEU	2.2
2	D	263	SER	2.2
1	B	63	ILE	2.2
2	D	242	SER	2.2
1	B	251	VAL	2.2
1	B	277	ALA	2.2
2	D	202	VAL	2.1
2	D	232	LEU	2.1
2	D	200	ASN	2.1
2	D	187	LEU	2.1
2	D	147	SER	2.1
2	D	219	ASP	2.1
2	D	287	SER	2.1
1	A	16	GLY	2.1
1	A	225	GLU	2.1
2	D	218	SER	2.1
1	B	27	HIS	2.1
2	D	268	MET	2.1
1	B	25	GLU	2.0
1	B	10	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates

There are no oligosaccharides in this entry.

## 6.4 Ligands



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	EDO	A	310	4/4	0.57	0.23	92,93,94,94	0
5	EDO	A	305	4/4	0.59	0.31	83,86,87,90	0
5	EDO	A	309	4/4	0.81	0.21	65,73,74,74	0
5	EDO	A	304	4/4	0.81	0.24	87,89,90,91	0
4	PGE	A	302	10/10	0.82	0.20	79,97,106,107	0
5	EDO	A	303	4/4	0.84	0.24	74,76,78,79	0
5	EDO	A	306	4/4	0.87	0.17	70,71,71,71	0
5	EDO	A	307	4/4	0.87	0.14	78,83,86,87	0
5	EDO	C	401	4/4	0.88	0.16	70,74,76,79	0
5	EDO	A	308	4/4	0.89	0.16	81,83,84,87	0
3	63I	B	301	33/33	0.92	0.10	68,85,95,95	0
3	63I	A	301	33/33	0.97	0.07	41,45,59,62	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around 63I B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 63I A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.