



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 11:26 PM UTC

PDB ID : 8FEN / pdb_00008fen
Title : Panicum vibratum Dihydroflavonol 4-reductase complexed with NADP and DHQ
Authors : Lewis, J.A.; Kang, C.
Deposited on : 2022-12-06
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	<div> <div>6%</div> <div>75%</div> <div>13%</div> <div>11%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5076 atoms, of which 2447 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called *Panicum virgatum* Dihydroflavonol 4-Reductase.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	326	4962	1608	2447	431	462	14	0	0	0

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0

- Molecule 3 is (2R,3R)-2-(3,4-DIHYDROXYPHENYL)-3,5,7-TRIHYDROXY-2,3-DIHYDRO-4H-CHROMEN-4-ONE (CCD ID: DQH) (formula: $C_{15}H_{12}O_7$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			22	15	7		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	44	Total	O	0	0
			44	44		

- Molecule 1: *Panicum virgatum* Dihydroflavonol 4-Reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	119.06Å 119.06Å 56.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.65 – 2.55 37.65 – 2.55	Depositor EDS
% Data completeness (in resolution range)	95.5 (37.65-2.55) 95.6 (37.65-2.55)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 2.54Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.215 , 0.238 0.215 , 0.238	Depositor DCC
R_{free} test set	1329 reflections (9.55%)	wwPDB-VP
Wilson B-factor (Å ²)	41.7	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 36.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5076	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DQH, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/2574	0.34	0/3495

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2515	2447	2516	38	0
2	A	48	0	24	4	0
3	A	22	0	12	1	0
4	A	44	0	0	3	0
All	All	2629	2447	2552	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:401:NAP:O4D	2:A:401:NAP:C1D	1.64	1.16
1:A:62:PRO:HB2	1:A:213:ILE:HD13	1.65	0.78
1:A:341:ILE:HG13	1:A:342:PRO:HD2	1.74	0.68
1:A:339:GLY:O	1:A:340:LEU:HD23	1.93	0.68
1:A:142:SER:O	2:A:401:NAP:H6N	1.95	0.66
1:A:234:HIS:HA	1:A:237:ILE:HD12	1.81	0.62
1:A:57:VAL:HG13	1:A:61:LYS:HB2	1.86	0.57
1:A:239:LYS:HE2	1:A:303:LEU:HD21	1.88	0.56
1:A:341:ILE:CG1	1:A:342:PRO:HD2	2.36	0.56
1:A:213:ILE:C	1:A:213:ILE:HD12	2.31	0.55
1:A:39:LYS:NZ	1:A:251:ASP:OD1	2.40	0.54
1:A:225:LEU:HD13	1:A:340:LEU:HD11	1.90	0.54
1:A:206:PRO:O	2:A:401:NAP:H4N	2.09	0.53
1:A:28:ALA:CB	1:A:49:ALA:HB1	2.40	0.52
1:A:287:PRO:HD2	1:A:288:GLU:OE1	2.09	0.52
1:A:334:THR:O	1:A:337:GLU:HG3	2.10	0.51
1:A:50:THR:CG2	1:A:78:ALA:HB3	2.41	0.50
1:A:234:HIS:HA	1:A:237:ILE:CD1	2.43	0.49
1:A:35:TRP:CB	1:A:246:LEU:HD21	2.42	0.48
1:A:35:TRP:HB2	1:A:246:LEU:HD21	1.95	0.48
1:A:39:LYS:HE3	1:A:247:ASP:OD1	2.13	0.47
1:A:177:MET:HA	1:A:180:VAL:HG22	1.97	0.47
1:A:217:MET:HE1	1:A:340:LEU:HD21	1.96	0.46
1:A:54:PRO:HG2	1:A:77:LYS:HB2	1.97	0.46
1:A:171:VAL:O	4:A:501:HOH:O	2.21	0.46
1:A:50:THR:HG23	1:A:78:ALA:HB3	1.97	0.45
1:A:109:ASP:O	1:A:110:PRO:C	2.59	0.45
1:A:341:ILE:CD1	1:A:342:PRO:HD2	2.45	0.45
1:A:21:GLY:N	4:A:508:HOH:O	2.49	0.45
1:A:62:PRO:CB	1:A:213:ILE:HD13	2.43	0.43
1:A:332:ILE:HG22	1:A:333:ARG:N	2.31	0.43
1:A:26:THR:HA	1:A:50:THR:HB	2.01	0.43
1:A:113:GLU:O	1:A:113:GLU:HG3	2.18	0.42
1:A:122:VAL:O	1:A:126:MET:HG2	2.19	0.42
1:A:49:ALA:HB3	1:A:75:ILE:HD13	2.03	0.41
1:A:148:ASN:OD1	3:A:402:DQH:O23	2.28	0.41
1:A:32:VAL:HB	2:A:401:NAP:H51N	2.03	0.40
1:A:269:SER:O	4:A:502:HOH:O	2.21	0.40
1:A:225:LEU:HD13	1:A:340:LEU:CD1	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	324/366 (88%)	302 (93%)	18 (6%)	4 (1%)	10 13

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	229	THR
1	A	56	ASN
1	A	301	GLY
1	A	206	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	265/286 (93%)	263 (99%)	2 (1%)	73 84

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ARG
1	A	341	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	240	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAP	A	401	-	50,52,52	4.55	23 (46%)	71,80,80	1.75	19 (26%)
3	DQH	A	402	-	24,24,24	2.95	8 (33%)	36,36,36	1.15	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	401	-	-	5/35/67/67	0/5/5/5
3	DQH	A	402	-	-	0/4/20/20	0/3/3/3

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NAP	O4D-C1D	17.63	1.64	1.40
2	A	401	NAP	PN-O3	13.33	1.73	1.59
2	A	401	NAP	C3B-C2B	-12.55	1.25	1.53
3	A	402	DQH	O12-C4	11.51	1.53	1.38
2	A	401	NAP	C7N-N7N	8.29	1.48	1.33
2	A	401	NAP	O4B-C4B	-6.86	1.29	1.45
2	A	401	NAP	O4D-C4D	-5.76	1.32	1.45
2	A	401	NAP	C3B-C4B	5.08	1.65	1.53
2	A	401	NAP	P2B-O2B	4.82	1.68	1.59
2	A	401	NAP	O3D-C3D	-4.34	1.32	1.43
2	A	401	NAP	O7N-C7N	-4.18	1.16	1.24
2	A	401	NAP	C6A-N6A	4.15	1.44	1.34
2	A	401	NAP	O4B-C1B	4.01	1.51	1.42
2	A	401	NAP	C1B-N9A	-3.91	1.35	1.46
3	A	402	DQH	O23-C18	3.53	1.43	1.36
2	A	401	NAP	C5A-N7A	-3.53	1.32	1.39
3	A	402	DQH	C10-C9	-3.43	1.46	1.52
3	A	402	DQH	O24-C17	3.42	1.43	1.36
3	A	402	DQH	O27-C10	3.30	1.48	1.42
2	A	401	NAP	C3N-C7N	3.07	1.55	1.50
2	A	401	NAP	O3B-C3B	3.00	1.50	1.43
2	A	401	NAP	O2B-C2B	2.93	1.54	1.44
2	A	401	NAP	PA-O5B	2.83	1.70	1.59
2	A	401	NAP	C4A-N9A	-2.34	1.32	1.37
2	A	401	NAP	C2A-N1A	2.31	1.38	1.33
2	A	401	NAP	O2D-C2D	2.31	1.48	1.43
3	A	402	DQH	O30-C2	2.31	1.41	1.36
2	A	401	NAP	PN-O5D	2.14	1.67	1.59
3	A	402	DQH	C5-C4	2.04	1.42	1.38
3	A	402	DQH	C16-C15	2.01	1.42	1.38
2	A	401	NAP	C4N-C3N	-2.00	1.36	1.39

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NAP	N3A-C2A-N1A	-5.66	120.02	128.58
2	A	401	NAP	C5A-C4A-N3A	-4.95	119.89	126.72
2	A	401	NAP	N9A-C8A-N7A	-4.44	107.64	113.94
2	A	401	NAP	C4A-N9A-C1B	-3.40	118.67	126.63
2	A	401	NAP	C2A-N3A-C4A	3.37	120.07	111.83
2	A	401	NAP	N3A-C4A-N9A	3.05	132.36	127.17
2	A	401	NAP	C4D-O4D-C1D	-3.05	107.13	109.92
2	A	401	NAP	C5A-N7A-C8A	3.02	108.20	103.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	DQH	C2-C3-C4	2.74	120.02	117.41
2	A	401	NAP	C1B-N9A-C8A	2.64	132.95	127.09
3	A	402	DQH	O12-C4-C3	-2.57	119.06	121.64
3	A	402	DQH	O12-C4-C5	2.56	120.66	116.29
2	A	401	NAP	C4A-N9A-C8A	2.52	108.38	105.74
2	A	401	NAP	C2B-C3B-C4B	2.42	107.19	101.99
2	A	401	NAP	C2B-C1B-N9A	-2.29	109.98	113.75
2	A	401	NAP	C4B-O4B-C1B	-2.26	104.47	109.47
2	A	401	NAP	C4A-C5A-N7A	-2.22	108.04	110.58
2	A	401	NAP	C6N-N1N-C2N	-2.19	120.02	121.88
2	A	401	NAP	C3N-C7N-N7N	2.17	120.42	117.74
3	A	402	DQH	O12-C11-C14	2.12	111.97	107.44
2	A	401	NAP	P2B-O2B-C2B	-2.11	117.79	123.43
2	A	401	NAP	N6A-C6A-N1A	-2.04	113.83	118.38
2	A	401	NAP	O7N-C7N-N7N	-2.02	119.69	122.62

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	NAP	PN-O3-PA-O5B
2	A	401	NAP	C5D-O5D-PN-O3
2	A	401	NAP	C5D-O5D-PN-O1N
2	A	401	NAP	C5D-O5D-PN-O2N
2	A	401	NAP	PA-O3-PN-O1N


There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	NAP	4	0
3	A	402	DQH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/366 (89%)	0.36	22 (6%) 24 23	30, 51, 85, 118	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	341	ILE	6.3
1	A	21	GLY	4.8
1	A	342	PRO	4.3
1	A	340	LEU	3.4
1	A	105	PHE	3.3
1	A	339	GLY	3.3
1	A	107	SER	3.1
1	A	229	THR	3.0
1	A	228	ILE	2.9
1	A	227	LEU	2.7
1	A	299	ASP	2.7
1	A	216	GLY	2.6
1	A	213	ILE	2.6
1	A	287	PRO	2.6
1	A	302	ASP	2.6
1	A	215	ALA	2.5
1	A	338	LYS	2.5
1	A	291	ILE	2.4
1	A	214	SER	2.4
1	A	152	ARG	2.4
1	A	17	ALA	2.3
1	A	104	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	DQH	A	402	22/22	0.76	0.18	51,57,64,65	0
2	NAP	A	401	48/48	0.95	0.11	35,46,65,124	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around NAP A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.