



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:15 PM UTC

PDB ID : 8FNY / pdb\_00008fny  
Title : Nucleotide-bound structure of a functional construct of eukaryotic elongation factor 2 kinase.  
Authors : Piserchio, A.; Isiorho, E.A.; Dalby, K.N.; Ghose, R.  
Deposited on : 2022-12-28  
Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	531	<div> <div>7%</div> <div>84%</div> <div>9%</div> <div>6%</div> </div>
1	C	531	<div> <div>7%</div> <div>85%</div> <div>7%</div> <div>8%</div> </div>
2	B	148	<div> <div>3%</div> <div>40%</div> <div>8%</div> <div>52%</div> </div>
2	D	148	<div> <div>%</div> <div>39%</div> <div>6%</div> <div>55%</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 9391 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Eukaryotic elongation factor 2 kinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	497	Total	C	N	O	P	S	0	0	0
			3949	2491	698	734	1	25			
1	C	489	Total	C	N	O	P	S	0	2	0
			3920	2474	693	727	1	25			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	484	GLY	-	linker	UNP O00418
A	485	GLY	-	linker	UNP O00418
A	486	GLY	-	linker	UNP O00418
A	487	GLY	-	linker	UNP O00418
A	488	GLY	-	linker	UNP O00418
A	489	GLY	-	linker	UNP O00418
C	484	GLY	-	linker	UNP O00418
C	485	GLY	-	linker	UNP O00418
C	486	GLY	-	linker	UNP O00418
C	487	GLY	-	linker	UNP O00418
C	488	GLY	-	linker	UNP O00418
C	489	GLY	-	linker	UNP O00418

- Molecule 2 is a protein called Calmodulin-1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	71	Total	C	N	O	S	0	0	0
			562	342	93	123	4			
2	D	67	Total	C	N	O	S	0	0	0
			529	323	87	115	4			

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
3	C	1	Total 31	C 10	N 5	O 13	P 3	0	0

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula:  $\text{C}_{10}\text{H}_{15}\text{N}_5\text{O}_{10}\text{P}_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		
5	C	1	Total	Zn	0	0
			1	1		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	2	Total	Ca	0	0
			2	2		
6	D	2	Total	Ca	0	0
			2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	138	Total	O	0	0
			138	138		
7	B	15	Total	O	0	0
			15	15		
7	C	144	Total	O	0	0
			144	144		
7	D	12	Total	O	0	0
			12	12		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Eukaryotic elongation factor 2 kinase




- Molecule 1: Eukaryotic elongation factor 2 kinase



- Molecule 2: Calmodulin-1



● Molecule 2: Calmodulin-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.16Å 83.35Å 88.98Å 65.35° 90.03° 86.47°	Depositor
Resolution (Å)	37.79 – 2.22 37.79 – 2.22	Depositor EDS
% Data completeness (in resolution range)	50.5 (37.79-2.22) 50.5 (37.79-2.22)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 2.22Å)	Xtriage
Refinement program	ISOLDE 1.4, PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.208 , 0.229 0.217 , 0.236	Depositor DCC
$R_{free}$ test set	1913 reflections (2.52%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.3	Xtriage
Anisotropy	0.096	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 29.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	9391	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, ZN, CA, ATP, TPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.14	0/4027	0.32	0/5430
1	C	0.10	0/3998	0.28	0/5387
2	B	0.15	0/567	0.31	0/759
2	D	0.11	0/534	0.27	0/717
All	All	0.12	0/9126	0.30	0/12293

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3949	0	3749	28	0
1	C	3920	0	3718	21	0
2	B	562	0	514	7	0
2	D	529	0	479	5	0
3	A	31	0	12	0	0
3	C	31	0	12	3	0
4	A	27	0	12	0	0
4	C	27	0	12	1	0
5	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	1	0	0	0	0
6	B	2	0	0	0	0
6	D	2	0	0	0	0
7	A	138	0	0	2	0
7	B	15	0	0	0	0
7	C	144	0	0	0	0
7	D	12	0	0	0	0
All	All	9391	0	8508	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:328:LEU:H	1:A:333:ARG:HH21	1.37	0.71
1:C:146:CYS:SG	3:C:801:ATP:H5'2	2.32	0.69
1:C:247:ARG:NH2	1:C:250:ASN:O	2.25	0.69
3:C:801:ATP:H8	3:C:801:ATP:H5'1	1.61	0.65
1:C:247:ARG:HH22	1:C:252:ARG:HG3	1.61	0.64
1:A:306:ALA:HB1	1:A:530:MET:HG2	1.80	0.60
1:C:306:ALA:HB1	1:C:530:MET:HG2	1.82	0.59
1:A:327:ASP:OD2	1:A:502:VAL:HG23	2.02	0.59
2:B:139:GLU:O	2:B:143:GLN:HG2	2.05	0.56
1:A:112:ALA:HB2	1:A:217:LEU:HD13	1.90	0.53
1:C:209:ILE:HG13	1:C:282:TYR:HB2	1.90	0.53
1:C:588:LYS:O	1:C:593:ASN:ND2	2.42	0.53
1:A:273:VAL:HG22	1:A:274:ASP:H	1.76	0.51
1:C:294:ASP:OD1	1:C:294:ASP:N	2.44	0.51
1:A:294:ASP:OD1	1:A:294:ASP:N	2.44	0.51
1:A:82:LYS:HG2	1:A:86:LYS:HD2	1.93	0.50
2:D:117:THR:HG23	2:D:120:GLU:H	1.78	0.49
2:B:109:MET:HG3	2:B:116:LEU:HD12	1.94	0.49
2:B:117:THR:HG22	2:B:120:GLU:OE2	2.13	0.48
1:A:209:ILE:HG13	1:A:282:TYR:HB2	1.94	0.48
2:B:101:SER:HA	2:B:135:GLN:HG2	1.96	0.47
1:C:183:GLU:OE1	1:C:186:ARG:NH1	2.49	0.46
1:A:248:ASP:HA	1:A:251:ILE:HD12	1.97	0.45
1:A:86:LYS:HA	1:A:89:ILE:HD12	1.97	0.45
1:A:530:MET:HB3	1:A:530:MET:HE2	1.78	0.45
1:A:663:PRO:HD2	1:A:666:MET:HG3	1.97	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:PHE:HD1	2:B:127:GLU:HG2	1.82	0.45
1:C:663:PRO:HD2	1:C:666:MET:HG3	1.98	0.45
1:C:187:LEU:HD12	1:C:295:PHE:HB3	1.99	0.45
1:C:242:ASN:HB2	1:C:288:HIS:HE2	1.81	0.44
2:D:144:MET:HE3	2:D:144:MET:HB3	1.91	0.44
1:A:610:ARG:HA	1:A:613:MET:HE2	1.99	0.44
1:C:530:MET:HB3	1:C:530:MET:HE2	1.79	0.44
1:C:675:LEU:HD12	1:C:692:LEU:HD12	1.99	0.44
1:C:247:ARG:HH11	1:C:247:ARG:HB3	1.83	0.44
2:B:144:MET:HE3	2:B:144:MET:HB3	1.89	0.44
1:A:588:LYS:O	1:A:593:ASN:ND2	2.51	0.43
1:A:171:ARG:NE	7:A:922:HOH:O	2.51	0.43
1:A:647:MET:HE3	1:A:647:MET:HB2	1.86	0.43
1:A:670:ARG:HA	1:A:670:ARG:HD2	1.83	0.43
1:A:114:ARG:O	1:A:126:ASP:HA	2.19	0.43
1:A:688:ARG:NH2	7:A:923:HOH:O	2.52	0.43
2:B:117:THR:HG23	2:B:120:GLU:H	1.84	0.43
1:A:675:LEU:HD12	1:A:692:LEU:HD12	2.01	0.42
1:C:574:TYR:HB3	1:C:593:ASN:O	2.19	0.42
1:C:242:ASN:HB2	1:C:288:HIS:NE2	2.35	0.42
3:C:801:ATP:H5'1	3:C:801:ATP:C8	2.49	0.42
4:C:802:ADP:H8	2:D:96:GLY:O	2.04	0.41
1:A:169:ALA:HA	1:A:227:HIS:O	2.19	0.41
1:A:177:ASP:OD1	1:A:178:ARG:N	2.53	0.41
2:D:130:ILE:HD12	2:D:140:GLU:HG2	2.02	0.41
1:A:289:THR:O	1:A:301:GLY:HA2	2.20	0.41
1:A:592:GLU:CD	1:A:592:GLU:H	2.27	0.41
1:C:107:ILE:O	1:C:132:LYS:NZ	2.52	0.41
1:C:162:LYS:H	1:C:162:LYS:HG3	1.68	0.40
1:C:243:SER:HA	1:C:307:LEU:HD13	2.03	0.40
2:D:108:VAL:O	2:D:112:LEU:HG	2.21	0.40
1:C:670:ARG:HD2	1:C:670:ARG:HA	1.82	0.40
1:A:210:MET:HE3	1:A:210:MET:HB3	1.96	0.40
1:A:215:ILE:HG13	1:A:226:PHE:HB2	2.04	0.40
1:A:241:SER:OG	1:A:245:PHE:HB3	2.20	0.40
1:C:178:ARG:HE	1:C:225:LEU:HD11	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	486/531 (92%)	476 (98%)	10 (2%)	0	100	100
1	C	478/531 (90%)	462 (97%)	15 (3%)	1 (0%)	43	50
2	B	69/148 (47%)	69 (100%)	0	0	100	100
2	D	65/148 (44%)	65 (100%)	0	0	100	100
All	All	1098/1358 (81%)	1072 (98%)	25 (2%)	1 (0%)	48	56

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	512	LEU

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	396/437 (91%)	389 (98%)	7 (2%)	51	66
1	C	392/437 (90%)	385 (98%)	7 (2%)	51	66
2	B	59/126 (47%)	57 (97%)	2 (3%)	32	42
2	D	55/126 (44%)	54 (98%)	1 (2%)	51	66
All	All	902/1126 (80%)	885 (98%)	17 (2%)	50	64

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	LYS
1	A	232	ILE
1	A	274	ASP
1	A	344	GLN
1	A	514	LEU
1	A	592	GLU
1	A	638	LEU
2	B	94	LYS
2	B	148	LYS
1	C	128	GLU
1	C	232	ILE
1	C	274	ASP
1	C	334	ASP
1	C	520	LYS
1	C	592	GLU
1	C	603	LYS
2	D	143	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	136	GLN
1	A	206	GLN
1	A	286	GLN
1	A	722	GLN
1	C	576	GLN
1	C	642	ASN
1	C	710	ASN
1	C	722	GLN
2	D	111	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	TPO	C	348	1	8,10,11	1.28	2 (25%)	10,14,16	1.83	1 (10%)
1	TPO	A	348	1	8,10,11	1.77	3 (37%)	10,14,16	1.72	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	C	348	1	-	1/9/11/13	-
1	TPO	A	348	1	-	4/9/11/13	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	348	TPO	P-O1P	3.49	1.61	1.50
1	A	348	TPO	P-O2P	2.10	1.62	1.54
1	C	348	TPO	P-OG1	2.10	1.63	1.59
1	A	348	TPO	P-OG1	2.05	1.63	1.59
1	C	348	TPO	P-O2P	2.00	1.62	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	348	TPO	P-OG1-CB	-5.06	109.59	123.33
1	A	348	TPO	P-OG1-CB	-4.49	111.12	123.33

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	348	TPO	O-C-CA-CB
1	A	348	TPO	CB-OG1-P-O2P
1	A	348	TPO	CB-OG1-P-O3P
1	C	348	TPO	O-C-CA-CB
1	A	348	TPO	CB-OG1-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ADP	A	802	-	28,29,29	0.74	1 (3%)	43,45,45	0.88	2 (4%)
3	ATP	A	801	-	32,33,33	1.14	1 (3%)	48,52,52	0.82	2 (4%)
4	ADP	C	802	-	28,29,29	0.70	0	43,45,45	0.92	1 (2%)
3	ATP	C	801	-	32,33,33	1.19	1 (3%)	48,52,52	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	A	802	-	-	0/16/32/32	0/3/3/3
3	ATP	A	801	-	-	5/22/38/38	0/3/3/3
4	ADP	C	802	-	-	0/16/32/32	0/3/3/3
3	ATP	C	801	-	-	6/22/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	801	ATP	PB-O3B	-6.41	1.52	1.59
3	A	801	ATP	PB-O3B	-6.11	1.52	1.59

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	802	ADP	PB-O1B	2.28	1.57	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	801	ATP	O3A-PA-O1A	-3.08	101.45	110.70
4	C	802	ADP	O2'-C2'-C1'	-2.63	101.06	110.10
4	A	802	ADP	O2'-C2'-C1'	-2.61	101.12	110.10
4	A	802	ADP	O3B-PB-O2B	2.25	116.25	107.80
3	A	801	ATP	O2A-PA-O5'	2.24	117.70	107.57

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	ATP	C5'-O5'-PA-O1A
3	A	801	ATP	C5'-O5'-PA-O3A
3	C	801	ATP	C5'-O5'-PA-O1A
3	C	801	ATP	C5'-O5'-PA-O3A
3	C	801	ATP	PG-O3B-PB-O2B
3	A	801	ATP	C4'-C5'-O5'-PA
3	A	801	ATP	C5'-O5'-PA-O2A
3	C	801	ATP	C4'-C5'-O5'-PA
3	A	801	ATP	PB-O3A-PA-O2A
3	C	801	ATP	PG-O3B-PB-O1B
3	C	801	ATP	PA-O3A-PB-O2B

There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	802	ADP	1	0
3	C	801	ATP	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring




in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




## Ligand ATP A 801



## Ligand ADP C 802





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	496/531 (93%)	0.56	39 (7%)	18 16	17, 32, 66, 86	0
1	C	488/531 (91%)	0.49	37 (7%)	20 17	15, 32, 64, 81	2 (0%)
2	B	71/148 (47%)	0.61	4 (5%)	30 27	23, 39, 65, 72	0
2	D	67/148 (45%)	0.72	2 (2%)	52 50	24, 40, 59, 70	0
All	All	1122/1358 (82%)	0.54	82 (7%)	21 18	15, 33, 65, 86	2 (0%)

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	340	THR	5.1
1	C	647	MET	5.0
1	C	250	ASN	5.0
1	A	648	THR	4.9
2	D	81	SER	4.7
1	A	512	LEU	4.5
1	C	661	ASP	4.4
1	A	357	CYS	4.3
1	A	513	ASP	4.2
1	A	77	ASN	4.2
2	B	78	ASP	4.2
1	A	514	LEU	4.0
1	C	349	ILE	4.0
1	C	247	ARG	3.9
1	A	495	HIS	3.9
1	C	249	ASP	3.8
1	A	590	THR	3.8
1	C	339	ASN	3.8
1	C	316	ARG	3.8
1	C	514	LEU	3.8
1	A	344	GLN	3.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	245	PHE	3.6
1	C	540	CYS	3.5
1	C	80	HIS	3.5
1	A	515	GLU	3.5
1	C	547	ASP	3.2
1	C	518	ILE	3.2
1	A	725	GLU	3.2
1	A	336	VAL	3.1
1	C	669	ALA	3.1
1	A	248	ASP	3.1
1	A	346	ALA	3.1
1	C	660	GLN	3.0
1	A	81	PHE	3.0
1	C	590	THR	3.0
1	C	513	ASP	2.9
1	C	346	ALA	2.9
1	C	95	MET	2.9
1	A	342	LEU	2.9
1	C	246	VAL	2.9
1	A	340	THR	2.8
1	C	724	GLU	2.8
1	C	511	ALA	2.8
1	A	333	ARG	2.8
1	A	94	HIS	2.7
1	A	703	ALA	2.6
1	A	356	LYS	2.6
1	C	702	GLU	2.5
1	A	79	PHE	2.5
1	C	79	PHE	2.5
1	A	334	ASP	2.5
1	C	544	GLU	2.4
1	C	515	GLU	2.4
1	A	511	ALA	2.4
1	A	647	MET	2.4
1	C	179	ASP	2.4
1	C	545	GLU	2.3
1	C	353	THR	2.3
1	C	701	MET	2.3
1	C	519	GLY	2.3
1	A	345	SER	2.3
1	A	636	GLU	2.3
1	C	591	GLU	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	357	CYS	2.2
1	A	337	ASN	2.2
1	A	547	ASP	2.2
2	B	148	LYS	2.1
1	A	80	HIS	2.1
1	A	338	GLN	2.1
2	D	117	THR	2.1
1	A	541	GLU	2.1
2	B	81	SER	2.1
2	B	147	ALA	2.1
1	A	662	GLU	2.1
1	C	350	LEU	2.1
1	A	273	VAL	2.1
1	A	591	GLU	2.0
1	C	512	LEU	2.0
1	A	316	ARG	2.0
1	A	540	CYS	2.0
1	A	339	ASN	2.0
1	A	542	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	A	348	11/12	0.92	0.12	30,39,50,55	0
1	TPO	C	348	11/12	0.93	0.13	43,50,59,60	0

## 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ATP	C	801	31/31	0.88	0.13	48,48,48,48	0
3	ATP	A	801	31/31	0.89	0.12	46,46,46,46	0
6	CA	B	202	1/1	0.95	0.05	36,36,36,36	0
4	ADP	C	802	27/27	0.97	0.07	24,24,24,24	0
4	ADP	A	802	27/27	0.97	0.07	21,21,21,21	0
6	CA	D	201	1/1	0.97	0.04	22,22,22,22	0
6	CA	D	202	1/1	0.97	0.04	35,35,35,35	0
6	CA	B	201	1/1	0.98	0.05	24,24,24,24	0
5	ZN	A	803	1/1	0.99	0.03	28,28,28,28	0
5	ZN	C	803	1/1	1.00	0.02	31,31,31,31	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








**Electron density around ATP A 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ADP C 802:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around ADP A 802:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.