



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 07:26 PM UTC

PDB ID : 8PMV / pdb_00008pmv
Title : transcription factor BARHL2 bound to TAAGC DNA sequence
Authors : Morgunova, E.; Popov, A.; Yin, Y.; Taipale, J.
Deposited on : 2023-06-29
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	62	<div> <div>8%</div> <div>79%</div> <div>19%</div> <div>.</div> </div>
1	D	62	<div> <div>16%</div> <div>66%</div> <div>27%</div> <div>..</div> </div>
1	G	62	<div> <div>6%</div> <div>82%</div> <div>18%</div> </div>
1	J	62	<div> <div>18%</div> <div>76%</div> <div>18%</div> <div>• 5%</div> </div>
2	B	12	<div> <div>67%</div> <div>33%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	12	
2	H	12	
2	K	12	
3	C	12	
3	F	12	
3	I	12	
3	L	12	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4148 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BarH-like 2 homeobox protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	62	Total	C	N	O	S	0	0	0
			530	328	106	95	1			
1	D	60	Total	C	N	O	S	0	0	0
			516	319	103	93	1			
1	G	62	Total	C	N	O	S	0	0	0
			530	328	106	95	1			
1	J	59	Total	C	N	O	S	0	0	0
			509	315	102	91	1			

- Molecule 2 is a DNA chain called DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	P	0	0	0
			243	116	45	71	11			
2	E	12	Total	C	N	O	P	0	0	0
			243	117	45	70	11			
2	H	12	Total	C	N	O	P	0	0	0
			244	117	45	71	11			
2	K	12	Total	C	N	O	P	0	0	0
			244	117	45	71	11			

- Molecule 3 is a DNA chain called DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	12	Total	C	N	O	P	0	0	0
			242	116	46	69	11			
3	F	12	Total	C	N	O	P	0	0	0
			242	116	46	69	11			
3	I	12	Total	C	N	O	P	0	0	0
			242	116	46	69	11			
3	L	12	Total	C	N	O	P	0	0	0
			242	116	46	69	11			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	12	Total O 12 12	0	0
4	B	10	Total O 10 10	0	0
4	C	7	Total O 7 7	0	0
4	D	15	Total O 15 15	0	0
4	E	6	Total O 6 6	0	0
4	F	10	Total O 10 10	0	0
4	G	10	Total O 10 10	0	0
4	H	5	Total O 5 5	0	0
4	I	23	Total O 23 23	0	0
4	J	14	Total O 14 14	0	0
4	K	3	Total O 3 3	0	0
4	L	6	Total O 6 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: BarH-like 2 homeobox protein




- Molecule 1: BarH-like 2 homeobox protein



- Molecule 1: BarH-like 2 homeobox protein




- Molecule 1: BarH-like 2 homeobox protein



- Molecule 2: DNA



- Molecule 2: DNA



● Molecule 2: DNA



● Molecule 2: DNA



● Molecule 3: DNA



● Molecule 3: DNA



● Molecule 3: DNA



● Molecule 3: DNA



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.08Å 133.77Å 64.99Å 90.00° 92.74° 90.00°	Depositor
Resolution (Å)	46.63 – 2.10 46.63 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.63-2.10) 99.8 (46.63-2.10)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.251 , 0.278 0.256 , 0.286	Depositor DCC
R_{free} test set	2041 reflections (5.23%)	wwPDB-VP
Wilson B-factor (Å ²)	52.6	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.054 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4148	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.11	0/539	1.48	1/723 (0.1%)
1	D	1.01	0/525	1.54	3/704 (0.4%)
1	G	1.05	0/539	1.53	1/723 (0.1%)
1	J	0.92	0/518	1.48	1/694 (0.1%)
2	B	0.85	0/270	1.42	1/413 (0.2%)
2	E	0.75	0/272	1.23	1/419 (0.2%)
2	H	0.88	0/273	1.44	3/420 (0.7%)
2	K	0.79	0/273	1.24	1/420 (0.2%)
3	C	0.80	0/271	1.57	5/416 (1.2%)
3	F	0.76	0/271	1.49	4/416 (1.0%)
3	I	0.75	0/271	1.48	4/416 (1.0%)
3	L	0.73	0/271	1.51	4/416 (1.0%)
All	All	0.91	0/4293	1.46	29/6180 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	21	DA	O3'-P-O5'	-8.15	91.77	104.00
1	D	280	TYR	CB-CA-C	7.72	123.17	110.81
3	I	23	DC	C2'-C3'-O3'	-7.64	100.04	111.50
3	C	23	DC	C2'-C3'-O3'	-7.57	100.14	111.50
1	A	280	TYR	CB-CA-C	7.48	123.16	110.74
3	L	22	DG	C4'-C3'-O3'	-7.34	98.99	110.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	280	TYR	CB-CA-C	7.29	122.83	110.74
3	F	23	DC	C2'-C3'-O3'	-7.06	100.91	111.50
3	F	21	DA	C4'-C3'-O3'	-6.99	99.52	110.00
1	G	280	TYR	CB-CA-C	6.82	122.06	110.74
3	L	21	DA	C4'-C3'-O3'	-6.68	99.98	110.00
3	I	22	DG	C4'-C3'-O3'	-6.61	100.09	110.00
1	D	285	THR	CA-CB-OG1	-6.20	100.30	109.60
3	I	18	DC	P-O5'-C5'	-5.95	111.08	120.00
2	K	8	DC	P-O3'-C3'	-5.93	111.30	120.20
2	E	8	DC	P-O3'-C3'	-5.93	111.31	120.20
3	F	23	DC	O3'-P-O5'	-5.88	95.19	104.00
2	H	4	DT	C2'-C3'-O3'	-5.86	102.71	111.50
3	L	16	DC	P-O3'-C3'	-5.82	111.47	120.20
3	C	21	DA	O3'-P-O5'	-5.82	95.27	104.00
3	F	22	DG	C4'-C3'-O3'	-5.54	101.68	110.00
3	C	21	DA	C4'-C3'-O3'	-5.43	101.85	110.00
3	I	21	DA	C4'-C3'-O3'	-5.38	101.93	110.00
1	D	280	TYR	N-CA-C	-5.31	104.74	111.11
2	H	2	DG	P-O5'-C5'	-5.28	112.08	120.00
3	C	18	DC	P-O5'-C5'	-5.25	112.12	120.00
2	B	2	DG	C4'-C3'-O3'	-5.19	102.21	110.00
2	H	2	DG	C4'-C3'-O3'	-5.15	102.27	110.00
3	C	22	DG	C4'-C3'-O3'	-5.06	102.41	110.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	282	ASN	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	530	0	533	12	0
1	D	516	0	516	13	0
1	G	530	0	533	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	J	509	0	509	6	0
2	B	243	0	133	3	0
2	E	243	0	134	3	0
2	H	244	0	137	2	0
2	K	244	0	137	4	0
3	C	242	0	136	2	0
3	F	242	0	136	5	0
3	I	242	0	136	1	0
3	L	242	0	136	3	0
4	A	12	0	0	4	0
4	B	10	0	0	0	0
4	C	7	0	0	0	0
4	D	15	0	0	1	0
4	E	6	0	0	0	0
4	F	10	0	0	1	0
4	G	10	0	0	1	0
4	H	5	0	0	0	0
4	I	23	0	0	3	0
4	J	14	0	0	0	0
4	K	3	0	0	0	0
4	L	6	0	0	1	0
All	All	4148	0	3176	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:18:DC:O5'	4:F:101:HOH:O	1.76	1.00
2:B:11:DT:C3'	2:B:11:DT:C5'	2.56	0.83
1:G:290:GLN:NE2	4:G:301:HOH:O	2.12	0.82
1:G:262:ARG:HD2	1:G:273:ASP:OD1	1.80	0.81
1:D:236:ARG:NH2	2:E:4:DT:O2	2.14	0.81
1:J:236:ARG:NH2	2:K:4:DT:O2	2.14	0.81
1:J:262:ARG:HD2	1:J:273:ASP:OD1	1.82	0.79
1:D:288:LYS:NZ	3:F:17:DG:OP2	2.14	0.78
1:D:262:ARG:NH1	3:F:14:DA:OP1	2.19	0.75
1:A:256:TYR:CE2	1:D:291:THR:HA	2.29	0.68
2:K:1:DC:H2''	2:K:2:DG:C8	2.30	0.66
2:B:11:DT:C5'	2:B:11:DT:O4'	2.43	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:1:DC:H2''	2:E:2:DG:C8	2.36	0.59
1:A:248:GLU:OE1	4:A:301:HOH:O	2.16	0.59
1:D:253:ARG:NH1	4:D:302:HOH:O	2.36	0.58
1:D:277:LYS:NZ	3:F:14:DA:OP2	2.36	0.58
1:A:281:GLN:NE2	3:C:15:DC:H2'	2.21	0.56
1:J:277:LYS:NZ	3:L:14:DA:OP2	2.30	0.56
1:A:252:GLU:HG3	4:A:310:HOH:O	2.11	0.51
1:A:248:GLU:HG3	4:A:310:HOH:O	2.11	0.50
1:A:281:GLN:HE22	3:C:15:DC:H2'	1.76	0.50
1:A:274:THR:CG2	4:I:101:HOH:O	2.60	0.49
1:J:265:LEU:CD2	1:J:269:LEU:HD22	2.43	0.49
1:J:246:GLN:HB2	1:J:269:LEU:HD21	1.95	0.48
1:D:248:GLU:O	1:D:252:GLU:HG3	2.14	0.48
1:A:256:TYR:CZ	1:D:291:THR:HA	2.48	0.48
1:A:274:THR:HG22	4:I:101:HOH:O	2.13	0.47
1:G:249:ARG:O	1:G:252:GLU:HG3	2.13	0.47
1:D:290:GLN:O	1:D:291:THR:C	2.59	0.46
1:A:260:GLN:HA	1:A:263:MET:HE2	1.98	0.45
1:A:281:GLN:NE2	4:A:302:HOH:O	2.37	0.45
2:K:8:DC:H2''	2:K:9:DG:C8	2.52	0.45
1:D:263:MET:HG2	1:D:273:ASP:OD2	2.17	0.44
1:D:234:LYS:HE2	1:D:235:ALA:O	2.18	0.44
3:I:23:DC:OP1	4:I:101:HOH:O	2.21	0.43
3:L:24:DG:O3'	4:L:101:HOH:O	2.20	0.43
1:D:288:LYS:CE	3:F:17:DG:OP2	2.68	0.42
2:B:8:DC:H2''	2:B:9:DG:C8	2.55	0.42
1:G:236:ARG:NH2	2:H:4:DT:O2	2.53	0.42
2:E:8:DC:H2''	2:E:9:DG:C8	2.54	0.41
1:J:236:ARG:NH2	3:L:22:DG:N2	2.69	0.41
2:K:1:DC:H2''	2:K:2:DG:N7	2.36	0.41
1:A:255:LYS:HB2	1:D:287:TRP:CE2	2.56	0.41
2:H:8:DC:H2''	2:H:9:DG:C8	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	60/62 (97%)	60 (100%)	0	0	100	100
1	D	58/62 (94%)	57 (98%)	1 (2%)	0	100	100
1	G	60/62 (97%)	60 (100%)	0	0	100	100
1	J	57/62 (92%)	56 (98%)	1 (2%)	0	100	100
All	All	235/248 (95%)	233 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	56/56 (100%)	52 (93%)	4 (7%)	13	11
1	D	55/56 (98%)	50 (91%)	5 (9%)	9	6
1	G	56/56 (100%)	52 (93%)	4 (7%)	13	11
1	J	54/56 (96%)	49 (91%)	5 (9%)	8	6
All	All	221/224 (99%)	203 (92%)	18 (8%)	11	8

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	231	LYS
1	A	244	LEU
1	A	270	ASN
1	A	274	THR
1	D	234	LYS
1	D	257	LEU
1	D	265	LEU
1	D	270	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	290	GLN
1	G	231	LYS
1	G	244	LEU
1	G	255	LYS
1	G	270	ASN
1	J	244	LEU
1	J	252	GLU
1	J	263	MET
1	J	265	LEU
1	J	270	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	245	ASN
1	A	246	GLN
1	A	281	GLN
1	G	243	GLN
1	G	290	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	62/62 (100%)	0.93	5 (8%) 18 19	37, 57, 82, 105	0
1	D	60/62 (96%)	1.16	10 (16%) 4 4	43, 64, 90, 108	0
1	G	62/62 (100%)	0.70	4 (6%) 25 26	41, 58, 83, 97	0
1	J	59/62 (95%)	1.32	11 (18%) 3 3	58, 91, 111, 122	0
2	B	12/12 (100%)	-0.10	0 100 100	45, 61, 77, 83	0
2	E	12/12 (100%)	0.36	1 (8%) 17 18	58, 67, 93, 106	0
2	H	12/12 (100%)	-0.37	0 100 100	43, 61, 67, 69	0
2	K	12/12 (100%)	0.23	0 100 100	54, 70, 80, 81	0
3	C	12/12 (100%)	-0.42	0 100 100	53, 58, 63, 63	0
3	F	12/12 (100%)	0.21	0 100 100	50, 67, 97, 104	0
3	I	12/12 (100%)	-0.46	0 100 100	48, 54, 60, 61	0
3	L	12/12 (100%)	0.29	0 100 100	62, 69, 77, 80	0
All	All	339/344 (98%)	0.72	31 (9%) 15 15	37, 64, 101, 122	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	291	THR	5.4
1	G	292	ALA	4.4
1	J	232	PRO	4.0
1	J	287	TRP	3.6
1	A	273	ASP	3.6
1	J	237	THR	3.2
1	J	256	TYR	2.9
1	J	259	VAL	2.8
1	J	239	PHE	2.8
1	D	288	LYS	2.8
1	J	265	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	237	THR	2.8
1	A	267	ALA	2.7
1	D	238	ALA	2.6
1	A	289	ARG	2.5
1	G	287	TRP	2.5
1	J	238	ALA	2.5
1	A	290	GLN	2.4
1	A	263	MET	2.4
1	J	289	ARG	2.4
1	D	273	ASP	2.3
1	D	235	ALA	2.3
1	G	273	ASP	2.2
1	J	257	LEU	2.2
1	D	259	VAL	2.1
1	D	254	GLN	2.1
1	D	244	LEU	2.1
1	D	270	ASN	2.1
1	G	291	THR	2.1
2	E	1	DC	2.1
1	J	271	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.