Full wwPDB X-ray Structure Validation Report ⓘ

Nov 16, 2026 – 02:37 PM EST

PDB ID	:	11EW / pdb_000011ew
Title	:	Crystal structure of C2-DHR1 domains of DOCK5 in complex with a small molecule DEL hit
Authors	:	Deng, Y.; Liu, W.; Gajiwala, K.S.
Deposited on	:	2026-02-19
Resolution	:	2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	13299 (2.04-2.00)
Clashscore	190562	1022 (2.02-2.02)
Ramachandran outliers	187476	1014 (2.02-2.02)
Sidechain outliers	187428	1014 (2.02-2.02)
RSRZ outliers	180081	13314 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	5% 75% 15% • 9%
1	B	419	6% 70% 17% • 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6238 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dedicator of cytokinesis protein 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	382	Total	C	N	O	S	0	0	0
			3055	1944	523	570	18			
1	B	374	Total	C	N	O	S	0	1	0
			2995	1904	513	559	19			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	219	GLY	-	expression tag	UNP Q9H7D0
A	220	SER	-	expression tag	UNP Q9H7D0
B	219	GLY	-	expression tag	UNP Q9H7D0
B	220	SER	-	expression tag	UNP Q9H7D0

- Molecule 2 is 1-{(1s,4s)-4-[(3-chloro-4-ethoxyphenyl)carbamoyl]cyclohexyl}-N-methyl-2-(pyrimidin-5-yl)-1H-1,3-benzimidazole-5-carboxamide (CCD ID: A1C8N) (formula: $C_{28}H_{29}ClN_6O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			38	28	1	6	3		
2	B	1	Total	C	Cl	N	O	0	0
			38	28	1	6	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	56	Total	O	0	0
			56	56		
3	B	56	Total	O	0	0
			56	56		


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dedicator of cytokinesis protein 5

Chain A:






• Molecule 1: Dedicator of cytokinesis protein 5

Chain B:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	50.86Å 80.61Å 101.51Å 90.00° 95.55° 90.00°	Depositor
Resolution (Å)	33.68 – 2.03 33.68 – 2.03	Depositor EDS
% Data completeness (in resolution range)	53.3 (33.68-2.03) 53.3 (33.68-2.03)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.25 (at 2.03Å)	Xtriage
Refinement program	BUSTER 2.11.8	Depositor
R, R_{free}	0.222 , 0.296 0.217 , 0.288	Depositor DCC
R_{free} test set	1478 reflections (2.80%)	wwPDB-VP
Wilson B-factor (Å^2)	53.5	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 43.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	6238	wwPDB-VP
Average B, all atoms (Å^2)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.98% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1C8N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.74	0/3112	1.07	1/4191 (0.0%)
1	B	0.75	0/3052	1.08	9/4111 (0.2%)
All	All	0.74	0/6164	1.08	10/8302 (0.1%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	474	ASP	CA-CB-CG	7.42	120.02	112.60
1	B	263	GLY	CA-C-N	5.92	128.49	120.38
1	B	263	GLY	C-N-CA	5.92	128.49	120.38
1	B	417	PHE	CA-C-N	5.57	129.50	120.60
1	B	417	PHE	C-N-CA	5.57	129.50	120.60
1	B	309	LYS	CA-C-N	5.12	131.32	121.54
1	B	309	LYS	C-N-CA	5.12	131.32	121.54
1	B	416	ASN	CA-C-N	5.12	129.36	121.53
1	B	416	ASN	C-N-CA	5.12	129.36	121.53
1	B	474	ASP	CA-CB-CG	5.05	117.65	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3055	0	3069	36	0
1	B	2995	0	2995	41	0
2	A	38	0	0	0	0
2	B	38	0	0	0	0
3	A	56	0	0	0	0
3	B	56	0	0	2	0
All	All	6238	0	6064	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:523:THR:HA	1:A:555:MET:HG3	1.07	1.04
1:B:476:GLU:HG2	1:B:583:LYS:HZ1	1.26	0.96
1:A:523:THR:HA	1:A:555:MET:CG	2.01	0.85
1:A:476:GLU:HA	1:A:583:LYS:HZ1	1.43	0.81
1:B:356:TYR:HD2	1:B:385:LYS:HZ2	1.27	0.81
1:A:229:LYS:NZ	1:A:343:GLN:HG2	1.98	0.79
1:A:229:LYS:HZ1	1:A:343:GLN:HG2	1.51	0.74
1:A:476:GLU:HA	1:A:583:LYS:NZ	2.04	0.72
1:A:476:GLU:HG2	1:A:583:LYS:NZ	2.03	0.72
1:A:476:GLU:HG2	1:A:583:LYS:HZ1	1.55	0.71
1:B:476:GLU:HG2	1:B:583:LYS:NZ	2.02	0.71
1:B:476:GLU:CG	1:B:583:LYS:HZ1	2.01	0.70
1:A:523:THR:CA	1:A:555:MET:HG3	2.04	0.68
1:B:238:ASP:HB3	1:B:306:MET:HE3	1.81	0.63
1:A:476:GLU:CA	1:A:583:LYS:HZ1	2.09	0.62
1:B:465:ASN:H	1:B:533:HIS:HD2	1.47	0.61
1:B:517:ILE:HG23	1:B:521:GLU:HB2	1.83	0.60
1:B:314:HIS:HE1	1:B:316:CYS:SG	2.24	0.60
1:B:567:HIS:HE1	1:B:624:THR:OG1	1.86	0.59
1:A:244:ALA:HB2	1:A:257:ASN:HD22	1.66	0.59
1:B:517:ILE:HG22	1:B:522:VAL:HG23	1.84	0.59
1:B:555:MET:HE2	1:B:561:THR:HG23	1.85	0.59
1:B:356:TYR:HD2	1:B:385:LYS:NZ	2.01	0.58
1:B:324:VAL:HG22	1:B:353:MET:HE1	1.85	0.58
1:B:314:HIS:CE1	1:B:316:CYS:SG	2.99	0.56
1:A:526:HIS:HE1	1:A:585:TYR:OH	1.90	0.55
1:B:229:LYS:O	1:B:230:ASN:HB2	2.08	0.54
1:B:556:ASN:HB3	1:B:557:PRO:HD2	1.89	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330:THR:HG23	1:A:334:HIS:CE1	2.43	0.53
1:B:379:LEU:O	1:B:383:LEU:HB2	2.08	0.53
1:A:248:PRO:HD2	1:A:293:ARG:O	2.09	0.53
1:B:456:ASP:CG	1:B:464:LYS:HZ2	2.17	0.53
1:A:303:VAL:HG21	1:A:306:MET:HE3	1.91	0.52
1:A:523:THR:HG22	1:A:555:MET:HE3	1.91	0.51
1:A:578:LYS:HB3	1:A:584:PHE:CD2	2.45	0.51
1:B:356:TYR:HA	1:B:359:GLN:HB2	1.91	0.51
1:B:229:LYS:HG3	1:B:343:GLN:HE21	1.75	0.51
1:A:227:ASN:HB3	1:A:402:SER:HB2	1.93	0.51
1:A:446:ILE:HG12	1:A:626:ILE:HG12	1.94	0.49
1:A:356:TYR:CD1	1:A:385:LYS:NZ	2.80	0.48
1:A:231:PHE:CD1	1:A:399:LEU:HD23	2.48	0.48
1:B:356:TYR:HB2	1:B:385:LYS:HD2	1.95	0.48
1:B:494:SER:HB2	3:B:9150:HOH:O	2.13	0.48
1:B:584:PHE:O	1:B:587:THR:OG1	2.32	0.48
1:A:476:GLU:CG	1:A:583:LYS:HZ1	2.24	0.47
1:B:472:VAL:HG11	1:B:517:ILE:CD1	2.45	0.46
1:A:330:THR:CG2	1:A:334:HIS:CE1	2.98	0.46
1:A:479:LEU:HD21	1:A:494:SER:HB3	1.98	0.46
1:B:578:LYS:HB3	1:B:584:PHE:CD2	2.51	0.46
1:B:244:ALA:HB2	1:B:257:ASN:HD22	1.81	0.46
1:A:327:MET:HE1	1:A:403:LEU:HD11	1.98	0.45
1:B:231:PHE:CD1	1:B:399:LEU:HD23	2.51	0.45
1:A:519:ILE:HG13	1:A:631:LEU:HD11	1.98	0.45
1:B:472:VAL:HG11	1:B:517:ILE:HD11	1.99	0.45
1:B:282:THR:HG21	1:B:410:LEU:HD13	1.98	0.45
1:B:356:TYR:CD2	1:B:385:LYS:HG3	2.53	0.44
1:B:243:MET:HB3	1:B:258:TYR:HB3	2.00	0.43
1:A:327:MET:CE	1:A:403:LEU:HD11	2.47	0.43
1:A:484:ILE:HG21	1:A:496:TYR:HB2	2.00	0.43
1:A:470:MET:HB3	1:A:496:TYR:HB3	1.99	0.42
1:B:308:LEU:HD22	1:B:312:LYS:HA	2.01	0.42
1:B:497:LYS:HE3	1:B:497:LYS:HB2	1.86	0.42
1:B:314:HIS:HD2	3:B:9148:HOH:O	2.03	0.42
1:A:229:LYS:HZ3	1:A:229:LYS:HG3	1.60	0.42
1:A:469:THR:HG23	1:A:497:LYS:HD3	2.01	0.41
1:B:476:GLU:CB	1:B:583:LYS:HZ1	2.32	0.41
1:A:476:GLU:CB	1:A:583:LYS:HZ1	2.32	0.41
1:A:530:THR:OG1	1:A:532:ARG:NH1	2.54	0.41
1:B:243:MET:HG2	1:B:296:LEU:HD11	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:567:HIS:CE1	1:B:624:THR:OG1	2.71	0.41
1:A:446:ILE:HB	1:A:515:VAL:HB	2.02	0.41
1:B:521:GLU:O	1:B:525:CYS:SG	2.79	0.41
1:B:240:GLU:OE2	1:B:316:CYS:SG	2.79	0.41
1:B:476:GLU:CB	1:B:583:LYS:NZ	2.85	0.40
1:A:228:PHE:HZ	1:A:231:PHE:HB2	1.86	0.40
1:A:577:LYS:HE3	1:B:438:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	372/419 (89%)	351 (94%)	20 (5%)	1 (0%)	36 31
1	B	365/419 (87%)	338 (93%)	25 (7%)	2 (0%)	24 17
All	All	737/838 (88%)	689 (94%)	45 (6%)	3 (0%)	30 22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	389	ALA
1	B	310	GLU
1	B	390	LYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	340/375 (91%)	312 (92%)	28 (8%)	10 5
1	B	333/375 (89%)	295 (89%)	38 (11%)	5 1
All	All	673/750 (90%)	607 (90%)	66 (10%)	7 2

All (66) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	229	LYS
1	A	252	THR
1	A	270	GLU
1	A	271	ILE
1	A	272	GLU
1	A	274	LEU
1	A	284	LEU
1	A	290	ILE
1	A	295	SER
1	A	333	ILE
1	A	341	GLU
1	A	362	LEU
1	A	379	LEU
1	A	383	LEU
1	A	385	LYS
1	A	401	VAL
1	A	415	LYS
1	A	423	ARG
1	A	454	GLU
1	A	481	GLU
1	A	491	GLU
1	A	505	LYS
1	A	519	ILE
1	A	520	GLU
1	A	583	LYS
1	A	600	LEU
1	A	617	LYS
1	A	625	LEU
1	B	229	LYS
1	B	243	MET
1	B	261	ARG
1	B	264	SER
1	B	269	LYS
1	B	270	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	271	ILE
1	B	272	GLU
1	B	274	LEU
1	B	308	LEU
1	B	310	GLU
1	B	312	LYS
1	B	315	THR
1	B	324	VAL
1	B	340	GLU
1	B	342	LYS
1	B	351	ILE
1	B	358	ARG
1	B	359	GLN
1	B	376	ASN
1	B	382	VAL
1	B	391	GLU
1	B	401	VAL
1	B	413	VAL
1	B	436	ILE
1	B	454	GLU
1	B	506	GLN
1	B	516	SER
1	B	517	ILE
1	B	522	VAL
1	B	530	THR
1	B	555	MET
1	B	558	ASP
1	B	561	THR
1	B	577	LYS
1	B	580	GLU
1	B	587	THR
1	B	600	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	227	ASN
1	A	257	ASN
1	A	278	GLN
1	A	314	HIS
1	A	334	HIS
1	A	343	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	349	GLN
1	A	350	GLN
1	A	384	ASN
1	A	473	HIS
1	A	526	HIS
1	A	533	HIS
1	A	556	ASN
1	A	567	HIS
1	B	227	ASN
1	B	257	ASN
1	B	278	GLN
1	B	314	HIS
1	B	343	GLN
1	B	376	ASN
1	B	397	GLN
1	B	414	GLN
1	B	503	GLN
1	B	506	GLN
1	B	526	HIS
1	B	533	HIS
1	B	556	ASN
1	B	567	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	A1C8N	B	9001	-	42,42,42	1.00	2 (4%)	59,59,59	1.08	3 (5%)
2	A1C8N	A	9001	-	42,42,42	0.90	1 (2%)	59,59,59	1.22	5 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1C8N	B	9001	-	-	4/25/35/35	0/5/5/5
2	A1C8N	A	9001	-	-	6/25/35/35	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	9001	A1C8N	C28-N30	3.20	1.37	1.33
2	B	9001	A1C8N	C15-N19	2.47	1.52	1.48
2	A	9001	A1C8N	C28-N30	2.08	1.36	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	9001	A1C8N	C25-C24-C23	3.23	122.99	119.25
2	A	9001	A1C8N	C18-C11-C9	3.23	117.59	110.79
2	B	9001	A1C8N	C13-C14-C15	2.69	115.18	109.98
2	B	9001	A1C8N	C25-C24-C23	2.48	122.11	119.25
2	A	9001	A1C8N	C31-N30-C28	-2.33	117.35	121.80
2	A	9001	A1C8N	C14-C13-C11	2.30	115.22	111.18
2	B	9001	A1C8N	C2-O3-C4	-2.29	113.41	118.08
2	A	9001	A1C8N	C2-O3-C4	-2.15	113.68	118.08

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	9001	A1C8N	O29-C28-N30-C31
2	A	9001	A1C8N	C24-C28-N30-C31
2	B	9001	A1C8N	C24-C28-N30-C31
2	A	9001	A1C8N	C18-C11-C9-O10
2	A	9001	A1C8N	C18-C11-C9-N8
2	A	9001	A1C8N	O29-C28-N30-C31
2	B	9001	A1C8N	C18-C11-C9-N8
2	B	9001	A1C8N	C18-C11-C9-O10
2	A	9001	A1C8N	C6-C7-N8-C9
2	A	9001	A1C8N	C38-C7-N8-C9

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1C8N B 9001




Bond lengths



Bond angles




Torsions




Rings


Ligand A1C8N A 9001




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	382/419 (91%)	0.63	23 (6%)	27	27	32, 61, 90, 105	0
1	B	374/419 (89%)	0.65	25 (6%)	24	23	17, 61, 91, 104	1 (0%)
All	All	756/838 (90%)	0.64	48 (6%)	26	25	17, 61, 91, 105	1 (0%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	274	LEU	3.9
1	A	436	ILE	3.7
1	B	356	TYR	3.6
1	A	438	LEU	3.5
1	A	363	ILE	3.5
1	A	602	ALA	3.4
1	B	446	ILE	3.3
1	B	458	GLY	3.2
1	B	433	PHE	3.2
1	B	557	PRO	3.1
1	B	436	ILE	3.1
1	A	632	THR	2.8
1	B	437	ILE	2.7
1	A	413	VAL	2.6
1	B	352	ALA	2.6
1	B	271	ILE	2.6
1	B	290	ILE	2.5
1	A	252	THR	2.5
1	A	362	LEU	2.5
1	A	600	LEU	2.5
1	A	308	LEU	2.4
1	B	628	SER	2.4
1	B	629	THR	2.4
1	B	502	TYR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	582	ALA	2.3
1	A	554	LEU	2.3
1	A	433	PHE	2.3
1	A	333	ILE	2.3
1	B	413	VAL	2.2
1	B	590	GLY	2.2
1	B	410	LEU	2.2
1	B	418	SER	2.2
1	A	236	GLY	2.1
1	A	289	LEU	2.1
1	A	631	LEU	2.1
1	A	595	MET	2.1
1	A	382	VAL	2.1
1	B	546	ALA	2.1
1	A	410	LEU	2.1
1	B	289	LEU	2.1
1	B	440	GLY	2.1
1	B	527	ILE	2.1
1	B	438	LEU	2.1
1	A	243	MET	2.1
1	B	353	MET	2.1
1	A	357	ILE	2.1
1	B	444	ASN	2.0
1	A	566	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	A1C8N	A	9001	38/38	0.93	0.08	42,47,51,51	0
2	A1C8N	B	9001	38/38	0.93	0.07	37,42,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around A1C8N A 9001:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around A1C8N B 9001:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers

There are no such residues in this entry.