



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 30, 2026 – 10:13 AM EDT

PDB ID : 11TS / pdb_000011ts
Title : Crystal structure of apo alpha/beta-hydrolase macrolide esterase EstT from *Sphingobacterium faecium* (S126A mutant)
Authors : Hemmings, M.Z.; Blanchet, J.; Kelly, E.T.R.; Berghuis, A.M.
Deposited on : 2026-03-12
Resolution : 3.19 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1370 (3.20-3.20)
Clashscore	180529	1497 (3.20-3.20)
Ramachandran outliers	177936	1479 (3.20-3.20)
Sidechain outliers	177891	1478 (3.20-3.20)
RSRZ outliers	164620	1371 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	
1	B	311	
1	C	311	
1	D	311	
1	E	311	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 9644 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha/beta-hydrolase macrolide esterase EstT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	278	Total 2113	C 1341	N 354	O 411	S 7	0	0	0
1	B	284	Total 2105	C 1326	N 349	O 422	S 8	0	0	0
1	C	283	Total 2094	C 1323	N 358	O 407	S 6	0	0	0
1	D	250	Total 1829	C 1162	N 310	O 350	S 7	0	0	0
1	E	229	Total 1489	C 938	N 261	O 287	S 3	0	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	126	ALA	SER	engineered mutation	UNP A0ABU1EVH8
A	307	LEU	-	expression tag	UNP A0ABU1EVH8
A	308	GLU	-	expression tag	UNP A0ABU1EVH8
A	309	HIS	-	expression tag	UNP A0ABU1EVH8
A	310	HIS	-	expression tag	UNP A0ABU1EVH8
A	311	HIS	-	expression tag	UNP A0ABU1EVH8
B	126	ALA	SER	engineered mutation	UNP A0ABU1EVH8
B	307	LEU	-	expression tag	UNP A0ABU1EVH8
B	308	GLU	-	expression tag	UNP A0ABU1EVH8
B	309	HIS	-	expression tag	UNP A0ABU1EVH8
B	310	HIS	-	expression tag	UNP A0ABU1EVH8
B	311	HIS	-	expression tag	UNP A0ABU1EVH8
C	126	ALA	SER	engineered mutation	UNP A0ABU1EVH8
C	307	LEU	-	expression tag	UNP A0ABU1EVH8
C	308	GLU	-	expression tag	UNP A0ABU1EVH8
C	309	HIS	-	expression tag	UNP A0ABU1EVH8
C	310	HIS	-	expression tag	UNP A0ABU1EVH8
C	311	HIS	-	expression tag	UNP A0ABU1EVH8
D	126	ALA	SER	engineered mutation	UNP A0ABU1EVH8


Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	307	LEU	-	expression tag	UNP A0ABU1EVH8
D	308	GLU	-	expression tag	UNP A0ABU1EVH8
D	309	HIS	-	expression tag	UNP A0ABU1EVH8
D	310	HIS	-	expression tag	UNP A0ABU1EVH8
D	311	HIS	-	expression tag	UNP A0ABU1EVH8
E	126	ALA	SER	engineered mutation	UNP A0ABU1EVH8
E	307	LEU	-	expression tag	UNP A0ABU1EVH8
E	308	GLU	-	expression tag	UNP A0ABU1EVH8
E	309	HIS	-	expression tag	UNP A0ABU1EVH8
E	310	HIS	-	expression tag	UNP A0ABU1EVH8
E	311	HIS	-	expression tag	UNP A0ABU1EVH8

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	4	Total O 4 4	0	0
2	B	2	Total O 2 2	0	0
2	C	5	Total O 5 5	0	0
2	D	3	Total O 3 3	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.92Å 82.32Å 392.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.44 – 3.19 29.44 – 3.19	Depositor EDS
% Data completeness (in resolution range)	83.6 (29.44-3.19) 83.4 (29.44-3.19)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.74 (at 3.18Å)	Xtrriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.265 , 0.296 0.265 , 0.297	Depositor DCC
R_{free} test set	1180 reflections (4.19%)	wwPDB-VP
Wilson B-factor (Å ²)	59.0	Xtrriage
Anisotropy	0.133	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	9644	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/2156	0.73	2/2927 (0.1%)
1	B	0.33	0/2148	0.74	0/2928
1	C	0.37	1/2139 (0.0%)	0.74	0/2913
1	D	0.33	0/1865	0.64	0/2541
1	E	0.30	0/1508	0.68	0/2064
All	All	0.34	1/9816 (0.0%)	0.71	2/13373 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	161	ILE	C-N	8.14	1.44	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	56	ALA	CA-C-N	5.38	131.83	121.54
1	A	56	ALA	C-N-CA	5.38	131.83	121.54

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	242	ARG	Sidechain
1	E	242	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2113	0	1993	3	0
1	B	2105	0	1902	5	0
1	C	2094	0	1890	8	0
1	D	1829	0	1633	9	0
1	E	1489	0	1171	7	0
2	A	4	0	0	0	0
2	B	2	0	0	0	0
2	C	5	0	0	0	0
2	D	3	0	0	0	0
All	All	9644	0	8589	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:PRO:HB3	1:A:243:LEU:HD11	1.80	0.62
1:C:149:MET:HG2	1:C:253:ILE:HB	1.82	0.61
1:D:149:MET:HG2	1:D:253:ILE:HB	1.82	0.61
1:E:250:THR:O	1:E:276:SER:HA	2.09	0.53
1:E:31:LYS:HA	1:E:35:ILE:O	2.11	0.51
1:D:239:TYR:HA	1:D:242:ARG:HD3	1.92	0.51
1:B:225:MET:O	1:B:228:HIS:HB2	2.11	0.50
1:D:152:GLY:HA2	1:D:265:ASN:HB3	1.93	0.50
1:A:32:THR:HG21	1:A:110:SER:HB3	1.94	0.49
1:A:149:MET:HE2	1:A:149:MET:HB3	1.69	0.47
1:D:123:VAL:HG13	1:D:147:THR:HB	1.97	0.47
1:C:239:TYR:HA	1:C:242:ARG:HD3	1.97	0.46
1:C:30:ILE:HD13	1:C:111:ILE:HA	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:30:ILE:HD13	1:E:114:GLY:HA3	1.99	0.45
1:B:197:MET:HG2	1:B:286:HIS:HB3	1.98	0.45
1:B:187:VAL:O	1:B:191:ILE:HB	2.18	0.44
1:D:61:LEU:HD11	1:D:218:ARG:HE	1.83	0.44
1:E:239:TYR:HA	1:E:242:ARG:HD3	2.00	0.44
1:D:288:LEU:HG	1:D:293:TRP:HE1	1.83	0.44
1:D:74:LYS:HB3	1:D:301:GLU:HG3	2.00	0.43
1:C:214:ALA:O	1:C:218:ARG:HB2	2.19	0.43
1:C:153:PRO:HG3	1:C:269:LEU:HG	2.00	0.43
1:D:30:ILE:HD12	1:D:114:GLY:HA3	2.00	0.42
1:B:149:MET:HE2	1:B:149:MET:HB3	1.75	0.41
1:E:53:VAL:HG12	1:E:81:TYR:HE1	1.85	0.41
1:C:52:LEU:HB3	1:C:63:TRP:CE2	2.56	0.41
1:E:105:THR:HG21	1:E:134:ILE:HG22	2.02	0.41
1:C:74:LYS:HB3	1:C:301:GLU:HG3	2.03	0.40
1:B:25:MET:HA	1:B:41:SER:O	2.22	0.40
1:C:109:ILE:HG13	1:C:139:PHE:CG	2.56	0.40
1:E:109:ILE:HD13	1:E:109:ILE:HA	1.93	0.40
1:D:130:LEU:HD22	1:D:154:TRP:HE3	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/311 (88%)	263 (96%)	11 (4%)	0	100	100
1	B	282/311 (91%)	271 (96%)	10 (4%)	1 (0%)	30	64
1	C	279/311 (90%)	268 (96%)	11 (4%)	0	100	100
1	D	244/311 (78%)	237 (97%)	7 (3%)	0	100	100
1	E	217/311 (70%)	206 (95%)	11 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1296/1555 (83%)	1245 (96%)	50 (4%)	1 (0%)	48 80

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	157	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	218/270 (81%)	218 (100%)	0	100 100
1	B	210/270 (78%)	210 (100%)	0	100 100
1	C	204/270 (76%)	204 (100%)	0	100 100
1	D	170/270 (63%)	169 (99%)	1 (1%)	84 92
1	E	104/270 (38%)	104 (100%)	0	100 100
All	All	906/1350 (67%)	905 (100%)	1 (0%)	92 98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	144	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	182	ASN
1	A	188	ASN
1	A	220	ASN
1	A	286	HIS
1	A	305	ASN
1	B	33	ASN
1	B	44	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	106	ASN
1	B	188	ASN
1	C	69	GLN
1	C	217	ASN
1	C	221	ASN
1	C	262	HIS
1	D	69	GLN
1	D	221	ASN
1	E	47	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	278/311 (89%)	0.08	6 (2%) 62 46	24, 44, 62, 73	0
1	B	284/311 (91%)	0.12	1 (0%) 89 81	25, 45, 74, 84	0
1	C	283/311 (90%)	0.35	9 (3%) 50 35	26, 50, 90, 101	0
1	D	250/311 (80%)	0.42	8 (3%) 50 35	34, 62, 97, 108	0
1	E	229/311 (73%)	0.98	29 (12%) 9 7	44, 77, 104, 109	0
All	All	1324/1555 (85%)	0.37	53 (4%) 43 29	24, 53, 96, 109	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	155	GLY	5.5
1	E	85	ASP	4.4
1	E	96	SER	4.3
1	E	155	GLY	4.0
1	C	308	GLU	3.8
1	E	264	LYS	3.6
1	A	156	ASP	3.5
1	E	192	GLN	3.4
1	E	305	ASN	3.3
1	C	159	PRO	3.2
1	E	292	ASP	3.2
1	D	152	GLY	3.0
1	D	292	ASP	3.0
1	E	293	TRP	3.0
1	C	156	ASP	3.0
1	C	160	THR	3.0
1	E	64	ASP	2.9
1	E	150	SER	2.9
1	E	156	ASP	2.9
1	C	34	GLY	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	94	PRO	2.8
1	E	295	SER	2.8
1	D	182	ASN	2.8
1	E	87	GLY	2.8
1	C	167	SER	2.8
1	C	268	PHE	2.7
1	E	190	LEU	2.7
1	E	38	CYS	2.7
1	C	306	ASP	2.6
1	D	91	ASN	2.6
1	A	157	SER	2.6
1	E	90	THR	2.6
1	E	92	TYR	2.5
1	E	86	VAL	2.5
1	C	96	SER	2.4
1	D	96	SER	2.4
1	D	103	ASP	2.4
1	E	194	ALA	2.4
1	D	186	VAL	2.4
1	E	63	TRP	2.3
1	E	141	ASP	2.3
1	D	185	SER	2.3
1	A	168	ILE	2.2
1	E	193	GLY	2.2
1	E	296	ILE	2.2
1	A	306	ASP	2.2
1	E	269	LEU	2.2
1	E	286	HIS	2.2
1	B	280	THR	2.1
1	A	172	HIS	2.1
1	E	91	ASN	2.1
1	E	284	THR	2.0
1	E	84	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.