



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 19, 2026 – 04:22 PM EDT

PDB ID : 14AA / pdb_000014aa
Title : PanDDA analysis group deposition of ground-state model of Enterovirus D68 3Dpol
Authors : Biswas, I.; Ruiz, F.X.; Saini, M.; Balcomb, B.H.; von Delft, F.; Arnold, E.
Deposited on : 2026-03-18
Resolution : 1.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1904 (1.36-1.32)
Clashscore	180529	2038 (1.36-1.32)
Ramachandran outliers	177936	2016 (1.36-1.32)
Sidechain outliers	177891	2016 (1.36-1.32)
RSRZ outliers	164620	1903 (1.36-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	467	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	IPA	A	501	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4038 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3Dpol RNA Dependent RNA Polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	457	3639	2336	604	678	21	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	ALA	VAL	conflict	UNP F1T146
A	458	GLY	-	expression tag	UNP F1T146
A	459	SER	-	expression tag	UNP F1T146
A	460	SER	-	expression tag	UNP F1T146
A	461	GLY	-	expression tag	UNP F1T146
A	462	HIS	-	expression tag	UNP F1T146
A	463	HIS	-	expression tag	UNP F1T146
A	464	HIS	-	expression tag	UNP F1T146
A	465	HIS	-	expression tag	UNP F1T146
A	466	HIS	-	expression tag	UNP F1T146
A	467	HIS	-	expression tag	UNP F1T146

- Molecule 2 is ISOPROPYL ALCOHOL (CCD ID: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	3	1		
2	A	1	Total	C	O	0	0
			4	3	1		
2	A	1	Total	C	O	0	0
			4	3	1		
2	A	1	Total	C	O	0	0
			4	3	1		
2	A	1	Total	C	O	0	0
			4	3	1		
2	A	1	Total	C	O	0	0
			4	3	1		
2	A	1	Total	C	O	0	0
			4	3	1		
2	A	1	Total	C	O	0	0
			4	3	1		
2	A	1	Total	C	O	0	0
			4	3	1		
2	A	1	Total	C	O	0	0
			4	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	3	1		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	326	Total	O	0	0
			326	326		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3Dpol RNA Dependent RNA Polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.23Å 82.83Å 58.57Å 90.00° 108.62° 90.00°	Depositor
Resolution (Å)	55.56 – 1.34 55.50 – 1.34	Depositor EDS
% Data completeness (in resolution range)	73.0 (55.56-1.34) 73.0 (55.50-1.34)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 1.34Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.178 , 0.200 0.187 , 0.206	Depositor DCC
R_{free} test set	5730 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	17.9	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4038	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG, IPA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.16	6/3731 (0.2%)	1.23	10/5052 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	60	GLY	C-O	6.70	1.30	1.23
1	A	286	SER	C-O	6.03	1.31	1.23
1	A	3	ILE	C-O	5.90	1.29	1.24
1	A	284	SER	CA-CB	-5.71	1.43	1.53
1	A	287	SER	C-O	5.30	1.30	1.24
1	A	283	PRO	C-O	-5.21	1.17	1.24

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	289	THR	CA-CB-OG1	-7.57	98.24	109.60
1	A	130	HIS	CB-CA-C	6.19	120.59	110.88
1	A	409	HIS	N-CA-C	-6.01	104.64	111.07
1	A	284	SER	CA-CB-OG	-5.97	99.15	111.10
1	A	58	TYR	CA-C-O	-5.60	115.93	122.37
1	A	67	ASP	CA-CB-CG	5.32	117.92	112.60
1	A	129	ARG	CA-C-N	5.18	127.18	120.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	129	ARG	C-N-CA	5.18	127.18	120.44
1	A	284	SER	CB-CA-C	-5.16	100.15	110.42
1	A	305	LEU	CA-C-O	-5.03	115.54	120.82

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	286	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3639	0	3598	18	0
2	A	60	0	120	9	0
3	A	7	0	10	2	0
4	A	6	0	8	0	0
5	A	326	0	0	7	0
All	All	4038	0	3736	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:290:SER:H	2:A:501:IPA:H11	1.42	0.84
1:A:290:SER:H	2:A:501:IPA:C1	2.07	0.66
1:A:168:LYS:HG2	2:A:503:IPA:H33	1.79	0.65
3:A:510:PEG:H31	2:A:512:IPA:H32	1.81	0.61
1:A:142:GLU:OE1	5:A:601:HOH:O	2.16	0.60
1:A:284:SER:OG	1:A:285:GLY:N	2.29	0.60
1:A:362:THR:O	5:A:602:HOH:O	2.17	0.59
1:A:247:LYS:HG2	1:A:251:LYS:HE3	1.88	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:452:LYS:NZ	5:A:609:HOH:O	2.42	0.53
2:A:501:IPA:H13	5:A:823:HOH:O	2.08	0.53
1:A:268:VAL:CG1	1:A:275:LYS:HE3	2.39	0.52
1:A:457:PHE:C	5:A:607:HOH:O	2.53	0.52
1:A:117:LEU:HD12	3:A:510:PEG:H22	1.97	0.47
1:A:168:LYS:CG	2:A:503:IPA:H33	2.45	0.45
2:A:515:IPA:H11	5:A:915:HOH:O	2.17	0.44
1:A:168:LYS:CB	2:A:503:IPA:H33	2.49	0.42
2:A:501:IPA:C1	5:A:823:HOH:O	2.67	0.41
1:A:342:ALA:O	1:A:345:HIS:HB3	2.20	0.41
1:A:69:TYR:CD1	1:A:306:ILE:HG22	2.56	0.41
1:A:432:ILE:O	1:A:438:GLY:HA3	2.21	0.41
1:A:233:TYR:CG	1:A:324:ASP:HB3	2.57	0.40
1:A:268:VAL:HG11	1:A:275:LYS:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	455/467 (97%)	447 (98%)	7 (2%)	1 (0%)	44 19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	284	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	399/408 (98%)	399 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	IPA	A	502	-	3,3,3	0.15	0	3,3,3	0.58	0
2	IPA	A	515	-	3,3,3	0.17	0	3,3,3	0.76	0
2	IPA	A	507	-	3,3,3	0.44	0	3,3,3	0.60	0
3	PEG	A	510	-	6,6,6	0.19	0	5,5,5	0.10	0
2	IPA	A	513	-	3,3,3	0.34	0	3,3,3	0.30	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IPA	A	517	-	3,3,3	0.16	0	3,3,3	0.39	0
2	IPA	A	508	-	3,3,3	0.28	0	3,3,3	0.30	0
2	IPA	A	501	-	3,3,3	0.74	0	3,3,3	0.22	0
2	IPA	A	504	-	3,3,3	0.10	0	3,3,3	0.17	0
2	IPA	A	505	-	3,3,3	0.23	0	3,3,3	0.06	0
2	IPA	A	503	-	3,3,3	1.03	0	3,3,3	0.64	0
2	IPA	A	506	-	3,3,3	0.61	0	3,3,3	0.44	0
4	GOL	A	511	-	5,5,5	0.09	0	5,5,5	0.29	0
2	IPA	A	514	-	3,3,3	0.50	0	3,3,3	0.89	0
2	IPA	A	516	-	3,3,3	0.61	0	3,3,3	0.20	0
2	IPA	A	512	-	3,3,3	0.20	0	3,3,3	0.26	0
2	IPA	A	509	-	3,3,3	0.59	0	3,3,3	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	511	-	-	4/4/4/4	-
3	PEG	A	510	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	511	GOL	O1-C1-C2-C3
4	A	511	GOL	C1-C2-C3-O3
4	A	511	GOL	O1-C1-C2-O2
4	A	511	GOL	O2-C2-C3-O3
3	A	510	PEG	C1-C2-O2-C3
3	A	510	PEG	O2-C3-C4-O4

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	515	IPA	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	510	PEG	2	0
2	A	501	IPA	4	0
2	A	503	IPA	3	0
2	A	512	IPA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	457/467 (97%)	0.68	42 (9%) 16 20	11, 19, 38, 53	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	285	GLY	9.1
1	A	17	ALA	4.5
1	A	382	PHE	4.5
1	A	415	TYR	4.2
1	A	357	THR	3.9
1	A	310	LYS	3.8
1	A	311	GLY	3.7
1	A	130	HIS	3.5
1	A	165	GLU	3.3
1	A	358	SER	3.3
1	A	286	SER	3.2
1	A	457	PHE	3.1
1	A	12	CYS	3.0
1	A	15	ALA	3.0
1	A	59	THR	3.0
1	A	16	PRO	3.0
1	A	359	PHE	2.9
1	A	380	TYR	2.9
1	A	284	SER	2.8
1	A	13	ILE	2.8
1	A	212	ILE	2.8
1	A	341	GLU	2.7
1	A	28	GLN	2.6
1	A	405	ASN	2.6
1	A	161	LYS	2.6
1	A	172	ILE	2.6
1	A	18	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	355	LYS	2.5
1	A	121	LYS	2.5
1	A	44	PRO	2.5
1	A	446	TYR	2.5
1	A	349	VAL	2.4
1	A	306	ILE	2.4
1	A	338	LEU	2.4
1	A	312	ILE	2.3
1	A	287	SER	2.3
1	A	451	ARG	2.3
1	A	14	ASN	2.2
1	A	289	THR	2.1
1	A	196	SER	2.1
1	A	47	LYS	2.0
1	A	447	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	IPA	A	512	4/4	0.63	0.23	59,60,62,67	0
2	IPA	A	505	4/4	0.73	0.24	53,53,55,57	0
2	IPA	A	509	4/4	0.76	0.19	24,33,36,36	0
2	IPA	A	513	4/4	0.77	0.21	46,48,49,52	0
2	IPA	A	517	4/4	0.77	0.23	40,46,47,47	0
3	PEG	A	510	7/7	0.77	0.19	53,64,72,74	0
4	GOL	A	511	6/6	0.78	0.21	62,64,64,66	0
2	IPA	A	504	4/4	0.80	0.23	42,42,44,44	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	IPA	A	508	4/4	0.83	0.23	32,35,36,46	0
2	IPA	A	502	4/4	0.84	0.22	28,33,37,39	0
2	IPA	A	507	4/4	0.84	0.18	32,33,37,43	0
2	IPA	A	515	4/4	0.84	0.20	30,30,36,38	0
2	IPA	A	516	4/4	0.86	0.15	30,30,31,35	0
2	IPA	A	506	4/4	0.86	0.15	19,25,27,38	0
2	IPA	A	503	4/4	0.87	0.15	17,23,26,30	0
2	IPA	A	501	4/4	0.87	0.17	17,18,27,38	0
2	IPA	A	514	4/4	0.94	0.16	17,24,26,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.