



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 27, 2026 – 10:30 PM EDT

PDB ID : 14UM / pdb_000014um
Title : Crystal Structure of FtsZ with F2146-0687
Authors : Meeks, K.R.; Kreitler, D.F.; Lazo, E.O.
Deposited on : 2026-07-15
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	379	 3% 67% 11% • 21%
1	B	379	 4% 61% 16% • 22%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell division protein FtsZ.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	0	18	0
			2258	1400	393	454	11			
1	B	296	Total	C	N	O	S	0	58	0
			2543	1577	438	516	12			

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is N-(thiophen-2-yl)acetamide (CCD ID: OK7) (formula: C₆H₇NOS) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	1
			9	6	1	1	1		
3	B	1	Total	C	N	O	S	0	1
			9	6	1	1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	159	Total	O	0	16
			170	170		
4	B	149	Total	O	0	40
			176	176		

4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	88.97Å 88.97Å 180.14Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.52 – 1.95 44.52 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.9 (44.52-1.95) 99.9 (44.52-1.95)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.181 , 0.214 0.188 , 0.220	Depositor DCC
R_{free} test set	2958 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.053 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5170	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OK7, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2272	1.09	8/3067 (0.3%)
1	B	0.54	0/2560	1.02	4/3462 (0.1%)
All	All	0.59	0/4832	1.06	12/6529 (0.2%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	251	MET	CG-SD-CE	-7.14	85.19	100.90
1	A	184	ASP	CA-CB-CG	6.88	119.48	112.60
1	A	271	GLU	N-CA-CB	6.33	120.09	110.28
1	A	178	ASP	CA-CB-CG	6.26	118.86	112.60
1	B	49[A]	MET	CG-SD-CE	6.17	114.47	100.90
1	B	49[B]	MET	CG-SD-CE	6.17	114.47	100.90
1	B	280	GLN	N-CA-CB	5.82	118.78	110.16
1	A	135	PHE	CA-CB-CG	5.28	119.08	113.80
1	A	91	ARG	CB-CA-C	-5.22	102.02	109.84
1	A	131	ARG	CB-CA-C	-5.18	101.57	108.87
1	A	162	PRO	CB-CA-C	5.14	116.19	111.87
1	A	259	MET	CG-SD-CE	-5.08	89.73	100.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	0	2285	25	1
1	B	2543	0	2550	55	0
2	A	5	0	0	0	0
3	B	18	0	0	1	0
4	A	170	0	0	4	0
4	B	176	0	0	11	0
All	All	5170	0	4835	76	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22[B]:ASN:OD1	4:B:542[B]:HOH:O	1.81	0.99
1:A:50:SER:O	1:A:55[B]:LYS:NZ	2.08	0.86
1:A:60[B]:ARG:NH2	1:B:43[B]:ASP:O	2.17	0.77
1:B:214:ILE:HD12	1:B:290:ILE:HG12	1.67	0.76
1:A:301:ASP:OD2	4:A:503[B]:HOH:O	2.06	0.74
1:B:193:GLY:HA3	1:B:306[B]:THR:HG21	1.70	0.73
1:A:259:MET:SD	1:A:279:VAL:HG21	2.29	0.72
1:B:42[B]:THR:OG1	1:B:106:THR:OG1	2.03	0.72
1:A:139:ARG:HB3	4:A:642:HOH:O	1.92	0.70
1:B:8:LEU:HB3	4:B:611:HOH:O	1.91	0.70
1:B:103:GLY:O	4:B:501:HOH:O	2.11	0.67
1:B:259:MET:HE3	1:B:305[A]:VAL:HG11	1.76	0.66
1:A:82:ALA:HB2	1:B:45[B]:GLN:HE22	1.61	0.65
1:B:17:GLY:O	1:B:21[B]:VAL:HG23	1.99	0.63
1:B:28[B]:ILE:O	1:B:31:GLY:N	2.23	0.63
1:B:26[B]:ARG:NH1	4:B:506:HOH:O	2.33	0.61
1:A:283:ALA:CB	1:A:289:ILE:HD11	2.31	0.61
1:A:278:LEU:HD12	4:A:502[B]:HOH:O	2.02	0.59
1:B:41[A]:ASN:CG	4:B:505[A]:HOH:O	2.44	0.59
1:B:108:THR:HG22	4:B:649[A]:HOH:O	2.02	0.58
1:B:43[B]:ASP:OD2	4:B:502[B]:HOH:O	2.17	0.58
1:B:42[A]:THR:HG22	1:B:58[A]:VAL:CG2	2.34	0.58
1:A:60[B]:ARG:NH1	1:B:57[B]:ASP:OD2	2.36	0.57
1:A:131:ARG:NH1	1:A:145:GLU:OE1	2.36	0.57
1:B:26[A]:ARG:NH1	1:B:184:ASP:HB3	2.22	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:54[B]:VAL:HG11	1:B:89[B]:LEU:HD22	1.89	0.54
1:A:60[B]:ARG:NE	1:B:44[B]:ALA:O	2.35	0.54
1:B:47[B]:LEU:HD11	4:B:578[B]:HOH:O	2.07	0.53
1:A:299:LEU:HD21	1:A:304:ARG:NH1	2.24	0.52
1:B:250:SER:HB2	1:B:312:PHE:CE1	2.45	0.51
1:B:88[B]:GLU:OE2	1:B:91:ARG:NH2	2.40	0.51
1:B:24[B]:VAL:O	1:B:28[B]:ILE:HG13	2.11	0.50
1:B:56[B]:LEU:HG	1:B:58[B]:VAL:CG1	2.42	0.50
1:B:23[B]:ALA:HA	1:B:26[B]:ARG:CZ	2.40	0.50
1:A:242:ILE:HD11	1:A:279:VAL:HG22	1.92	0.50
1:B:41[A]:ASN:OD1	1:B:47[A]:LEU:HA	2.12	0.50
1:B:233:ARG:HD2	1:B:303[B]:VAL:HG13	1.92	0.50
1:B:57[A]:ASP:OD2	4:B:503[A]:HOH:O	2.20	0.49
1:B:228[B]:ALA:CB	1:B:236[B]:LYS:HD3	2.43	0.49
1:B:165:ARG:HE	1:B:168:GLN:HE22	1.59	0.49
1:A:242:ILE:CD1	1:A:279:VAL:HG22	2.42	0.49
1:B:45[A]:GLN:HG3	4:B:531:HOH:O	2.12	0.49
1:B:38[B]:ILE:HA	1:B:54[B]:VAL:O	2.12	0.47
1:A:259:MET:HE2	1:A:291:PHE:HD1	1.80	0.47
1:B:42[A]:THR:HG22	1:B:58[A]:VAL:HG21	1.96	0.47
1:B:90[B]:LEU:O	1:B:91:ARG:C	2.57	0.47
1:B:41[A]:ASN:OD1	1:B:47[A]:LEU:HD12	2.15	0.47
1:B:86:ILE:HA	1:B:89[A]:LEU:HG	1.97	0.47
1:B:228[B]:ALA:HB1	1:B:236[B]:LYS:HD3	1.97	0.46
1:A:47[A]:LEU:HD12	1:A:57[A]:ASP:HB2	1.97	0.46
1:A:82:ALA:CB	1:B:45[B]:GLN:HE22	2.29	0.46
1:B:88[B]:GLU:CD	1:B:91:ARG:HE	2.24	0.45
1:A:99:THR:HA	1:A:128:VAL:O	2.16	0.45
1:A:200:THR:HG23	1:A:294:VAL:HG13	1.97	0.45
1:B:72:PRO:O	1:B:109:GLY:HA3	2.17	0.45
1:B:47[B]:LEU:HD23	1:B:49[B]:MET:HE2	1.99	0.45
1:A:59[A]:GLY:HA2	4:A:592:HOH:O	2.16	0.44
1:B:87:GLU:OE2	1:B:91:ARG:NH2	2.50	0.44
1:B:238[A]:ALA:HB1	1:B:259:MET:HE1	1.99	0.44
1:B:214:ILE:HG22	1:B:215:MET:HG3	1.99	0.44
1:A:223:MET:HE2	1:A:225:ILE:HG23	2.00	0.43
1:B:229[B]:ARG:HA	1:B:301[B]:ASP:O	2.19	0.43
1:B:47[B]:LEU:HD21	1:B:49[B]:MET:HG2	2.00	0.43
1:B:53[B]:ASP:C	1:B:54[B]:VAL:CG2	2.92	0.42
1:A:41[A]:ASN:HB2	1:A:47[A]:LEU:HD21	2.02	0.42
1:A:272:ILE:HD13	1:A:293:THR:HG21	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:41[B]:ASN:ND2	1:B:46[B]:ALA:O	2.45	0.42
1:B:56[A]:LEU:HD23	1:B:58[A]:VAL:HG13	2.00	0.42
1:B:97:PHE:HA	1:B:126:VAL:O	2.20	0.42
1:A:95:MET:HE2	1:A:97:PHE:CZ	2.55	0.42
3:B:401[A]:OK7:SAA	3:B:401[A]:OK7:OAI	2.78	0.42
1:B:259:MET:HE3	1:B:305[A]:VAL:CG1	2.48	0.41
1:A:45[B]:GLN:O	1:A:49[B]:MET:HG2	2.20	0.41
1:B:47[B]:LEU:CD2	1:B:49[B]:MET:HE2	2.51	0.41
1:B:27[A]:MET:HG2	1:B:188:LEU:HG	2.03	0.40
1:B:212:LYS:HE3	4:B:571[B]:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:ASN:ND2	1:A:285:PRO:O[5_555]	1.93	0.27

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/379 (82%)	304 (98%)	7 (2%)	0	100	100
1	B	350/379 (92%)	340 (97%)	10 (3%)	0	100	100
All	All	661/758 (87%)	644 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/281 (82%)	226 (98%)	5 (2%)	45	35
1	B	261/281 (93%)	248 (95%)	13 (5%)	22	9
All	All	492/562 (88%)	474 (96%)	18 (4%)	36	16

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48[A]	LEU
1	A	48[B]	LEU
1	A	154	SER
1	A	220	THR
1	A	286	ASP
1	B	24[A]	VAL
1	B	24[B]	VAL
1	B	89[A]	LEU
1	B	89[B]	LEU
1	B	143	GLN
1	B	185[A]	GLU
1	B	185[B]	GLU
1	B	207	ASP
1	B	214	ILE
1	B	236[A]	LYS
1	B	236[B]	LYS
1	B	252	GLU
1	B	267	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	143	GLN
1	B	168	GLN
1	B	192	GLN
1	B	255	GLN
1	B	288	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	OK7	B	401[A]	-	8,9,9	0.37	0	8,11,11	0.50	0
2	PO4	A	401	-	4,4,4	1.75	1 (25%)	6,6,6	0.98	0
3	OK7	B	402[A]	-	8,9,9	0.19	0	8,11,11	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OK7	B	401[A]	-	-	0/4/4/4	0/1/1/1
3	OK7	B	402[A]	-	-	1/4/4/4	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	PO4	P-O4	-2.55	1.47	1.54

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	402[A]	OK7	SAA-CAE-NAF-CAG


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401[A]	OK7	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/379 (78%)	0.17	13 (4%) 39 43	14, 35, 66, 104	18 (6%)
1	B	296/379 (78%)	0.30	14 (4%) 36 41	14, 35, 66, 106	58 (19%)
All	All	594/758 (78%)	0.23	27 (4%) 38 43	14, 35, 66, 106	76 (12%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	142	ASN	5.8
1	B	7	TYR	5.7
1	B	8	LEU	5.7
1	B	141	SER	5.6
1	A	172	ALA	5.3
1	B	6	ASN	5.0
1	B	312	PHE	4.8
1	A	69	GLY	4.3
1	A	285	PRO	3.6
1	A	7	TYR	3.4
1	A	170	GLY	3.1
1	B	45[A]	GLN	3.0
1	B	172	ALA	2.9
1	B	140	ARG	2.9
1	B	139	ARG	2.9
1	A	173	ALA	2.8
1	A	312	PHE	2.8
1	A	171	ASP	2.5
1	B	214	ILE	2.4
1	B	29[A]	GLU	2.4
1	A	8	LEU	2.2
1	A	282	ALA	2.2
1	A	49[A]	MET	2.2
1	B	207	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	173	ALA	2.0
1	A	177	MET	2.0
1	A	251	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OK7	B	401[A]	9/9	0.89	0.17	43,44,54,55	9
3	OK7	B	402[A]	9/9	0.89	0.16	41,44,49,51	9
2	PO4	A	401	5/5	0.97	0.07	37,38,41,50	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OK7 B 401 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around OK7 B 402 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.