



# Full wwPDB NMR Structure Validation Report ⓘ

Mar 24, 2026 – 01:00 PM UTC

PDB ID : 1G5V / pdb\_00001g5v  
BMRB ID : 4899  
Title : SOLUTION STRUCTURE OF THE TUDOR DOMAIN OF THE HUMAN SMN PROTEIN  
Authors : Selenko, P.; Sprangers, R.; Stier, G.; Buehler, D.; Fischer, U.; Sattler, M.  
Deposited on : 2000-11-02

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 83%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	88	

## 2 Ensemble composition and analysis

This entry contains 10 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average, lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:92-A:144 (53)	0.46	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	2, 3, 9, 10
2	1, 4
3	6, 7
Single-model clusters	5; 8

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 864 atoms, of which 423 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called SURVIVAL MOTOR NEURON PROTEIN 1.


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	56	864	278	423	72	88	3	0

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1



#### 4.2.2 Score per residue for model 2 (medoid)

- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1



ASN  
ALA  
GLN  
GLU  
GLU  
ASN  
GLU  
SER  
GLN  
VAL  
SER  
THR  
ASP  
GLU

### 4.2.3 Score per residue for model 3

- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1

Chain A: 41% 19% 36%

LYS LYS ASN THR ALA ALA SER LEU Q90 Q91 Q92 K93 K97 K98 S99 E104 D105 G106 G107 A111 T112 T113 I116 Y125 Y126 Y127 Y128 R133 S139 D140 L141 L142 S143 P144 I145 CYS GLU VAL ALA ASN ASN ILE GLN GLN ALA GLN GLN GLN GLN VAL VAL ALA ASN ASN ILE GLN GLN GLN GLN GLN VAL SER THR ASP GLU VAL

SER  
THR  
ASP  
GLU

### 4.2.4 Score per residue for model 4

- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1

Chain A: 45% 15% 36%

LYS LYS ASN THR ALA ALA SER LEU Q90 Q91 Q92 K93 S103 E104 D105 G106 G107 I113 I116 K119 R120 Y127 T128 G129 Y130 R133 S143 P144 I145 CYS GLU VAL ALA ASN ASN ILE GLN GLN ALA GLN GLN GLN GLN VAL SER THR ASP GLU VAL

### 4.2.5 Score per residue for model 5

- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1

Chain A: 38% 23% 36%

LYS LYS ASN THR ALA ALA SER LEU Q90 Q91 Q92 K93 D96 K97 S99 E104 D105 G106 G107 S115 K119 R120 E121 Y127 Y128 R133 Q136 M137 L138 S139 S143 P144 I145 CYS GLU VAL ALA ASN ASN ILE GLN GLN GLN GLN VAL SER THR ASP GLU VAL

GLU  
SER  
GLN  
VAL  
SER  
THR  
ASP  
GLU

### 4.2.6 Score per residue for model 6


- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1

Chain A: 45% 14% 36%

LYS LYS ASN THR ALA ALA SER LEU Q90 Q91 D96 K97 G98 S99 E104 D105 G106 T112 D117 F118 K119 T128 R133 S139 D140 S143 P144 I145 CYS GLU VAL ALA ASN ASN ILE GLN GLN GLN GLN VAL SER THR ASP GLU VAL


### 4.2.7 Score per residue for model 7

- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1




### 4.2.8 Score per residue for model 8

- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1




### 4.2.9 Score per residue for model 9

- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1



### 4.2.10 Score per residue for model 10

- Molecule 1: SURVIVAL MOTOR NEURON PROTEIN 1



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *mixed torsion and Cartesian angle dynamics simulated annealing protocol*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *back calculated data agree with experimental NOESY spectrum, structures with acceptable covalent geometry, structures with favorable non-bond energy, structures with the least restraint violations, structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	0.3

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	622
Number of shifts mapped to atoms	615
Number of unparsed shifts	0
Number of shifts with mapping errors	7
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	83%

## 6 Model quality i

### 6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	415	396	396	2±1
All	All	4150	3960	3960	22

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:111:ALA:HB1	1:A:125:VAL:HB	0.58	1.75	3	3
1:A:96:ASP:HB3	1:A:113:ILE:HD12	0.56	1.75	8	3
1:A:127:TYR:HB3	1:A:130:TYR:HB2	0.56	1.77	4	2
1:A:102:TRP:CD1	1:A:127:TYR:HH	0.55	2.20	5	1
1:A:126:VAL:HG22	1:A:133:ARG:HG2	0.51	1.82	1	1
1:A:113:ILE:HG21	1:A:116:ILE:HD11	0.47	1.86	4	4
1:A:126:VAL:HG22	1:A:133:ARG:HG3	0.46	1.86	5	2
1:A:123:CYS:SG	1:A:138:LEU:HG	0.44	2.53	9	1
1:A:137:ASN:ND2	1:A:139:SER:HB3	0.43	2.28	1	1
1:A:137:ASN:HD21	1:A:139:SER:HB3	0.43	1.74	1	1
1:A:98:CYS:HB2	1:A:142:LEU:O	0.41	2.16	8	1
1:A:99:SER:HB2	1:A:142:LEU:HB2	0.41	1.92	8	1
1:A:111:ALA:HA	1:A:126:VAL:O	0.40	2.16	2	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	53/88 (60%)	45±2 (84±4%)	8±2 (14±4%)	1±1 (1±2%)	12	60
All	All	530/880 (60%)	447 (84%)	76 (14%)	7 (1%)	12	60

All 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	106	GLY	2
1	A	144	PRO	1
1	A	119	LYS	1
1	A	113	ILE	1
1	A	121	GLU	1
1	A	128	THR	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	46/77 (60%)	34±3 (74±7%)	12±3 (26±7%)	2	22
All	All	460/770 (60%)	339 (74%)	121 (26%)	2	22

All 27 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	128	THR	9
1	A	143	SER	8
1	A	119	LYS	7
1	A	120	ARG	7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	139	SER	7
1	A	93	LYS	6
1	A	97	LYS	6
1	A	99	SER	6
1	A	101	ILE	6
1	A	107	CYS	6
1	A	115	SER	5
1	A	103	SER	5
1	A	96	ASP	4
1	A	134	GLU	4
1	A	136	GLN	4
1	A	108	ILE	4
1	A	140	ASP	4
1	A	105	ASP	4
1	A	104	GLU	3
1	A	112	THR	3
1	A	142	LEU	3
1	A	121	GLU	2
1	A	133	ARG	2
1	A	137	ASN	2
1	A	117	ASP	2
1	A	122	THR	1
1	A	123	CYS	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 83% for the well-defined parts and 83% for the entire structure.

### 7.1 Chemical shift list 1

File name: working\_cs.cif

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	622
Number of shifts mapped to atoms	615
Number of unparsed shifts	0
Number of shifts with mapping errors	7
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. All 7 occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	146	CYS	H	8.528	0.02	1
1	A	146	CYS	HA	4.497	0.006	1
1	A	146	CYS	HB2	2.913	0.006	1
1	A	146	CYS	HB3	2.913	0.006	1
1	A	146	CYS	CA	58.255	0.02	1
1	A	146	CYS	CB	27.751	0.02	1
1	A	146	CYS	N	122.007	0.02	1

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	57	0.25 $\pm$ 0.17	None needed (< 0.5 ppm)

*Continued on next page...*

Continued from previous page...

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\beta$	53	0.35 $\pm$ 0.27	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
$^{15}\text{N}$	55	3.31 $\pm$ 0.53	Should be applied

### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 83%, i.e. 573 atoms were assigned a chemical shift out of a possible 690. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	212/265 (80%)	108/108 (100%)	53/106 (50%)	51/51 (100%)
Sidechain	319/364 (88%)	214/236 (91%)	103/116 (89%)	2/12 (17%)
Aromatic	42/61 (69%)	21/29 (72%)	19/30 (63%)	2/2 (100%)
Overall	573/690 (83%)	343/373 (92%)	175/252 (69%)	55/65 (85%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 83%, i.e. 615 atoms were assigned a chemical shift out of a possible 738. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	224/280 (80%)	114/114 (100%)	56/112 (50%)	54/54 (100%)
Sidechain	349/397 (88%)	234/257 (91%)	111/126 (88%)	4/14 (29%)
Aromatic	42/61 (69%)	21/29 (72%)	19/30 (63%)	2/2 (100%)
Overall	615/738 (83%)	369/400 (92%)	186/268 (69%)	60/70 (86%)

### 7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	104	GLU	HB3	-0.05	0.95 – 3.05	-9.8
1	A	104	GLU	HB2	0.39	1.00 – 3.05	-8.0

### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from

the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

