



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2025 – 04:36 AM EDT

PDB ID : 1LLB / pdb_000011lb
Title : Crystal Structure Of AmpC beta-Lactamase From E. Coli In Complex With ATMO-penicillin
Authors : Trehan, I.; Morandi, F.; Blaszcak, L.C.; Shoichet, B.K.
Deposited on : 2002-04-26
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7106 (1.74-1.70)
Clashscore	180529	7746 (1.74-1.70)
Ramachandran outliers	177936	7654 (1.74-1.70)
Sidechain outliers	177891	7654 (1.74-1.70)
RSRZ outliers	164620	7104 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	 4% 85% 14%
1	B	358	 % 87% 13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PCN	B	964[B]	-	-	X	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6242 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	358	Total 2796	C 1802	N 475	O 513	S 6	28	0	0
1	B	358	Total 2796	C 1802	N 475	O 513	S 6	8	0	0

- Molecule 2 is 2-{1-[2-(2-AMINO-THIAZOL-4-YL)-2-METHOXYIMINO-ACETYLAMINO]-2-OXO-ETHYL}-5,5-DIMETHYL-THIAZOLIDINE-4-CARBOXYLIC ACID (CCD ID: PCN) (formula: C₁₄H₁₉N₅O₅S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	B	1	Total 52	C 28	N 10	O 10	S 4	0	1


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	313	Total 313	O 313	0	0
3	B	285	Total 285	O 285	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: beta-lactamase



- Molecule 1: beta-lactamase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	118.66Å 76.73Å 98.17Å 90.00° 116.19° 90.00°	Depositor
Resolution (Å)	20.00 – 1.72 20.00 – 1.72	Depositor EDS
% Data completeness (in resolution range)	97.0 (20.00-1.72) 94.8 (20.00-1.72)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.26 (at 1.72Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.178 , 0.199 0.175 , 0.196	Depositor DCC
R_{free} test set	2411 reflections (3.03%)	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtrriage
Anisotropy	0.199	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6242	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PCN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/2876	1.01	11/3928 (0.3%)
1	B	0.61	0/2876	1.01	13/3928 (0.3%)
All	All	0.63	0/5752	1.01	24/7856 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	121	VAL	N-CA-C	-10.87	99.28	108.63
1	A	121	VAL	N-CA-C	-10.70	99.43	108.63
1	A	189	ILE	N-CA-C	-8.22	102.86	111.58
1	A	312	TRP	N-CA-C	-7.87	95.90	108.73
1	B	189	ILE	N-CA-C	-7.79	103.32	111.58
1	B	312	TRP	N-CA-C	-7.22	96.97	108.73
1	B	117	LEU	N-CA-C	-6.89	100.89	109.64
1	A	45	TYR	N-CA-C	6.87	120.65	109.59
1	A	319	THR	N-CA-C	-6.22	100.34	109.18
1	B	118	PRO	N-CA-C	6.17	121.18	111.38
1	B	44	GLY	N-CA-C	6.14	122.97	111.80
1	B	99	LYS	N-CA-C	6.00	119.44	111.75
1	B	54	VAL	N-CA-C	-5.99	101.07	109.21
1	B	222	GLY	N-CA-C	5.78	124.88	115.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	99	LYS	N-CA-C	5.74	118.31	111.71
1	A	169	SER	N-CA-C	-5.69	103.02	110.53
1	A	118	PRO	N-CA-C	5.53	120.16	111.38
1	B	151	ALA	N-CA-C	5.51	117.36	108.32
1	A	202	GLY	N-CA-C	-5.39	104.45	113.02
1	B	45	TYR	N-CA-C	5.38	118.17	109.40
1	B	202	GLY	N-CA-C	-5.20	104.75	113.02
1	A	44	GLY	N-CA-C	5.12	121.54	112.02
1	B	97	THR	N-CA-C	5.04	119.58	113.38
1	A	124	GLU	N-CA-C	5.01	118.89	112.87

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	45	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2796	0	2772	28	0
1	B	2796	0	2771	37	0
2	B	52	0	35	20	0
3	A	313	0	0	0	0
3	B	285	0	0	1	0
All	All	6242	0	5578	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:293:LEU:HD11	2:B:964[B]:PCN:C52	1.93	0.98

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:GLU:H	1:A:124:GLU:CD	1.86	0.83
1:B:293:LEU:HD11	2:B:964[B]:PCN:H522	1.61	0.78
1:B:211:VAL:HG23	2:B:964[B]:PCN:H191	1.49	0.77
1:A:287:SER:OG	1:A:350:VAL:HG21	1.85	0.76
1:B:293:LEU:HD11	2:B:964[B]:PCN:H521	1.69	0.73
1:B:289:ASN:ND2	2:B:964[B]:PCN:O63	2.22	0.72
1:A:17:THR:HB	1:A:18:PRO:HD3	1.73	0.71
1:B:280:PRO:HG3	1:B:354:TRP:CZ2	2.26	0.70
1:A:243:ILE:CD1	1:A:252:ILE:HD12	2.24	0.68
1:A:279:ASN:HD22	1:A:280:PRO:HD2	1.62	0.64
1:A:334:LEU:HG	1:A:357:LEU:HD22	1.79	0.64
1:B:243:ILE:CD1	1:B:252:ILE:HD12	2.30	0.62
1:B:293:LEU:HD21	2:B:964[A]:PCN:C52	2.30	0.61
1:A:164:LYS:HB2	1:A:165:PRO:HD3	1.83	0.60
1:A:124:GLU:CD	1:A:124:GLU:N	2.58	0.58
2:B:964[B]:PCN:O63	3:B:1160:HOH:O	2.17	0.58
1:B:211:VAL:HG23	2:B:964[B]:PCN:N19	2.16	0.57
2:B:964[A]:PCN:HC6	2:B:964[A]:PCN:O12	2.06	0.56
2:B:964[B]:PCN:C9	2:B:964[B]:PCN:S4	2.94	0.56
1:A:243:ILE:HD12	1:A:252:ILE:HD12	1.89	0.55
1:B:17:THR:HB	1:B:18:PRO:HD3	1.89	0.55
1:A:360:LEU:O	1:A:361:GLN:CB	2.55	0.54
1:A:169:SER:OG	1:A:172:GLN:HG3	2.07	0.54
1:A:232:ARG:HA	1:A:235:GLN:HE21	1.73	0.53
1:B:5:PRO:HD2	1:B:8:ILE:HD12	1.91	0.53
1:A:316:THR:HG22	1:A:325:TYR:CD1	2.43	0.52
1:A:275:ASP:O	1:A:278:VAL:HG12	2.10	0.51
1:B:334:LEU:HG	1:B:357:LEU:HD22	1.91	0.51
1:B:280:PRO:HG3	1:B:354:TRP:CE2	2.45	0.51
2:B:964[A]:PCN:H512	2:B:964[A]:PCN:C13	2.43	0.48
1:B:293:LEU:HD21	2:B:964[B]:PCN:H523	1.95	0.47
1:A:279:ASN:O	1:A:282:SER:HB3	2.15	0.47
1:B:221:TYR:CE1	2:B:964[B]:PCN:S16	3.08	0.47
1:A:265:MET:HG2	1:A:266:TYR:N	2.29	0.46
1:A:287:SER:HB3	1:A:346:ASN:HB3	1.97	0.46
1:A:344:TYR:CZ	1:A:349:ARG:HG2	2.50	0.46
1:B:343:ASN:C	1:B:343:ASN:HD22	2.24	0.46
1:A:276:TRP:CD2	1:A:277:PRO:HA	2.51	0.45
1:B:33:ILE:HD12	1:B:33:ILE:N	2.31	0.45
1:B:320:GLY:H	2:B:964[B]:PCN:H192	1.63	0.45
1:B:319:THR:HA	2:B:964[B]:PCN:N18	2.32	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:276:TRP:NE1	1:B:332:LYS:HE2	2.31	0.44
1:A:180:GLN:HB2	1:A:181:PRO:HD3	1.99	0.44
1:B:276:TRP:CD2	1:B:277:PRO:HA	2.52	0.44
1:A:344:TYR:CE2	1:A:349:ARG:HG2	2.53	0.44
1:B:171:GLU:CG	1:B:175:GLN:HE21	2.31	0.43
1:A:316:THR:HG22	1:A:325:TYR:HD1	1.81	0.43
1:B:320:GLY:H	2:B:964[A]:PCN:H192	1.66	0.43
2:B:964[B]:PCN:S4	2:B:964[B]:PCN:O91	2.77	0.43
1:B:164:LYS:HB2	1:B:165:PRO:HD3	2.01	0.43
1:B:293:LEU:HD21	2:B:964[A]:PCN:H521	2.01	0.42
1:B:243:ILE:HD12	1:B:252:ILE:HD12	2.01	0.42
1:B:343:ASN:C	1:B:343:ASN:ND2	2.75	0.42
1:B:83:ILE:HG22	1:B:92:TYR:CE2	2.55	0.42
1:B:26:PRO:HB3	1:B:48:ILE:HD11	2.00	0.42
1:A:280:PRO:O	1:A:284:ILE:HG13	2.20	0.42
1:A:107:LEU:HD22	1:B:303:PRO:HD3	2.01	0.41
1:B:171:GLU:HG2	1:B:175:GLN:NE2	2.35	0.41
1:B:293:LEU:CD1	2:B:964[B]:PCN:H522	2.42	0.41
1:A:34:TYR:CE2	1:A:35:GLN:HG3	2.56	0.41
1:B:262:THR:O	1:B:262:THR:HG23	2.21	0.41
1:A:275:ASP:O	1:A:278:VAL:CG1	2.68	0.41
1:B:293:LEU:HD11	2:B:964[A]:PCN:H521	2.02	0.41
1:B:91:LYS:HD3	1:B:92:TYR:CZ	2.55	0.41
1:B:273:MET:HG3	1:B:313:VAL:HG13	2.02	0.41
1:A:119:LEU:HA	1:A:151:ALA:HA	2.02	0.40
1:A:302:THR:HA	1:A:303:PRO:HA	2.00	0.40
1:B:302:THR:HA	1:B:303:PRO:HA	1.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	356/358 (99%)	349 (98%)	6 (2%)	1 (0%)	37	23
1	B	356/358 (99%)	350 (98%)	6 (2%)	0	100	100
All	All	712/716 (99%)	699 (98%)	12 (2%)	1 (0%)	48	32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	289	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/292 (100%)	288 (99%)	3 (1%)	73	63
1	B	291/292 (100%)	291 (100%)	0	100	100
All	All	582/584 (100%)	579 (100%)	3 (0%)	86	82

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	123	ASP
1	A	124	GLU
1	A	262	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	7	GLN
1	A	9	ASN
1	A	23	GLN
1	A	102	ASN
1	A	137	ASN
1	A	175	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	180	GLN
1	A	190	ASN
1	A	235	GLN
1	A	279	ASN
1	B	23	GLN
1	B	56	GLN
1	B	139	GLN
1	B	175	GLN
1	B	180	GLN
1	B	285	ASN
1	B	343	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PCN	B	964[B]	1	19,27,27	2.74	9 (47%)	23,39,39	2.04	6 (26%)
2	PCN	B	964[A]	1	19,27,27	2.83	7 (36%)	23,39,39	1.80	4 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCN	B	964[B]	1	-	7/15/40/40	0/2/2/2
2	PCN	B	964[A]	1	-	7/15/40/40	0/2/2/2

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	964[A]	PCN	C10-N11	7.68	1.40	1.29
2	B	964[B]	PCN	C10-N11	5.97	1.38	1.29
2	B	964[B]	PCN	C9-N8	4.99	1.44	1.34
2	B	964[A]	PCN	C9-N8	4.39	1.43	1.34
2	B	964[A]	PCN	O12-N11	3.90	1.49	1.40
2	B	964[B]	PCN	C15-S16	-3.88	1.64	1.70
2	B	964[B]	PCN	C14-N18	-3.69	1.26	1.37
2	B	964[A]	PCN	C14-N18	-3.59	1.26	1.37
2	B	964[B]	PCN	C14-C10	-3.33	1.42	1.47
2	B	964[A]	PCN	C15-S16	-3.15	1.65	1.70
2	B	964[B]	PCN	O1-C1	3.00	1.31	1.20
2	B	964[A]	PCN	O1-C1	2.90	1.31	1.20
2	B	964[A]	PCN	C10-C9	-2.81	1.45	1.50
2	B	964[B]	PCN	O12-N11	2.73	1.47	1.40
2	B	964[B]	PCN	C10-C9	-2.48	1.46	1.50
2	B	964[B]	PCN	C3-S4	2.31	1.89	1.84

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	964[A]	PCN	O91-C9-C10	5.48	126.68	120.45
2	B	964[B]	PCN	O12-N11-C10	5.44	117.49	111.50
2	B	964[B]	PCN	O91-C9-C10	5.06	126.20	120.45
2	B	964[A]	PCN	O12-N11-C10	4.01	115.92	111.50
2	B	964[B]	PCN	C14-C10-C9	3.43	124.50	118.17
2	B	964[B]	PCN	O91-C9-N8	-2.58	118.54	123.09
2	B	964[B]	PCN	C6-C5-S4	-2.52	99.07	103.86
2	B	964[A]	PCN	C6-C5-S4	-2.23	99.62	103.86
2	B	964[B]	PCN	C3-C2-N8	2.17	114.52	109.93
2	B	964[A]	PCN	O91-C9-N8	-2.08	119.43	123.09

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	964[B]	PCN	C1-C2-N8-C9
2	B	964[B]	PCN	N7-C6-C61-O62
2	B	964[B]	PCN	N7-C6-C61-O63
2	B	964[A]	PCN	N7-C6-C61-O62
2	B	964[A]	PCN	N7-C6-C61-O63
2	B	964[A]	PCN	N11-C10-C9-N8
2	B	964[B]	PCN	C10-N11-O12-C13
2	B	964[A]	PCN	N11-C10-C9-O91
2	B	964[A]	PCN	C14-C10-C9-O91
2	B	964[B]	PCN	N11-C10-C9-O91
2	B	964[B]	PCN	N11-C10-C9-N8
2	B	964[A]	PCN	C14-C10-C9-N8
2	B	964[A]	PCN	C1-C2-N8-C9
2	B	964[B]	PCN	C3-C2-N8-C9


There are no ring outliers.

2 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	964[B]	PCN	14	0
2	B	964[A]	PCN	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	357/358 (99%)	-0.06	15 (4%) 41 45	12, 21, 39, 62	5 (1%)
1	B	358/358 (100%)	-0.12	5 (1%) 73 77	14, 22, 35, 45	4 (1%)
All	All	715/716 (99%)	-0.09	20 (2%) 55 59	12, 22, 37, 62	9 (1%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	287	SER	6.5
1	A	284	ILE	4.9
1	A	291	ILE	4.5
1	A	292	ALA	4.3
1	A	286	GLY	4.2
1	A	293	LEU	4.1
1	A	283	ILE	3.4
1	A	354	TRP	2.9
1	A	294	ALA	2.8
1	A	285	ASN	2.8
1	A	278	VAL	2.7
1	B	244	ASN	2.7
1	B	56	GLN	2.6
1	A	4	ALA	2.5
1	A	288	ASP	2.5
1	B	64	SER	2.3
1	B	354	TRP	2.2
1	A	289	ASN	2.1
1	A	361	GLN	2.0
1	B	172	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCN	B	964[A]	26/26	0.61	0.18	29,35,37,43	26
2	PCN	B	964[B]	26/26	0.61	0.18	21,23,24,25	26




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PCN B 964 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around PCN B 964 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.