



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 3, 2023 – 02:29 pm GMT

PDB ID : 1WA6  
Title : The structure of ACC oxidase  
Authors : Zhang, Z.; Ren, J.-S.; Clifton, I.J.; Schofield, C.J.  
Deposited on : 2004-10-25  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	319	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	X	1308	-	X	-	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-AMINOCYCLOPROPANE-1-CARBOXYLATE OXIDASE 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	X	298	2372	1523	394	438	4	13	0	0	0

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	X	1	Total	Fe	0	0
			1	1		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	X	1	Total	O	P	0	0
			5	4	1		
3	X	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	X	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	X	72	Total	O	0	0
			72	72		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1-AMINOCYCLOPROPANE-1-CARBOXYLATE OXIDASE 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.34Å 107.06Å 108.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.55 39.84 – 2.55	Depositor EDS
% Data completeness (in resolution range)	93.4 (40.00-2.55) 93.4 (39.84-2.55)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.82 (at 2.54Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.186 , 0.236 0.191 , 0.241	Depositor DCC
$R_{free}$ test set	654 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.7	Xtrriage
Anisotropy	0.519	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 42.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.010 for -h,-l,-k	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2460	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FE2, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	X	1.47	19/2411 (0.8%)	1.30	23/3228 (0.7%)

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	87	GLU	CD-OE1	-8.80	1.16	1.25
1	X	43	ARG	CG-CD	8.63	1.73	1.51
1	X	286	MSE	SE-CE	-7.85	1.49	1.95
1	X	167	LYS	CD-CE	7.39	1.69	1.51
1	X	139	GLU	CD-OE1	6.56	1.32	1.25
1	X	59	LYS	CD-CE	6.55	1.67	1.51
1	X	51	LYS	CD-CE	6.36	1.67	1.51
1	X	103	VAL	CA-CB	6.17	1.67	1.54
1	X	148	TYR	CD1-CE1	5.75	1.48	1.39
1	X	29	GLU	CD-OE1	5.71	1.31	1.25
1	X	289	TYR	CB-CG	-5.63	1.43	1.51
1	X	191	LYS	CA-C	5.59	1.67	1.52
1	X	25	LYS	CE-NZ	5.50	1.62	1.49
1	X	159	VAL	CB-CG2	5.37	1.64	1.52
1	X	242	GLY	C-O	-5.30	1.15	1.23
1	X	21	MSE	SE-CE	-5.20	1.64	1.95
1	X	231	SER	CB-OG	-5.19	1.35	1.42
1	X	52	MSE	SE-CE	-5.18	1.64	1.95
1	X	254	GLY	C-O	5.17	1.31	1.23

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	189	ASP	CB-CG-OD2	9.77	127.09	118.30
1	X	26	ASP	CB-CG-OD2	9.71	127.03	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	10	ASP	CB-CG-OD2	8.91	126.32	118.30
1	X	116	ASP	CB-CG-OD2	8.89	126.30	118.30
1	X	47	ASP	CB-CG-OD2	8.58	126.03	118.30
1	X	50	GLU	OE1-CD-OE2	-8.30	113.34	123.30
1	X	43	ARG	NE-CZ-NH1	8.28	124.44	120.30
1	X	244	ARG	NE-CZ-NH1	7.78	124.19	120.30
1	X	235	ARG	NE-CZ-NH1	-7.48	116.56	120.30
1	X	130	ASP	CB-CG-OD2	7.27	124.84	118.30
1	X	256	ASP	CB-CG-OD2	7.21	124.79	118.30
1	X	210	ARG	NE-CZ-NH2	-7.18	116.71	120.30
1	X	300	ARG	NE-CZ-NH2	7.08	123.84	120.30
1	X	190	ASP	CB-CG-OD1	6.62	124.26	118.30
1	X	179	ASP	CB-CG-OD2	6.58	124.22	118.30
1	X	105	ASP	CB-CG-OD2	5.81	123.53	118.30
1	X	115	ARG	NE-CZ-NH2	-5.65	117.47	120.30
1	X	197	LEU	CB-CG-CD2	-5.33	101.94	111.00
1	X	244	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	X	200	ASP	CB-CG-OD2	5.21	122.99	118.30
1	X	235	ARG	NE-CZ-NH2	5.16	122.88	120.30
1	X	210	ARG	NE-CZ-NH1	5.10	122.85	120.30
1	X	219	ASP	CB-CG-OD2	5.08	122.87	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	2372	0	2372	67	0
2	X	1	0	0	0	0
3	X	10	0	0	2	0
4	X	5	0	0	0	0
5	X	72	0	0	3	2
All	All	2460	0	2372	67	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 14.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:307:MSE:SE	1:X:307:MSE:CE	2.14	1.46
1:X:190:ASP:CG	1:X:191:LYS:H	1.60	1.03
1:X:52:MSE:CE	1:X:110:TYR:HA	1.93	0.99
1:X:52:MSE:HE1	1:X:110:TYR:HA	1.53	0.89
1:X:68:LEU:HD22	1:X:84:MSE:HE3	1.59	0.82
1:X:93:LYS:H	1:X:99:ASN:ND2	1.77	0.82
1:X:197:LEU:N	1:X:197:LEU:HD23	1.97	0.79
1:X:197:LEU:HD23	1:X:197:LEU:H	1.51	0.76
1:X:91:PHE:O	1:X:92:LEU:HD23	1.87	0.75
1:X:93:LYS:H	1:X:99:ASN:HD21	1.32	0.74
1:X:190:ASP:CG	1:X:191:LYS:N	2.40	0.73
1:X:180:ALA:HB1	1:X:286:MSE:HE1	1.76	0.68
1:X:190:ASP:OD1	1:X:191:LYS:N	2.27	0.66
1:X:64:ARG:HG2	1:X:84:MSE:HE1	1.77	0.65
1:X:94:HIS:O	1:X:95:LEU:HD12	1.97	0.64
1:X:52:MSE:HE3	1:X:110:TYR:HA	1.79	0.64
1:X:46:MSE:HE2	1:X:187:PHE:CE1	2.33	0.63
1:X:90:PHE:HE1	1:X:92:LEU:HD21	1.65	0.61
1:X:192:VAL:HG21	1:X:241:ASP:OD2	2.01	0.59
1:X:150:SER:HB3	1:X:256:ASP:OD2	2.01	0.59
1:X:84:MSE:HE2	1:X:85:ASP:H	1.66	0.59
1:X:93:LYS:N	1:X:99:ASN:ND2	2.49	0.58
1:X:52:MSE:HE2	1:X:110:TYR:HD1	1.70	0.56
1:X:84:MSE:HA	1:X:84:MSE:CE	2.36	0.56
1:X:132:LEU:O	1:X:136:LEU:HB2	2.06	0.55
1:X:225:THR:HA	1:X:265:LEU:HD23	1.88	0.55
1:X:25:LYS:O	1:X:29:GLU:HG3	2.06	0.55
1:X:185:LEU:HD23	1:X:215:VAL:HG22	1.88	0.55
1:X:142:TYR:CD2	1:X:142:TYR:C	2.79	0.55
1:X:93:LYS:N	1:X:99:ASN:HD21	1.99	0.54
1:X:188:GLN:OE1	1:X:193:SER:HB3	2.07	0.54
1:X:184:ILE:HG21	3:X:1308:PO4:O1	2.08	0.54
1:X:90:PHE:CE1	1:X:92:LEU:HD21	2.42	0.54
1:X:286:MSE:HA	1:X:286:MSE:HE2	1.89	0.54
1:X:52:MSE:CE	1:X:110:TYR:HD1	2.21	0.53
1:X:180:ALA:CB	1:X:286:MSE:HE1	2.39	0.53
1:X:84:MSE:HE2	1:X:84:MSE:HA	1.90	0.53
1:X:115:ARG:HD2	5:X:2032:HOH:O	2.08	0.52

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:136:LEU:HD13	1:X:229:TYR:CZ	2.45	0.52
1:X:150:SER:CB	1:X:256:ASP:OD2	2.60	0.50
1:X:122:GLU:O	1:X:126:GLU:HG3	2.13	0.48
1:X:188:GLN:OE1	1:X:193:SER:CB	2.62	0.48
1:X:63:GLN:NE2	5:X:2018:HOH:O	2.47	0.48
1:X:241:ASP:OD1	1:X:241:ASP:C	2.51	0.48
1:X:46:MSE:HE2	1:X:187:PHE:CD1	2.49	0.47
1:X:195:LEU:HD23	1:X:206:VAL:HG21	1.97	0.47
1:X:111:ARG:O	1:X:115:ARG:HG3	2.15	0.46
1:X:71:SER:O	1:X:74:LEU:HB2	2.15	0.46
1:X:285:TYR:HD2	1:X:286:MSE:CE	2.28	0.46
1:X:242:GLY:O	1:X:243:ALA:HB3	2.17	0.45
1:X:52:MSE:HE3	1:X:110:TYR:CA	2.47	0.45
1:X:262:ALA:HB3	1:X:265:LEU:HD22	2.00	0.44
1:X:145:ASN:ND2	5:X:2045:HOH:O	2.25	0.43
1:X:23:MSE:HE3	3:X:1309:PO4:O2	2.18	0.43
1:X:192:VAL:O	1:X:192:VAL:HG12	2.19	0.42
1:X:171:ILE:O	1:X:172:LYS:C	2.57	0.42
1:X:99:ASN:ND2	1:X:99:ASN:C	2.73	0.42
1:X:194:GLY:HA3	1:X:237:ILE:O	2.20	0.42
1:X:21:MSE:HE1	1:X:131:LEU:HD13	2.02	0.41
1:X:50:GLU:O	1:X:54:LYS:HG3	2.20	0.41
1:X:189:ASP:O	1:X:190:ASP:C	2.59	0.41
1:X:181:GLY:HA3	1:X:252:ASN:OD1	2.21	0.41
1:X:136:LEU:HD13	1:X:229:TYR:OH	2.21	0.40
1:X:155:PHE:HA	1:X:250:PHE:O	2.22	0.40
1:X:12:VAL:HG12	1:X:21:MSE:HE2	2.04	0.40
1:X:45:VAL:O	1:X:49:VAL:HG23	2.21	0.40
1:X:56:HIS:CE1	1:X:60:CYS:SG	3.14	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:X:2025:HOH:O	5:X:2064:HOH:O[2_665]	0.52	1.68
5:X:2023:HOH:O	5:X:2050:HOH:O[2_665]	1.08	1.12

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	257/261 (98%)	242 (94%)	15 (6%)	20   26

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	9	LEU
1	X	36	LEU
1	X	43	ARG
1	X	44	GLU
1	X	59	LYS
1	X	74	LEU
1	X	84	MSE
1	X	97	ILE
1	X	99	ASN
1	X	104	PRO
1	X	136	LEU
1	X	167	LYS
1	X	169	ASP
1	X	210	ARG
1	X	265	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	56	HIS
1	X	63	GLN
1	X	78	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	X	99	ASN
1	X	196	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	X	1309	-	4,4,4	0.94	0	6,6,6	1.13	0
3	PO4	X	1308	2	4,4,4	0.49	0	6,6,6	2.47	4 (66%)
4	SO4	X	1310	-	4,4,4	0.51	0	6,6,6	1.62	1 (16%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	X	1308	PO4	O4-P-O2	3.87	120.38	107.97
4	X	1310	SO4	O4-S-O3	-3.47	94.26	109.06

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	X	1308	PO4	O2-P-O1	-2.91	100.23	110.89
3	X	1308	PO4	O3-P-O1	-2.22	102.77	110.89
3	X	1308	PO4	O3-P-O2	2.17	114.92	107.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	X	1309	PO4	1	0
3	X	1308	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	X	285/319 (89%)	-0.06	7 (2%) 57 63	32, 50, 73, 83	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	265	LEU	2.9
1	X	276	VAL	2.8
1	X	101	SER	2.7
1	X	100	ILE	2.7
1	X	110	TYR	2.4
1	X	97	ILE	2.3
1	X	275	GLN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PO4	X	1308	5/5	0.86	0.22	86,86,93,96	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	SO4	X	1310	5/5	0.88	0.14	78,81,87,88	0
3	PO4	X	1309	5/5	0.91	0.32	101,103,105,106	0
2	FE2	X	1307	1/1	0.99	0.14	58,58,58,58	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.