



# Full wwPDB X-ray Structure Validation Report i

Jan 28, 2026 – 06:08 PM JST

PDB ID : 21IP / pdb\_000021ip  
Title : The MIDN Catch-IRF4 (A8C) fusion protein  
Authors : Ding, J.; Chen, Z.; Zhong, Y.  
Deposited on : 2025-12-14  
Resolution : 3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.47


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1085 (3.32-3.28)
Clashscore	180529	1128 (3.32-3.28)
Ramachandran outliers	177936	1125 (3.32-3.28)
Sidechain outliers	177891	1124 (3.32-3.28)
RSRZ outliers	164620	1085 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	68	10%	72%	25%	.
2	B	73	4%	62%	23%	• 14%

## 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 978 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Midnolin, Interferon regulatory factor 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	66	Total	C	N	O	S	0	0	0
			496	312	79	98	7			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLN	deletion	UNP Q15306
A	?	-	GLY	deletion	UNP Q15306
A	173	CYS	ALA	engineered mutation	UNP Q15306

- Molecule 2 is a protein called Midnolin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	63	Total	C	N	O	S	0	0	0
			482	303	91	86	2			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	335	GLY	-	expression tag	UNP Q504T8
B	336	SER	-	expression tag	UNP Q504T8
B	337	GLY	-	expression tag	UNP Q504T8


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Midnolin, Interferon regulatory factor 4



- Molecule 2: Midnolin



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.40Å 88.40Å 72.69Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.20 – 3.30 44.20 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.3 (44.20-3.30) 99.4 (44.20-3.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.64 (at 3.32Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
$R$ , $R_{free}$	0.248 , 0.291 0.250 , 0.300	Depositor DCC
$R_{free}$ test set	288 reflections (10.32%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	92.4	Xtriage
Anisotropy	0.231	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 82.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	978	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.49	0/507	0.68	0/691
2	B	0.44	0/493	0.61	0/668
All	All	0.47	0/1000	0.65	0/1359

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	496	0	472	10	0
2	B	482	0	480	13	0
All	All	978	0	952	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:267:ALA:HB2	2:B:312:ASP:HB3	1.88	0.56
1:A:157:ALA:O	1:A:160:ALA:HB2	2.09	0.52

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:MET:HG3	1:A:175:ALA:HB2	1.94	0.49
1:A:177:PRO:HD2	2:B:277:PRO:O	2.12	0.49
2:B:266:GLY:O	2:B:286:GLY:HA2	2.13	0.48
1:A:133:LEU:HA	1:A:156:LEU:HD12	1.96	0.46
2:B:325:PRO:O	2:B:329:GLN:HG2	2.16	0.46
1:A:150:MET:HE1	2:B:327:LEU:HG	1.98	0.46
1:A:114:GLU:CD	2:B:330:LEU:HD11	2.41	0.46
1:A:114:GLU:HG3	2:B:330:LEU:HD21	1.98	0.45
1:A:132:PHE:CD1	1:A:139:LEU:HB2	2.53	0.43
1:A:125:THR:O	1:A:129:VAL:HG23	2.20	0.42
2:B:320:TYR:O	2:B:323:MET:HB2	2.19	0.42
2:B:272:PHE:HE1	2:B:280:PHE:HB2	1.85	0.42
1:A:121:LEU:O	2:B:303:ILE:HD13	2.20	0.41
2:B:318:ARG:HB2	2:B:328:ALA:HA	2.03	0.41
2:B:317:THR:O	2:B:323:MET:HG3	2.21	0.40
2:B:307:LEU:HA	2:B:307:LEU:HD23	1.87	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	64/68 (94%)	63 (98%)	1 (2%)	0	100 100
2	B	59/73 (81%)	56 (95%)	3 (5%)	0	100 100
All	All	123/141 (87%)	119 (97%)	4 (3%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	56/59 (95%)	52 (93%)	4 (7%)	12 37
2	B	53/59 (90%)	52 (98%)	1 (2%)	52 72
All	All	109/118 (92%)	104 (95%)	5 (5%)	23 52

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	SER
1	A	147	ASP
1	A	149	MET
1	A	170	THR
2	B	327	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	274	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	66/68 (97%)	0.66	7 (10%) 13 12	56, 78, 124, 141	0
2	B	63/73 (86%)	0.65	3 (4%) 36 28	61, 98, 145, 160	0
All	All	129/141 (91%)	0.66	10 (7%) 20 18	56, 91, 140, 160	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	149	MET	3.6
1	A	162	GLU	3.1
2	B	292	CYS	2.8
1	A	174	CYS	2.7
1	A	113	PRO	2.6
1	A	122	GLU	2.5
2	B	300	ARG	2.4
1	A	153	GLN	2.2
1	A	118	MET	2.2
2	B	308	GLN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates i

There are no oligosaccharides in this entry.

### 6.4 Ligands i

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.