



Full wwPDB EM Validation Report ⓘ

Oct 18, 2026 – 05:25 PM JST

PDB ID : 21RZ / pdb_000021rz
EMDB ID : EMD-67956
Title : cryoEM structure of PROTEIN TANC2
Authors : Xue, C.Y.; Liu, Z.M.; Liu, W.; Zhong, D.
Deposited on : 2025-12-23
Resolution : 3.45 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.2
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.51



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13836 (2.95 - 3.95)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1994	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6395 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein TANC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	815	6395	4064	1119	1179	33	0	0



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, POINT, POINT, POINT	Depositor
Number of particles used	59675, 59675, 59675, 59675	Depositor
Resolution determination method	FSC 0.143 CUT-OFF, FSC 0.143 CUT-OFF, FSC 0.143 CUT-OFF, FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION, PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI MORGAGNI	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.397	Depositor
Minimum map value	-1.005	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.030	Depositor
Recommended contour level	0.19	Depositor
Map size ($\text{\AA}$)	251.4, 251.4, 251.4	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.838, 0.838, 0.838	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.08	10/6517 (0.2%)	1.54	38/8817 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	473	HIS	C-O	134.44	3.13	1.23
1	A	473	HIS	CG-CD2	51.76	1.92	1.35
1	A	473	HIS	CE1-NE2	33.21	1.65	1.32
1	A	473	HIS	CD2-NE2	25.48	1.65	1.37
1	A	473	HIS	ND1-CE1	23.34	1.55	1.32
1	A	473	HIS	CG-ND1	17.07	1.57	1.38
1	A	473	HIS	CB-CG	-8.33	1.38	1.50
1	A	504	LEU	CG-CD1	-8.01	1.26	1.52
1	A	763	VAL	CB-CG2	-7.53	1.27	1.52
1	A	995	LEU	CG-CD2	-5.82	1.33	1.52

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	473	HIS	CA-CB-CG	-34.16	79.64	113.80
1	A	869	LEU	N-CA-CB	14.33	130.78	110.01
1	A	1181	VAL	CA-CB-CG1	13.46	133.29	110.40
1	A	473	HIS	CA-C-O	-13.10	106.12	121.32
1	A	712	VAL	CA-CB-CG1	12.99	132.48	110.40
1	A	911	VAL	CA-CB-CG1	12.62	131.86	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	473	HIS	CG-CD2-NE2	-11.94	95.26	107.20
1	A	577	VAL	CA-CB-CG1	11.43	129.83	110.40
1	A	473	HIS	O-C-N	10.63	136.18	121.83
1	A	867	ARG	N-CA-C	-9.54	97.05	110.50
1	A	836	LEU	CB-CG-CD1	7.87	134.31	110.70
1	A	763	VAL	CA-CB-CG1	7.78	123.62	110.40
1	A	804	LEU	CB-CG-CD1	7.58	133.45	110.70
1	A	609	ASP	CA-C-N	7.25	133.96	122.21
1	A	609	ASP	C-N-CA	7.25	133.96	122.21
1	A	771	LEU	CB-CG-CD1	6.96	131.59	110.70
1	A	497	HIS	CA-CB-CG	6.62	120.42	113.80
1	A	774	ARG	NE-CZ-NH2	6.27	124.84	119.20
1	A	724	PHE	CA-CB-CG	6.21	120.00	113.80
1	A	1038	LEU	CB-CG-CD1	6.17	129.21	110.70
1	A	839	LEU	CB-CG-CD1	5.95	128.54	110.70
1	A	1170	LEU	CB-CG-CD1	5.87	128.30	110.70
1	A	573	LEU	CB-CG-CD1	5.80	128.09	110.70
1	A	995	LEU	CB-CG-CD1	5.51	127.24	110.70
1	A	592	PHE	CA-CB-CG	5.45	119.25	113.80
1	A	473	HIS	CE1-NE2-CD2	5.38	114.38	109.00
1	A	447	ARG	NE-CZ-NH2	5.28	123.96	119.20
1	A	1163	ASP	CA-CB-CG	5.26	117.86	112.60
1	A	911	VAL	CB-CA-C	5.25	118.72	110.95
1	A	764	HIS	CA-C-N	5.21	126.35	119.84
1	A	764	HIS	C-N-CA	5.21	126.35	119.84
1	A	768	ARG	NE-CZ-NH2	5.20	123.88	119.20
1	A	723	ILE	CA-C-N	5.13	127.11	120.44
1	A	723	ILE	C-N-CA	5.13	127.11	120.44
1	A	680	PRO	N-CA-C	-5.10	101.97	112.47
1	A	656	TYR	CA-CB-CG	5.06	123.00	113.90
1	A	948	ASP	CA-CB-CG	5.04	117.64	112.60
1	A	763	VAL	CB-CA-C	5.01	118.05	111.33

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	473	HIS	Sidechain
1	A	580	ARG	Sidechain
1	A	656	TYR	Sidechain
1	A	756	ARG	Sidechain
1	A	867	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6395	0	6449	26	0
All	All	6395	0	6449	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:473:HIS:CD2	1:A:473:HIS:CG	1.92	1.53
1:A:473:HIS:CD2	1:A:473:HIS:C	2.18	1.19
1:A:473:HIS:CG	1:A:473:HIS:O	2.07	1.06
1:A:473:HIS:C	1:A:473:HIS:CE1	2.32	1.06
1:A:473:HIS:CD2	1:A:473:HIS:O	2.12	1.03
1:A:473:HIS:O	1:A:473:HIS:CE1	2.16	0.98
1:A:473:HIS:C	1:A:473:HIS:NE2	2.32	0.88
1:A:473:HIS:C	1:A:473:HIS:ND1	2.12	0.88
1:A:473:HIS:O	1:A:473:HIS:NE2	2.10	0.85
1:A:473:HIS:O	1:A:473:HIS:ND1	2.09	0.85
1:A:618:ILE:HG23	1:A:619:HIS:H	1.61	0.65
1:A:610:LEU:HD23	1:A:611:GLN:H	1.62	0.65
1:A:669:LEU:HD11	1:A:678:VAL:HG12	1.77	0.65
1:A:473:HIS:NE2	1:A:474:ASN:HA	2.19	0.57
1:A:1087:LEU:HA	1:A:1091:ALA:HB3	1.88	0.56
1:A:836:LEU:HA	1:A:839:LEU:HD22	1.88	0.56
1:A:753:ILE:HD11	1:A:763:VAL:HB	1.92	0.50
1:A:473:HIS:CE1	1:A:474:ASN:HA	2.46	0.50
1:A:470:GLU:O	1:A:471:PHE:C	2.54	0.49
1:A:678:VAL:O	1:A:679:VAL:C	2.55	0.49
1:A:473:HIS:NE2	1:A:474:ASN:CA	2.75	0.49
1:A:839:LEU:C	1:A:839:LEU:HD23	2.40	0.47
1:A:753:ILE:CD1	1:A:763:VAL:HB	2.47	0.45
1:A:683:LEU:HD22	1:A:687:TYR:CZ	2.55	0.41
1:A:468:VAL:HG22	1:A:558:ILE:HA	2.02	0.41
1:A:468:VAL:N	1:A:469:PRO:CD	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	807/1994 (40%)	699 (87%)	94 (12%)	14 (2%)	7	35

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	340	VAL
1	A	618	ILE
1	A	1182	VAL
1	A	323	ILE
1	A	344	ARG
1	A	868	LEU
1	A	446	ARG
1	A	476	ALA
1	A	765	PRO
1	A	1062	GLN
1	A	1093	VAL
1	A	1090	GLY
1	A	447	ARG
1	A	764	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	698/1730 (40%)	640 (92%)	58 (8%)	10	35

All (58) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	322	ASP
1	A	325	LEU
1	A	340	VAL
1	A	344	ARG
1	A	364	GLN
1	A	366	VAL
1	A	377	THR
1	A	394	ARG
1	A	439	ARG
1	A	470	GLU
1	A	473	HIS
1	A	475	VAL
1	A	491	GLN
1	A	497	HIS
1	A	504	LEU
1	A	539	ILE
1	A	565	MET
1	A	577	VAL
1	A	637	THR
1	A	646	LEU
1	A	654	TYR
1	A	655	LEU
1	A	656	TYR
1	A	666	LYS
1	A	694	LYS
1	A	712	VAL
1	A	719	THR
1	A	721	GLU
1	A	732	ILE
1	A	757	ASP
1	A	763	VAL
1	A	771	LEU
1	A	799	ARG
1	A	800	GLN
1	A	804	LEU
1	A	806	ARG
1	A	839	LEU
1	A	864	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	867	ARG
1	A	869	LEU
1	A	889	LEU
1	A	893	SER
1	A	911	VAL
1	A	933	ILE
1	A	955	LEU
1	A	966	VAL
1	A	995	LEU
1	A	996	ILE
1	A	1020	GLU
1	A	1044	ARG
1	A	1047	LEU
1	A	1062	GLN
1	A	1076	ARG
1	A	1102	THR
1	A	1126	ILE
1	A	1132	GLU
1	A	1170	LEU
1	A	1181	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	354	GLN
1	A	462	ASN
1	A	474	ASN
1	A	611	GLN
1	A	645	HIS
1	A	692	ASN
1	A	728	ASN
1	A	790	HIS
1	A	815	HIS
1	A	857	ASN
1	A	862	ASN
1	A	892	GLN
1	A	894	HIS
1	A	952	GLN
1	A	1027	ASN
1	A	1062	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-67956. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150

6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 154




Y Index: 156




Z Index: 154


6.3.2 Raw map



X Index: 155



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.19. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 47 nm³; this corresponds to an approximate mass of 42 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.290 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.290 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	-	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.70	7.77	6.15

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-67956 and PDB model 21RZ. Per-residue inclusion information can be found in [section 3](#) on [page 4](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.19 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.19).

9.4 Atom inclusion [i](#)



At the recommended contour level, 78% of all backbone atoms, 67% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.19) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6670	0.3640
A	0.6670	0.3640

