



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 02:19 AM JST

PDB ID : 22FD / pdb_000022fd
Title : Crystal structure of thioredoxin glutathione reductase from *Schistosoma japonicum* with the U597C mutation in complex with NADPH
Authors : Wang, S.Q.; Huang, S.Q.; Lin, T.W.
Deposited on : 2026-01-08
Resolution : 1.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	598	
1	B	598	

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9836 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called thioredoxin-disulfide reductase (NADPH).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	592	Total 4562	C 2883	N 780	O 874	S 25	0	4	0
1	B	592	Total 4553	C 2879	N 775	O 874	S 25	0	3	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	B	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 3 is 2'-MONOPHOSPHOADENOSINE-5'-DIPHOSPHATE (CCD ID: ATR) (formula: $C_{10}H_{16}N_5O_{13}P_3$).

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	290	Total 290	O 290	0	0
5	B	249	Total 249	O 249	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	84.22Å 86.36Å 183.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.37 – 1.90 60.37 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (60.37-1.90) 100.0 (60.37-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.190 , 0.227 0.199 , 0.235	Depositor DCC
R_{free} test set	5351 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtrriage
Anisotropy	0.378	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 24.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.013 for k,h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9836	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLA, FAD, ATR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/4650	1.02	6/6294 (0.1%)
1	B	0.58	0/4641	1.00	5/6284 (0.1%)
All	All	0.59	0/9291	1.01	11/12578 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
All	All	0	5

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	471	THR	CA-CB-OG1	-9.03	96.06	109.60
1	A	471	THR	OG1-CB-CG2	7.31	123.92	109.30
1	B	517	ASP	CA-CB-CG	7.00	119.60	112.60
1	A	517	ASP	CA-CB-CG	6.71	119.31	112.60
1	A	271	GLU	CB-CG-CD	6.27	123.25	112.60
1	B	304	PHE	CA-CB-CG	6.20	120.00	113.80
1	A	277	ASP	CA-CB-CG	5.55	118.15	112.60
1	A	304	PHE	CA-CB-CG	5.46	119.26	113.80
1	B	277	ASP	CA-CB-CG	5.23	117.83	112.60
1	B	37	VAL	N-CA-CB	5.13	118.28	110.58
1	B	147	THR	CA-CB-OG1	-5.12	101.92	109.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	13	ARG	Sidechain
1	A	78	ARG	Sidechain
1	B	566	ARG	Sidechain
1	B	596	CYS	Peptide
1	B	78	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4562	0	4551	20	0
1	B	4553	0	4536	28	0
2	A	53	0	31	0	0
2	B	53	0	31	1	0
3	A	31	0	11	4	0
3	B	31	0	11	2	0
4	A	7	0	2	1	0
4	B	7	0	2	1	0
5	A	290	0	0	1	0
5	B	249	0	0	1	0
All	All	9836	0	9175	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:352:ILE:HD11	1:B:389:PHE:HZ	1.51	0.76
1:A:317[B]:ARG:HH11	3:A:602:ATR:C6	2.03	0.71
1:A:175:LEU:HD23	1:B:175:LEU:HD23	1.76	0.66
1:B:173:HIS:HE1	4:B:603:MLA:O1A	1.78	0.66
1:A:317[B]:ARG:NH1	3:A:602:ATR:N6	2.44	0.66
1:B:352:ILE:HD11	1:B:389:PHE:CZ	2.30	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:HIS:HE1	4:A:603:MLA:O1B	1.81	0.64
1:A:317[B]:ARG:NH1	3:A:602:ATR:C6	2.61	0.63
1:B:162:LYS:HG3	5:B:845:HOH:O	2.01	0.60
1:B:210:TRP:CZ2	1:B:214:VAL:CG2	2.87	0.58
1:B:597:CYS:O	1:B:598:GLY:C	2.49	0.54
1:B:317[B]:ARG:NH2	3:B:602:ATR:N1	2.56	0.54
1:B:124:LYS:HD3	1:B:215:SER:OG	2.10	0.52
1:B:317[B]:ARG:HE	3:B:602:ATR:C2	2.24	0.51
1:B:210:TRP:CZ2	1:B:214:VAL:HG21	2.46	0.50
1:A:471:THR:HG23	5:A:886:HOH:O	2.11	0.49
1:A:359:ASP:OD1	1:A:361:GLU:HG2	2.13	0.48
1:A:162:LYS:HD2	1:A:162:LYS:C	2.39	0.47
1:A:34:LEU:CD1	1:A:88:VAL:HG11	2.44	0.47
1:A:175:LEU:HD23	1:B:175:LEU:CD2	2.43	0.47
1:B:28:CYS:CB	1:B:31:CYS:HG	2.29	0.46
1:B:162:LYS:HD2	1:B:162:LYS:C	2.42	0.45
1:B:185:LEU:H	1:B:185:LEU:HD23	1.81	0.45
1:A:108:TYR:O	1:A:250:GLY:HA2	2.17	0.45
1:B:7:THR:HG22	1:B:55:ASN:HD22	1.81	0.45
1:A:242:ASN:HD22	1:A:244:LYS:H	1.64	0.44
1:B:7:THR:CG2	1:B:55:ASN:HD22	2.29	0.44
1:B:108:TYR:O	1:B:250:GLY:HA2	2.16	0.44
1:A:318:SER:OG	3:A:602:ATR:O3P	2.26	0.44
1:A:346:LEU:O	1:A:374:TYR:HA	2.18	0.44
1:A:20:VAL:CG1	1:A:75:MET:HE3	2.47	0.44
1:A:556:LYS:HE2	1:A:556:LYS:HB2	1.85	0.44
1:B:285:PHE:CD1	1:B:286:PRO:HD2	2.54	0.43
1:B:7:THR:HG21	1:B:58:VAL:HG11	1.99	0.42
1:A:443:PRO:HG3	1:B:571:HIS:HB2	2.01	0.42
1:B:34:LEU:HD11	1:B:88:VAL:HG11	2.01	0.42
1:A:285:PHE:CD1	1:A:286:PRO:HD2	2.54	0.42
1:B:136:LEU:HD23	1:B:224:LEU:HB2	2.00	0.42
1:B:242:ASN:HD22	1:B:244:LYS:H	1.68	0.42
1:B:210:TRP:CE2	1:B:214:VAL:HG21	2.54	0.42
1:B:556:LYS:HB2	1:B:556:LYS:HE2	1.92	0.42
1:A:361:GLU:OE1	1:A:361:GLU:N	2.45	0.41
1:B:159:CYS:SG	2:B:601:FAD:C4X	3.09	0.41
1:A:561:LYS:NZ	1:A:565:ASP:OD1	2.52	0.41
1:B:346:LEU:O	1:B:374:TYR:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	594/598 (99%)	579 (98%)	15 (2%)	0	100	100
1	B	593/598 (99%)	577 (97%)	16 (3%)	0	100	100
All	All	1187/1196 (99%)	1156 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	495/496 (100%)	482 (97%)	13 (3%)	41	36
1	B	494/496 (100%)	481 (97%)	13 (3%)	41	36
All	All	989/992 (100%)	963 (97%)	26 (3%)	41	36

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	LYS
1	A	65	SER
1	A	143	PRO
1	A	159	CYS
1	A	242	ASN
1	A	320	LEU
1	A	357	PRO
1	A	379	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	399	LYS
1	A	442	THR
1	A	463	LEU
1	A	475	THR
1	A	562	GLU
1	B	80	LYS
1	B	89	LEU
1	B	103	GLU
1	B	159	CYS
1	B	242	ASN
1	B	320	LEU
1	B	379	LYS
1	B	411	LYS
1	B	442	THR
1	B	463	LEU
1	B	475	THR
1	B	476	PRO
1	B	531	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	ASN
1	A	156	ASN
1	A	173	HIS
1	A	209	ASN
1	A	225	ASN
1	A	242	ASN
1	B	52	GLN
1	B	90	ASN
1	B	93	ASN
1	B	156	ASN
1	B	173	HIS
1	B	209	ASN
1	B	225	ASN
1	B	242	ASN
1	B	363	ASN
1	B	496	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	601	-	53,58,58	0.82	3 (5%)	68,89,89	0.80	1 (1%)
4	MLA	A	603	-	6,6,6	1.48	0	7,7,7	0.76	0
2	FAD	B	601	-	53,58,58	0.74	1 (1%)	68,89,89	0.89	3 (4%)
3	ATR	B	602	-	27,33,33	0.79	0	35,52,52	1.20	4 (11%)
4	MLA	B	603	-	6,6,6	0.96	0	7,7,7	1.53	1 (14%)
3	ATR	A	602	-	27,33,33	0.82	0	35,52,52	1.21	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	601	-	-	4/30/50/50	0/6/6/6
4	MLA	A	603	-	-	0/4/4/4	-
2	FAD	B	601	-	-	3/30/50/50	0/6/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATR	B	602	-	-	4/17/37/37	0/3/3/3
4	MLA	B	603	-	-	1/4/4/4	-
3	ATR	A	602	-	-	7/17/37/37	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	FAD	C9-C8	-2.52	1.35	1.39
2	A	601	FAD	C1'-C2'	-2.47	1.49	1.52
2	B	601	FAD	C1'-C2'	-2.15	1.49	1.52
2	A	601	FAD	O2-C2	-2.09	1.20	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	ATR	PA-O3A-PB	3.12	143.53	132.83
2	A	601	FAD	O4B-C1B-C2B	-2.94	102.63	106.93
3	B	602	ATR	O3'-C3'-C4'	-2.61	103.49	111.05
3	B	602	ATR	O2'-C2'-C1'	-2.58	100.82	110.10
3	A	602	ATR	O3P-P2'-O2P	2.49	117.17	107.64
2	B	601	FAD	C5A-C6A-N6A	2.32	123.89	120.35
3	B	602	ATR	O2'-C2'-C3'	2.27	119.90	111.68
2	B	601	FAD	O4B-C1B-C2B	-2.25	103.64	106.93
2	B	601	FAD	O3B-C3B-C2B	2.24	119.06	111.82
4	B	603	MLA	O3A-C3-C2	-2.23	115.57	122.08
3	B	602	ATR	C1'-N9-C4	-2.18	122.81	126.64
3	A	602	ATR	O2P-P2'-O1P	2.17	119.17	110.68
3	A	602	ATR	O2'-P2'-O1P	-2.05	101.48	109.39

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	FAD	PA-O3P-P-O5'
3	A	602	ATR	C5'-O5'-PA-O1A
3	B	602	ATR	C2'-O2'-P2'-O1P
3	B	602	ATR	O4'-C4'-C5'-O5'
2	B	601	FAD	O4B-C4B-C5B-O5B
3	B	602	ATR	C3'-C4'-C5'-O5'
3	A	602	ATR	PB-O3A-PA-O1A
2	A	601	FAD	PA-O3P-P-O5'

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	602	ATR	PB-O3A-PA-O5'
3	B	602	ATR	C4'-C5'-O5'-PA
3	A	602	ATR	C2'-O2'-P2'-O2P
2	B	601	FAD	C3B-C4B-C5B-O5B
2	A	601	FAD	P-O3P-PA-O1A
3	A	602	ATR	C5'-O5'-PA-O3A
3	A	602	ATR	C2'-O2'-P2'-O3P
4	B	603	MLA	O1B-C1-C2-C3
2	A	601	FAD	P-O3P-PA-O2A
2	A	601	FAD	O4B-C4B-C5B-O5B
3	A	602	ATR	O4'-C4'-C5'-O5'


There are no ring outliers.


5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	603	MLA	1	0
2	B	601	FAD	1	0
3	B	602	ATR	2	0
4	B	603	MLA	1	0
3	A	602	ATR	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	592/598 (98%)	-0.03	13 (2%) 62 64	13, 30, 51, 85	4 (0%)
1	B	592/598 (98%)	0.19	11 (1%) 66 68	16, 34, 55, 93	3 (0%)
All	All	1184/1196 (98%)	0.08	24 (2%) 64 67	13, 32, 54, 93	7 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	360	THR	4.4
1	A	595	GLY	4.1
1	B	7	THR	4.0
1	A	596	CYS	3.6
1	A	243	LYS	3.5
1	A	362	ASN	3.5
1	B	210	TRP	3.4
1	A	358	VAL	3.4
1	A	597	CYS	3.2
1	A	359	ASP	3.1
1	B	53	LEU	2.7
1	B	33	LYS	2.7
1	A	594	THR	2.6
1	B	61	LYS	2.6
1	B	411	LYS	2.5
1	B	404	ALA	2.5
1	A	593	VAL	2.4
1	B	13	ARG	2.4
1	A	403	GLU	2.2
1	B	355	LEU	2.2
1	A	404	ALA	2.2
1	B	597	CYS	2.1
1	A	496	ASN	2.1
1	B	87	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ATR	A	602	31/31	0.70	0.20	39,64,171,188	0
3	ATR	B	602	31/31	0.77	0.16	44,64,142,151	0
4	MLA	A	603	7/7	0.90	0.09	35,41,45,45	0
4	MLA	B	603	7/7	0.91	0.09	31,36,40,41	0
2	FAD	B	601	53/53	0.94	0.08	28,34,40,43	0
2	FAD	A	601	53/53	0.97	0.06	23,27,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around ATR B 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.