



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 22, 2026 – 09:10 PM JST

PDB ID : 22WZ / pdb_000022wz
Title : GII.23/24/25 noroviruses recognize glycans via a conventional glycan-binding site
Authors : Li, H.B.; Cong, X.
Deposited on : 2026-01-26
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3717 (1.50-1.50)
Clashscore	180529	4048 (1.50-1.50)
Ramachandran outliers	177936	3970 (1.50-1.50)
Sidechain outliers	177891	3967 (1.50-1.50)
RSRZ outliers	164620	3718 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	
1	B	307	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5613 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	307	2428	1543	418	457	10	0	4	0
1	B	307	2419	1537	416	456	10	0	3	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	385	385	385	0	0
2	B	381	381	381	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: VP1



- Molecule 1: VP1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	108.84Å 53.49Å 109.42Å 90.00° 102.70° 90.00°	Depositor
Resolution (Å)	45.31 – 1.50 45.31 – 1.50	Depositor EDS
% Data completeness (in resolution range)	97.9 (45.31-1.50) 98.6 (45.31-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 1.49Å)	Xtrriage
Refinement program	PHENIX PHENIX	Depositor
R, R_{free}	0.184 , 0.203 0.193 , 0.212	Depositor DCC
R_{free} test set	4906 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	12.9	Xtrriage
Anisotropy	0.061	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 37.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5613	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.17	0/2507	0.35	0/3424
1	B	0.19	0/2498	0.34	0/3411
All	All	0.18	0/5005	0.35	0/6835

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2428	0	2341	20	0
1	B	2419	0	2332	13	0
2	A	385	0	0	7	1
2	B	381	0	0	4	1
All	All	5613	0	4673	33	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:ARG:NH2	2:A:402:HOH:O	2.22	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:GLN:O	1:A:240:GLN:HG3	1.93	0.68
1:A:67:THR:O	2:A:401:HOH:O	2.12	0.67
1:A:270:ARG:NH2	2:A:405:HOH:O	2.29	0.65
2:A:403:HOH:O	1:B:236:GLN:NE2	2.32	0.62
1:A:171:LEU:HG	1:A:215:ILE:HD11	1.83	0.61
1:A:208:ILE:HD13	1:A:275:THR:HG22	1.88	0.54
1:B:255:VAL:HG23	1:B:257:PRO:HD3	1.90	0.54
1:A:65:ARG:NH2	2:A:415:HOH:O	2.41	0.53
1:B:8:LEU:N	2:B:414:HOH:O	2.43	0.52
1:A:22:ILE:HD12	1:A:211:PHE:HB3	1.92	0.52
1:B:5:VAL:O	2:B:402:HOH:O	2.19	0.51
1:B:33:GLN:O	2:B:401:HOH:O	2.19	0.50
1:A:258:ASP:OD2	1:A:289:ASN:ND2	2.45	0.49
1:B:264:PHE:HB2	1:B:307:MET:HE2	1.96	0.48
1:B:264:PHE:CG	1:B:307:MET:HE2	2.49	0.48
1:B:258:ASP:OD2	1:B:289:ASN:ND2	2.47	0.47
1:A:293:ARG:NE	2:A:412:HOH:O	2.40	0.47
1:B:264:PHE:CB	1:B:307:MET:HE2	2.45	0.46
1:B:256:ASN:HB2	1:B:263:LEU:HD11	1.97	0.45
1:A:26:TYR:OH	1:A:281:ASP:OD1	2.23	0.45
1:A:240:GLN:NE2	2:A:420:HOH:O	2.49	0.45
1:B:44:LEU:HG	1:B:270:ARG:HA	1.99	0.44
1:A:66:ILE:O	1:A:161:ASN:HA	2.18	0.44
1:B:1:LYS:HE2	1:B:244:PRO:HD3	1.98	0.44
1:A:246:GLN:HB2	1:A:297:TRP:CG	2.54	0.43
1:B:256:ASN:O	1:B:260:GLY:N	2.48	0.43
1:A:25:LEU:HB2	1:A:284:ILE:HB	2.01	0.42
1:A:240:GLN:HG2	2:B:414:HOH:O	2.20	0.41
1:A:1:LYS:HA	1:A:2:PRO:HD3	1.89	0.41
1:A:141:LYS:HD3	1:A:141:LYS:HA	1.91	0.40
1:A:246:GLN:HB2	1:A:297:TRP:CD1	2.56	0.40
1:A:40:GLY:HA2	1:A:235:ILE:CD1	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:494:HOH:O	2:B:619:HOH:O[4_546]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	309/307 (101%)	303 (98%)	6 (2%)	0	100	100
1	B	308/307 (100%)	302 (98%)	6 (2%)	0	100	100
All	All	617/614 (100%)	605 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/265 (102%)	266 (99%)	3 (1%)	70	48
1	B	268/265 (101%)	268 (100%)	0	100	100
All	All	537/530 (101%)	534 (99%)	3 (1%)	84	70

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	VAL
1	A	118	GLN
1	A	286	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	174	ASN
1	B	161	ASN
1	B	207	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	307/307 (100%)	0.27	21 (6%) 25 25	6, 14, 29, 42	4 (1%)
1	B	307/307 (100%)	0.88	62 (20%) 3 3	7, 18, 39, 67	3 (0%)
All	All	614/614 (100%)	0.58	83 (13%) 8 7	6, 16, 35, 67	7 (1%)

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	255	VAL	8.5
1	B	260	GLY	7.1
1	A	285	VAL	6.8
1	B	291	TYR	6.7
1	B	9	THR	5.6
1	B	7	ILE	5.6
1	B	307	MET	5.6
1	B	257	PRO	5.5
1	A	2	PRO	5.4
1	B	262	VAL	5.4
1	A	301	PHE	5.2
1	B	263	LEU	5.2
1	B	8	LEU	5.0
1	B	259	THR	4.6
1	B	10	ILE	4.5
1	B	261	ARG	4.3
1	A	286	VAL	4.1
1	B	292	PHE	4.0
1	A	211	PHE	3.9
1	B	264	PHE	3.9
1	B	256	ASN	3.9
1	B	258	ASP	3.8
1	B	290	GLY	3.7
1	A	189	LEU	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	254	TYR	3.7
1	B	289	ASN	3.7
1	B	155	VAL	3.6
1	B	306	PRO	3.6
1	B	72	SER	3.6
1	B	304	LEU	3.6
1	B	293	ARG	3.5
1	B	1	LYS	3.5
1	A	72	SER	3.4
1	B	31	ASP	3.4
1	B	2	PRO	3.3
1	A	261	ARG	3.3
1	B	295	ASP	3.2
1	B	305	ALA	3.1
1	A	191	LEU	3.1
1	B	280	GLY	3.1
1	B	294	PHE	3.0
1	B	71	GLY	3.0
1	B	296	SER	3.0
1	A	255[A]	VAL	3.0
1	B	285	VAL	3.0
1	B	301	PHE	3.0
1	B	297	TRP	2.9
1	B	11	GLY	2.9
1	A	5	VAL	2.8
1	A	10	ILE	2.8
1	A	240	GLN	2.8
1	A	149	VAL	2.8
1	B	284	ILE	2.8
1	B	200	HIS	2.8
1	B	300	GLN	2.8
1	B	288	PRO	2.7
1	A	155	VAL	2.6
1	B	303	SER	2.6
1	B	251	LEU	2.6
1	B	287	PRO	2.6
1	B	32	ASN	2.6
1	A	299	ASN	2.5
1	B	34	ASN	2.5
1	B	278	LYS	2.5
1	B	246	GLN	2.5
1	B	74	GLN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	276	ILE	2.4
1	B	286	VAL	2.3
1	B	175	GLU	2.3
1	B	253	ARG	2.3
1	A	156[A]	ASN	2.2
1	B	282	GLY	2.2
1	A	300	GLN	2.1
1	B	211	PHE	2.1
1	A	263	LEU	2.1
1	A	203	TYR	2.1
1	B	13	MET	2.1
1	B	201	PRO	2.1
1	B	279	SER	2.1
1	B	277	ALA	2.1
1	B	252	LEU	2.0
1	B	26	TYR	2.0
1	A	175	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.